

Chulalongkorn University

## Chula Digital Collections

---

Chulalongkorn University Theses and Dissertations (Chula ETD)

---

2020

### Chemical constituents with alpha-glucosidase inhibitory activity from dendrobium braianense and dendrobium kentrophyllum

Mia Indria Permatasari

*Faculty of Pharmaceutical Sciences*

Follow this and additional works at: <https://digital.car.chula.ac.th/chulaetd>



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

---

#### Recommended Citation

Permatasari, Mia Indria, "Chemical constituents with alpha-glucosidase inhibitory activity from dendrobium braianense and dendrobium kentrophyllum" (2020). *Chulalongkorn University Theses and Dissertations (Chula ETD)*. 284.

<https://digital.car.chula.ac.th/chulaetd/284>

This Thesis is brought to you for free and open access by Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn University Theses and Dissertations (Chula ETD) by an authorized administrator of Chula Digital Collections. For more information, please contact [ChulaDC@car.chula.ac.th](mailto:ChulaDC@car.chula.ac.th).

CHEMICAL CONSTITUENTS WITH ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY FROM  
*DENDROBIUM BRAIANENSE* AND *DENDROBIUM KENTROPHYLLUM*



Miss Mia Indria Permatasari

A Thesis Submitted in Partial Fulfillment of the Requirements  
for the Degree of Master of Science in Pharmaceutical Sciences and Technology

Common Course

FACULTY OF PHARMACEUTICAL SCIENCES

Chulalongkorn University

Academic Year 2020

Copyright of Chulalongkorn University

องค์ประกอบทางเคมีที่มีฤทธิ์ยับยั้งเอนไซม์แอลฟา-กลูโคซิเดสของเอื้องคำปอกลาวและเอื้องก้างปลา  
ใหญ่



น.ส.ไมอา อินเดรีย เพอร์มาทาสารี

วิทยานิพนธ์นี้เป็นส่วนหนึ่งของการศึกษาตามหลักสูตรปริญญาวิทยาศาสตรมหาบัณฑิต  
สาขาวิชาเภสัชศาสตร์และเทคโนโลยี ไม่สังกัดภาควิชา/เทียบเท่า  
คณะเภสัชศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย  
ปีการศึกษา 2563  
ลิขสิทธิ์ของจุฬาลงกรณ์มหาวิทยาลัย

Thesis Title	CHEMICAL CONSTITUENTS WITH ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY FROM <i>DENDROBIUM BRAIANENSE</i> AND <i>DENDROBIUM KENTROPHYLLUM</i>
By	Miss Mia Indria Permatasari
Field of Study	Pharmaceutical Sciences and Technology
Thesis Advisor	Associate Professor BOONCHOO SRITULARAK, Ph.D.
Thesis Co Advisor	Professor KITTISAK LIKHITWITAYAWUID, Ph.D.

---

Accepted by the FACULTY OF PHARMACEUTICAL SCIENCES, Chulalongkorn  
University in Partial Fulfillment of the Requirement for the Master of Science

..... Dean of the FACULTY OF  
PHARMACEUTICAL SCIENCES  
(Assistant Professor RUNGPETCH SAKULBUMRUNGSI,  
Ph.D.)

#### THESIS COMMITTEE

..... Chairman  
(Assistant Professor TAKSINA CHUANASA, Ph.D.)  
..... Thesis Advisor  
(Associate Professor BOONCHOO SRITULARAK, Ph.D.)  
..... Thesis Co-Advisor  
(Professor KITTISAK LIKHITWITAYAWUID, Ph.D.)  
..... Examiner  
(Assistant Professor CHAISAK CHANSRINIYOM, Ph.D.)  
..... External Examiner  
(Duangpen Pattamadilok, Ph.D.)

ไมอา อินเดรีย เพอร์มาทาชารี : องค์ประกอบทางเคมีที่มีฤทธิ์ยับยั้งเอนไซม์แอลฟา-กลูโคซิเดสของเอื้องคำปอกลาวและเอื้องกำปลาใหญ่. (CHEMICAL CONSTITUENTS WITH ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY FROM *DENDROBIUM BRAIANENSE* AND *DENDROBIUM KENTROPHYLLUM*) อ.ที่ปรึกษาหลัก : รศ.บุญชู ศรีตุลารักษ์, อ.ที่ปรึกษาร่วม : ศ.กิตติศักดิ์ ลิขิตวิทยาวุฒิ

การศึกษาองค์ประกอบทางเคมีจากส่วนทั้งต้นของเอื้องคำปอกลาวสามารถแยกสกัดสารกลุ่มไบเบนซิลจำนวน 3 ชนิด ได้แก่ chrysotoxin moscatilin และ gigantol จากส่วนทั้งต้นของเอื้องกำปลาใหญ่พบสารกลุ่มฟลาโวนอยด์จำนวน 3 ชนิด ได้แก่ quercetin kaempferol และ rutin การศึกษาโครงสร้างของสารที่แยกได้อาศัยการวิเคราะห์คุณสมบัติทางสเปกโทรสโกปีร่วมกับการเปรียบเทียบกับข้อมูลที่มีรายงานมาก่อน เมื่อนำสารทั้งหมดที่แยกได้มาทดสอบฤทธิ์ยับยั้งเอนไซม์แอลฟา-กลูโคซิเดสพบสารที่มีฤทธิ์แรง ได้แก่ gigantol quercetin และ kaempferol เมื่อเปรียบเทียบกับ acarbose ที่เป็นสารควบคุมผลบวกร่วม สาร chrysotoxin moscatilin และ rutin พบว่าไม่มีฤทธิ์ยับยั้งเอนไซม์

จุฬาลงกรณ์มหาวิทยาลัย  
CHULALONGKORN UNIVERSITY

สาขาวิชา เกษษศาสตร์และเทคโนโลยี  
ปีการศึกษา 2563


ลายมือชื่อนิสิต .....  
ลายมือชื่อ อ.ที่ปรึกษาหลัก .....  
ลายมือชื่อ อ.ที่ปรึกษาร่วม .....

# # 6176139033 : MAJOR PHARMACEUTICAL SCIENCES AND TECHNOLOGY

KEYWORD:

Mia Indria Permatasari : CHEMICAL CONSTITUENTS WITH ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY FROM *DENDROBIUM BRAIANENSE* AND *DENDROBIUM KENTROPHYLLUM*. Advisor: Assoc. Prof. BOONCHOO SRITULARAK, Ph.D. Co-advisor: Prof. KITTISAK LIKHITWITAYAWUID, Ph.D.

To investigate the phytochemicals of the entire plants of *Dendrobium braianense* brought about in the isolation of three bibenzyls, which included chrysotoxin, moscatilin and gigantol. From the whole plant of *D. kentrophyllum* three flavonoids, namely, quercetin, kaempferol and rutin were isolated, the structures of which were decided by means of analysis of spectroscopic data to compare the values already reported. They were assessed for their alpha-glucosidase inhibitory activity. Gigantol, quercetin and kaempferol were found to exhibit strong alpha-glucosidase inhibitory activity when compared with the positive control acarbose, while chrysotoxin, moscatilin and rutin were devoid of activity.



Field of Study: Pharmaceutical Sciences  
and Technology

Student's Signature .....

Academic Year: 2020

Advisor's Signature .....

Co-advisor's Signature .....

## ACKNOWLEDGEMENTS

To begin with and first, I would like to appreciate my thesis advisor, Associate Professor Dr. Boonchoo Sritularak, and my co-advisor Professor Dr. Kittisak Likhitwitayawuid as they gave me expert guidance, immense knowledge, support and ceaseless positive thinking that lead to effective completion of this study.

I would also like to thanks the members of my thesis committee for their valuable suggestion, and express my gratitude to all teachers and staff members of the Department of Pharmacognosy and Pharmaceutical Botany, as well as the staff members of the Faculty of Pharmaceutical Sciences, Chulalongkorn University for providing all facilities.

I want to extend my sincere gratitude to the Graduate School, Chulalongkorn University for CU-ASEAN scholarship for my M.Sc. study.

Thanks to the Rector of Pharmacy the University Muhmmadiyah of Surakarta (Indonesia) and Mr Radikin (Head of Wadaslintang 2 Government Public Health) for giving me permission to pursue this M.Sc. degree at Chulalongkorn University.

I would like to appreciate all of my friends in Thailand for their warmest heart and kindest help to me during this time.

Eventually, my greatest gratitude is also expressed to parents of mine for their mortal support throughout the course of this M.Sc. degree to accomplish.

Mia Indria Permatasari

## TABLE OF CONTENTS

	Page
ABSTRACT (THAI) .....	iii
ABSTRACT (ENGLISH) .....	iv
ACKNOWLEDGEMENTS .....	v
TABLE OF CONTENTS .....	vi
LIST OF TABLES .....	1
LIST OF FIGURES .....	2
LIST OF SCHEMES .....	3
ABBREVIATIONS AND SYMBOLS .....	4
CHAPTER I INTRODUCTION .....	7
CHAPTER II LITERATURE REVIEW .....	11
1. Chemical constituents of <i>Dendrobium</i> species .....	11
2. Alpha-glucosidase inhibitory activity of <i>Dendrobium</i> species .....	77
CHAPTER III EXPERIMENTAL .....	80
1. Source of plant materials .....	80
1.1 <i>Dendrobium braianense</i> ( เอื้องคำป้อกลาว ) .....	80
1.2 <i>Dendrobium kentrophyllum</i> ( เอื้องกำป่าใหญ่ ) .....	80
2. General Techniques .....	80
2.1 Analytical thin-layer chromatography (TLC) .....	80
2.1.1 Normal phase thin-layer chromatography .....	80
2.1.2 Reverse phase thin-layer chromatography .....	81
2.2 Column chromatography (CC) .....	81




2.2.1 Vacuum liquid chromatography (VLC).....	81
2.2.2 Flash column chromatography (FCC), normal phase .....	81
2.2.3 Flash column chromatography (FCC), reverse phase .....	81
2.2.4 Gel filtration chromatography .....	82
2.2.5 Semi-preparative high pressure liquid chromatography (HPLC).....	82
2.3 Spectroscopy .....	83
2.3.1 Mass spectra.....	83
2.3.2 Proton and carbon-13 nuclear magnetic resonance ( $^1\text{H}$ and $^{13}\text{C}$ -NMR) spectra .....	83
2.4 Solvents .....	83
3. Extraction and isolation .....	83
3.1 Extraction and isolation of compounds from <i>Dendrobium braianense</i> ...	83
3.1.1 Extraction .....	83
3.1.2 Isolation.....	85
3.1.2.1 Isolation of compound DBra1 (Chrysotoxine) and compound DBra2 (Moscatilin).....	85
3.1.2.2 Isolation of compound DBra3 (Gigantol).....	85
3.2 Extraction and isolation of compounds from <i>Dendrobium kentrophyllum</i>	87
3.2.1 Extraction .....	87
3.2.2 Isolation .....	88
3.2.2.1 Isolation of compound DK 1 (Kaempferol) and compound DK 2 (Quercetin) .....	88
3.2.2.2 Isolation of compound DK3 (Rutin) .....	88
4. Physical and spectral data of isolated compounds.....	90
4.1 Compound DBra 1 (Chrysotoxine).....	90

4.2 Compound DBra 2 (Moscatalin) .....	90
4.3 Compound DBra 3 (Gigantol).....	90
4.4 Compound DK 1 (Kaempferol).....	90
4.5 Compound DK 2 (Quercetin).....	91
4.6 Compound DK 3 (Rutin).....	91
5. Assay for $\alpha$ -glucosidase inhibitory activity .....	92
5.1 Materials and instruments .....	92
5.2 Determination of $\alpha$ -glucosidase inhibitory activity .....	93
CHAPTER IV RESULTS AND DISCUSSION .....	94
1. Preliminary biological activity evaluation of <i>D. braianense</i> and <i>D. kentrophyllum</i> .....	94
2. Structure determination of isolated compounds from <i>D. braianense</i> .....	96
2.1 Identification of compound DBra1 (Chrysotoxine).....	96
2.2 Identification of compound DBra2 (Moscatalin).....	98
2.3 Identification of compound DBra3 (Gigantol).....	100
3. Structure determination of isolated compounds from <i>D. kentrophyllum</i> .....	102
3.1 Identification of compound DK1 (Kaempferol).....	102
3.2 Identification of compound DK2 (Quercetin) .....	104
3.3 Identification of compound DK3 (Rutin).....	106
4. Biological activity of isolated compounds of <i>Dendrobium braianense</i> and <i>Dendrobium kentrophyllum</i> .....	108
CHAPTER V CONCLUSION.....	109
APPENDICES.....	110
Appendix 1: The spectral data of compound DBra 1 .....	110

1.1 Mass spectrum of compound DBra 1 .....	110
1.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DBra 1 (Acetone- $d_6$ ).....	110
1.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 1 (Acetone- $d_6$ ) .....	111
1.4 HSQC spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 2.82 – 6.83 ppm, $\delta\text{C}$ 37.6 – 120.4] .....	111
1.5 HSQC spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 6.49 – 6.83 ppm, $\delta\text{C}$ 106.0 – 120.4].....	112
1.6 HMBC spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 2.82 – 6.83 ppm, $\delta\text{C}$ 37.6 – 149.3 ppm] .....	112
1.7 HMBC spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 2.82 – 6.83 ppm, $\delta\text{C}$ 106.0 – 149.3 ppm].....	113
1.8 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 2.82 – 3.77 ppm (F1), $\delta\text{H}$ 6.49 – 6.83 ppm (F2)].....	113
1.9 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 2.82 – 6.83 ppm].....	114
1.10 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 6.49 – 6.83 ppm].....	114
Appendix 2: The spectral data of compound DBra 2 .....	115
2.1 Mass spectrum of compound DBra 2 .....	115
2.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DBra 2 (Acetone- $d_6$ ).....	115
2.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 2 (Acetone- $d_6$ ) .....	116
Appendix 3: The spectral data of compound DBra 3 .....	116
3.1 Mass spectrum of compound DBra 3 .....	116
3.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DBra 3 (Acetone- $d_6$ ).....	117
3.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 3 (Acetone- $d_6$ ) .....	117

Appendix 4: The spectral data of compound DK 1.....	118
4.1. Mass spectrum of compound DK 1.....	118
4.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DK 1 (Acetone- $d_6$ ).....	118
4.3 $^{13}\text{C}$ -NMR (300 MHz) spectrum of compound DK 1 (Acetone- $d_6$ ) .....	119
4.4 HSQC NMR spectrum of compound DK 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 6.26 – 8.14 ppm, $\delta\text{C}$ 93.6 – 129.6 ppm].....	119
4.5 HMBC NMR spectrum of compound DK 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 6.26 – 12.17 ppm, $\delta\text{C}$ 93.6 – 164.1 ppm].....	120
4.6 HMBC NMR spectrum of compound DK 1 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 6.26 – 12.17 ppm, $\delta\text{C}$ 93.6 – 164.1 ppm].....	120
Appendix 5: The spectral data of compound DK 2.....	121
5.1 Mass spectrum of compound DK 2.....	121
5.2 $^1\text{H}$ -NMR (500 MHz) spectrum of compound DK 2 (Acetone- $d_6$ ).....	122
5.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DK 2 (Acetone- $d_6$ ).....	122
Appendix 6: The spectral data of compound DK 3.....	123
6.1 Mass spectrum of compound DK 3.....	123
6.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DK 3 (DMSO- $d_6$ ).....	123
6.3 $^{13}\text{C}$ -NMR (300 MHz) spectrum of compound DK 3 (DMSO- $d_6$ ).....	124
6.4 HSQC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 0.98 – 4.37 ppm, $\delta\text{C}$ 67.4 – 76.9 ppm].....	124
6.5 HSQC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 4.37 – 7.54 ppm, $\delta\text{C}$ 94.0 – 122.0 ppm].....	125
6.6 HMBC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta\text{H}$ 0.98 – 12.59 ppm, $\delta\text{C}$ 18.2 – 177.8 ppm].....	126

6.7 HMBC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta_H$ 3.22 – 12.59 ppm, $\delta_C$ 94.0 – 177.8 ppm].....	126
6.8 HMBC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta_H$ 0.98 – 4.37 ppm, $\delta_C$ 67.4 – 76.9 ppm].....	127
6.9 HMBC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta_H$ 0.98 – 7.54 ppm].....	127
6.10 HMBC spectrum of compound DK 3 (Acetone- $d_6$ ) expansion [ $\delta_H$ 3.03 – 5.33 ppm].....	128
REFERENCES .....	129
VITA.....	144



## LIST OF TABLES

Table 1 Stilbenoids from genus <i>Dendrobium</i> .....	14
Table 2 Flavonoids from genus <i>Dendrobium</i> .....	44
Table 3 Terpenoids from <i>Dendrobium</i> .....	49
Table 4 Miscellaneous compounds from <i>Dendrobium</i> .....	57
Table 5 Examples of $\alpha$ -glucosidase inhibitors from <i>Dendrobium</i> spp. ....	78
Table 6 $\alpha$ -glucosidase inhibitory activity of crude extracts from <i>D. braianense</i> and <i>D. Kentrophyllum</i> .....	95
Table 7 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of compound DBra1 in acetone- $d_6$ and $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) of chrysotoxine in $\text{CDCl}_3$ .....	97
Table 8 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of compound DBra2 in acetone- $d_6$ and $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) of moscatilin in acetone- $d_6$ .....	99
Table 9 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of compound DBra3 in acetone- $d_6$ and $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) of gigantol in acetone- $d_6$ .....	101
Table 10 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of DK1 in acetone- $d_6$ and $^1\text{H}$ NMR (600 MHz) and $^{13}\text{C}$ NMR (150 MHz) of kaempferol in $\text{DMSO}-d_6$ .....	103
Table 11 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of compound DK2 in acetone- $d_6$ and $^1\text{H}$ NMR (600 MHz) and $^{13}\text{C}$ NMR (150 MHz) of quercetin in $\text{DMSO}-d_6$ .....	105
Table 12 $^1\text{H}$ NMR (300 MHz) and $^{13}\text{C}$ NMR (75 MHz) spectral data of compound DK3 in acetone- $d_6$ and $^1\text{H}$ NMR (400 MHz) and $^{13}\text{C}$ NMR (100 MHz) of rutin in $\text{DMSO}-d_6$ .....	107
Table 13 $\alpha$ -Glucosidase inhibitory activity of <i>Dendrobium braianense</i> and <i>Dendrobium kentrophyllum</i> .....	108

## LIST OF FIGURES

Figure 1. <i>Dendrobium braianense</i> .....	9
Figure 2. <i>Dendrobium kentrophyllum</i> .....	9
Figure 3. Structures of stilbenoids from <i>Dendrobium</i> .....	27
Figure 4. Structures of flavonoids from <i>Dendrobium</i> .....	46
Figure 5. Structures of terpenoids from <i>Dendrobium</i> .....	52
Figure 6. Structures of miscellaneous compounds from <i>Dendrobium</i> .....	66
Figure 7. Structures of $\alpha$ -glucosidase inhibitors from <i>Dendrobium</i> spp.....	79
Figure 8. Chrysotoxine .....	96
Figure 9. Moscatilin .....	98
Figure 10. Gigantol .....	100
Figure 12. Kaempferol .....	102
Figure 13. Quercetin .....	104
Figure 14. Rutin .....	106

## LIST OF SCHEMES

Scheme 1 Extraction steps for <i>Dendrobium braianense</i> .....	84
Scheme 2 Isolation of compounds from Ethyl acetate extract of <i>Dendrobium braianense</i> .....	86
Scheme 3 Extraction steps for <i>Dendrobium kentrophyllum</i> .....	87
Scheme 4 Isolation of compounds from ethyl acetate extract of <i>Dendrobium kentrophyllum</i> .....	89
Scheme 5 Isolation of compounds from fraction H of <i>Dendrobium kentrophyllum</i> .....	89





## ABBREVIATIONS AND SYMBOLS

Acetone- $d_6$	=	Deuterated acetone
APCI-MS	=	Atmospheric Pressure Chemical Ionization Mass Spectrometry
br s	=	Broad singlet (for NMR spectra)
°C	=	Degree celsius
CC	=	Column chromatography
$CDCl_3$	=	Deuterated chloroform
$CH_2Cl_2$	=	Dichloromethane
cm	=	Centimeter
$^{13}C$ -NMR	=	Carbon-13 Nuclear Magnetic Resonance
1-D NMR	=	One-dimensional Nuclear Magnetic Resonance
2-D NMR	=	Two-dimensional Nuclear Magnetic Resonance
$d$	=	Doublet (for NMR spectra)
$dd$	=	Doublet of doublets (for NMR spectra)
$\delta$	=	Chemical shift
DMSO- $d_6$	=	Deuterated dimethylsulfoxide
EtOAc	=	Ethyl acetate
FCC	=	Flash Column Chromatography
g	=	Gram
Gal	=	Galactose
GF	=	Gel Filtration
Glc	=	Glucose
HMBC	=	$^1H$ -detected Heteronuclear Multiple Bond Correlation
HR-ESI-MS	=	High Resolution Electrospray Ionization Mass Spectrometry

$^1\text{H-NMR}$	=	Proton Nuclear Magnetic Resonance
HSQC	=	$^1\text{H}$ -detected Heteronuclear Single Quantum Coherence
Hz	=	Hertz
$\text{IC}_{50}$	=	Concentration exhibiting 50% inhibition
IR	=	Infrared
$J$	=	Coupling constant
Kg	=	Kilogram
L	=	Liter
$[\text{M}+\text{H}]^+$	=	Protonated molecular ion
$[\text{M}+\text{Na}]^+$	=	Sodium-adduct molecular ion
m	=	Multiplet (for NMR spectra)
MeOH	=	Methanol
mg	=	Milligram
$\mu\text{g}$	=	Microgram
min	=	Minute
ml	=	Milliliter
$\mu\text{l}$	=	Microliter
$\mu\text{M}$	=	Micromolar
mm	=	Millimeter
mM	=	Millimolar
MS	=	Mass spectrum
MW	=	Molecular weight
$m/z$	=	Mass to charge ratio
NA	=	Not applicable
nm	=	Nanometer
nM	=	Nanomolar
NMR	=	Nuclear Magnetic Resonance

NOESY	=	Nuclear Overhauser Effect Spectroscopy
OEt	=	Ethoxy group
OMe	=	Methoxy group
Rha	=	Rhamnose
<i>s</i>	=	Singlet (for NMR spectra)
<i>t</i>	=	Triplet (for NMR spectra)
TLC	=	Thin Layer Chromatography
UV-VIS	=	Ultraviolet and Visible spectrophotometry
VLC	=	Vacuum Liquid Column Chromatography
Xyl	=	Xylose



## CHAPTER I

### INTRODUCTION

Diabetes mellitus (DM) is characterized by high blood sugar and metabolic disorders of fat, carbohydrates as well as protein due to absolute or relative insulin deficiency. This disease has several symptoms including increased appetite, thirst, and frequency of urination (Fatimah, 2015). There are several complications of DM such as retinopathy, neuropathy, cardiovascular disease, stroke, and chronic kidney disease (Ahmad *et al.*, 2011).

Diabetes Mellitus type 2 (DM II) stems from insulin resistance or abnormal insulin secretion. Pancreatic beta cells still produce insulin, therefore DM II is regarded as non-insulin-dependent diabetes mellitus (NIDDM) (Fatimah, 2015). DM II is more common in adulthood and elderly people. The exact causes of DM are unknown, but they are related to lifestyle factors and genetics such as obesity, lack of physical activity, poor diet, and stress (Patel *et al.*, 2012).

Normal insulin has the work as the main regulator of fat, carbohydrate, and protein metabolism. First, it inhibits lipolysis of fat in adipose tissue which inhibits the release of fatty acid release in adipose tissue. Second, prevent gluconeogenesis in the liver which reducing the liver's glucose output. Third, it makes the glucose transporter type 4 (GLUT-4) transport the glucose into the muscle cells, which makes skeletal muscle increase the glucose uptake (Sears & Perry, 2015).

Hyperglycemia and more circulating fatty acids in DM II result from many factors, for instance, adipokines as well as inflammation of the diabetes genes. These factors affect how insulin is secreted in the pancreas and works. The reduced insulin secretion reduces the insulin signal in the target tissue. Insulin resistance has impacts on the action of insulin in every major target tissue, causing hyperglycemia and an

increase in circulating fatty acids. Insulin secretion and resistance were affected by the increase of glucose and fatty acids levels in the blood (Stumvoll *et al.*, 2005).

There are several ways to treat and prevent diabetes and its secondary complications. One of these is herbal medicine. There are different mechanisms of action for different antidiabetic drugs in the treatment of high glucose levels. The initial increase in insulin secretion from sulphonylureas and meglitinides. Decreased gastrointestinal glucose absorption (GIT) is a mechanism of  $\alpha$ -glucosidase inhibitors (AGIs) and dipeptide peptidase-4 inhibitors. Reduction in glucagon secretion from amylin analog  $\alpha$ -glucosidase inhibitors and dipeptide peptidase-4 inhibitors (Choudhury *et al.*, 2018).

*Dendrobium* plants have been reported to produce various classes of secondary metabolites, like phenanthrenes, bibenzyls, alkaloids, fluorenones as well as flavonoids, in which several compounds were found to possess antitumor, anti-inflammatory, antimutagenic, and  $\alpha$ -glucosidase inhibitory activities (Na Ranong, *et al.*, 2019).

*Dendrobium braianense* Gagnep. (เอื้องคำป้อกลาว), is distributed in South-Central China, Laos, Thailand, and Vietnam (Plants of the world online, 2017). The leaves of *D. braianense* gagnep have medium-sized, warm to cool growing epiphyte with pseudobulbous, slightly clavate stems carrying variably deciduous leaves that blooms in the winter and spring. The structural compound of the *D. braianense* gagnep flower has 1.6 inches or 4 cm (The Plant List, 2018).



**Figure 1.** *Dendrobium braianense*

*Dendrobium kentrophyllum* Hook. f. (เอื้องกำพร้าวัวใหญ่) stem is 10-30 cm long, erect or ascending; leaves, narrow, slightly curved outwards. Its flower has pale greenish-yellow or whitish-yellow petals. It flowering in October – November. Distribution in India, Borneo, Laos, Malaysia, Sumatra, and Thailand (Khurajam & Roy, 2016).



**Figure 2.** *Dendrobium kentrophyllum*

Prior to this report, there were no records on phytochemical constituents of *Dendrobium braianense* and *D. kentrophyllum*. Both plant crude extracts were

prepared through maceration of the dried powdered whole plants and methanol (MeOH). After being evaluated with  $\alpha$ -glucosidase inhibitory activity, MeOH extracts of both plants at concentration of 100  $\mu\text{g/mL}$  showed inhibition around 65% for *D. braianense* and around 80% for *D. kentrophyllum*. Isolation continues with liquid-liquid extraction of MeOH extracts with ethyl acetate (EtOAc), butanol (BuOH) and water ( $\text{H}_2\text{O}$ ) to give EtOAc, BuOH and water extracts respectively. Both EtOAc and BuOH extracts then were tested with  $\alpha$ -glucosidase inhibitory activity assay. The result showed that only EtOAc extracts of both plants have strong inhibitory effect with inhibition above 80% at the concentration of 100  $\mu\text{g/mL}$  for *D. braianense* and more than 90% for *D. kentrophyllum*. Therefore the EtOAc extracts from *D. braianense* and *D. kentrophyllum* were selected for further chemical investigation. Regarding the biological activity of *D. braianense* and *D. kentrophyllum*, so far, there have been no previous reports.

The information of phytochemical constituents from *Dendrobium braianense* and *Dendrobium kentrophyllum* and their  $\alpha$ -glucosidase inhibitory effect should be useful for the development of new anti-diabetic drugs from natural sources. To achieve this purpose, the objectives of this project are:

1. To isolate and purify the chemical constituents from *Dendrobium braianense* and *D. kentrophyllum* and analyze the chemical structure of each compound.
2. To investigate  $\alpha$ -glucosidase inhibitory activity of isolated compounds.

## CHAPTER II

### LITERATURE REVIEW

#### 1. Chemical constituents of *Dendrobium* species

Like other plants, orchids produce many phytochemicals. Chemical properties and biological activity have been studied in several of them. Dendrobine was the first alkaloid isolated from the orchid *Dendrobium nobile* and has been used in the Chinese traditional medicine "Shi-Hu" (Hossain, 2011). In Orchidaceae, *Dendrobium* is a large genus. In Chinese traditional medicine, dried *Dendrobium* spp. stems have been used to treat stomach, diabetes, kidney, and lung diseases (Xu *et al.*, 2017). In Thailand, some of these orchids are also used in traditional medicine (Chuakul, 2002). According to reports, several types of secondary metabolites are already isolated from *Dendrobium* spp. such as, bibenzyls, phenanthrenes, alkaloids, and flavonoids which have certain biological activities (Wang *et al.*, 2009).

Shi-Hu is deemed to be an important herb for replenishing yin (coldness, moistness, and so on) and is widely used in Chinese medicine. They are considered effective in some diseases or syndromes related to the deficiency of yin in the kidney, lung, and stomach—like thirst, fever, red tongue, faucitis, atrophic gastritis as well as diabetes (Bulpitt *et al.*, 2007).

There is some traditional belief that suggests the use of several orchids in treating diabetes. Some of these orchids have also been validated by several scholars throughout the world experimentally, while some are still under study. The examples of the orchids that have a pathological condition to treat diabetes are whole *Dendrobium aphyllum* plants (Roxb.), infusion of leaves from *Dendrobium auranticum*, leaves from *Dendrobium candidum* wall ex Lindl., whole plants of



*Dendrobium chrysotoxum*, and whole plants of *Dendrobium officinale* kimura et migo (Mukherjee & Jagtap, 2020).

Table 1 and Figure 3 show compounds belonging to stilbenoids which can be divided into stilbenes, bibenzyls, bisbibenzyls, and phenanthrenes derivatives. The main structural skeleton of this group composes of 2 aromatic rings linked by an ethylene bridge. Stilbenoids are constructed from *trans*-cinnamic acid and three malonyl-CoA units. In phenylpropanoid biosynthesis, L-phenylalanine is deaminated by phenylalanine ammonia-lyase (PAL) to give *trans*-cinnamic which is then hydroxylated by cinnamate-4-hydroxylase (C4H) to yield 4-coumaric acid. This compound is then activated by 4-coumaroyl CoA ligase (4CL) to form 4-coumaroyl CoA. Subsequently, three acetate extender units from malonyl-CoA are added to 4-coumaroyl CoA by stilbene synthase (STS) to make tetraketide intermediate which consequently folds and cyclizes to shape as a chalcone or a stilbene structure (Tsopmo *et al.*, 2013).

Bibenzyl is a type of active compound abundant in *Dendrobium* (Gong *et al.*, 2013). The Bibenzyls constitute small-molecular compounds of natural origin and are widely source. In previous studies, the bibenzyls were shown to have good anti-tumor, anti-diabetes and its complications, and neuroprotective effects. One of their primary sources is the *Dendrobium* genus. Currently, 89 bibenzyl derivatives, for example, moscatilin (**1**), gigantol (**2**), and chrysotoxine (**3**), have been found in 46 species of *Dendrobium*. Moscatilin (**1**) has anti-tumor, neuroprotection as well as improvement of retinal vascular diseases. Gigantol (**2**) has anti-tumor, anti-diabetes as well as its complications, antioxidant, anti-inflammatory, antibacterial, antiviral, and antimalarial effects. Chrysotoxine (**3**) exhibited anti-lung cancer activity (He *et al.*, 2020).

Flavonoids, one of the largest groups of low molecular weight secondary metabolites presenting common benzo- $\gamma$ -pyrone structure, are naturally occurring

phenolic compounds. Due to the presence of aromatic hydroxyl groups, flavonoids have strong antioxidant properties. Many of the biological actions of flavonoids have been attributed to their powerful antioxidant properties. Since the beginning of the 1980s, the beneficial effects of flavonoids have been studied in DM. Flavonoids with antioxidant and free radical scavenging properties, in particular, prevent autopoly (ADP-ribosyl)-ation of poly synthetase polymerase and by stabilizing Reg gene transcriptional complex, result in the regeneration of  $\beta$ -cells and protect pancreatic islets. Recently, many biologically active flavonoids for treating DM and its complications have been found (Kumar, S., & Pandey, A. K., 2013).

Alkaloids is a very diverse group of compounds related only to the occurrence of a nitrogen atom in a heterocyclic ring. Plants are estimated to produce almost 12,000 different alkaloids, organized into groups based on their carbon skeletal structures (Ziegler, J., & Facchini, P. J., 2008).



**Table 1.** Stilbenoids from the genus *Dendrobium*

Stilbenoids	Plant	Plant part	Reference
Aloifol I [1]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. williamsonii</i>	Whole plant	(Yang <i>et al.</i> , 2018)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Amoenylin [2]	<i>D. amoenum</i>	Whole plant	(Majumder <i>et al.</i> , 1999)
	<i>D. williamsonii</i>	Whole plant	(Yang <i>et al.</i> , 2018)
Batatasin [3]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)
Batatasin III [4]	<i>D. aphyllum</i>	Whole plant	(Chen <i>et al.</i> , 2008a)
		Stem	(Yang <i>et al.</i> , 2015)
	<i>D. cariniferum</i>	Stem	(Chen <i>et al.</i> , 2008b)
	<i>D. rotundatum</i>	Whole plant	(Majumder & Pal, 1992)
	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)
	<i>D. draconis</i>	Stem	(Sritularak, Anuwat, <i>et al.</i> , 2011)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. gratiosissimum</i>	Stem	(C. F. Zhang <i>et al.</i> , 2008)
	<i>D. loddigesii</i>	Stem	(Ito <i>et al.</i> , 2010)
	<i>D. loddigesii</i>	Stem	(Ma <i>et al.</i> , 2019)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Brittonin A [5]	<i>D. secundum</i>	Stem	(Sritularak, Duangrak, <i>et al.</i> , 2011)
Chrysotobibenzyl [6]	<i>D. aurantiacum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	var. <i>denneanum</i>		
	<i>D. capillipes</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)

Stilbenoids	Plant	Plant part	Reference
	<i>D. scabriliguae</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Chrysotoxine [7]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
Crepidatin [8]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Whole plant	(Liu <i>et al.</i> , 2009)
	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
	<i>D. crepidatum</i>	Whole plant	(Majumder & Chatterjee, 1989)
	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)
Cumulatin [9]	<i>D. cumulatum</i>	Whole plant	(Majumder & Pal, 1993)
4,5-dihydroxy-3,3'-dimethoxybibenzyl [10]	<i>D. nobile</i>	Stem	(H. Wang <i>et al.</i> , 1985; Ye & Zhao, 2002)
3,3'-Dihydroxy-4,5-dimethoxybibenzyl [11]	<i>D. williamsonii</i>	Whole plant	(Rungwichaniwat <i>et al.</i> , 2014)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
3,4'-Dihydroxy-5-methoxybibenzyl [12]	<i>D. amoenum</i>	Whole plant	(Majumder <i>et al.</i> , 1999)
3,4'-Dihydroxy-5,5'-dimethoxydihydrostilbene [13]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
4,5-Dihydroxy-3,3'-dimethoxybibenzyl [14]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)
Erianin [15]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Gigantol [16]	<i>D. aphyllum</i>	Whole plant	(Y. Chen <i>et al.</i> , 2008)
	<i>D. aphyllum</i>	Stem	(D. Yang <i>et al.</i> , 2015)
	<i>D. aurantiacum</i> var. <i>denneanum</i>	Whole plant	(Liu <i>et al.</i> , 2009)
	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)

Stilbenoids	Plant	Plant part	Reference
	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. devonianum</i>	Whole plant	(Sun <i>et al.</i> , 2014)
	<i>D. draconis</i>	Stem	(Sritularak, Anuwat, <i>et al.</i> , 2011)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. gratiosissimum</i>	Stem	(C. F. Zhang <i>et al.</i> , 2008)
	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2007)
	<i>D. officinale</i>	Stem	(G. Y. Zhao <i>et al.</i> , 2018)
	<i>D. polyanthum</i>	Stem	(Hu <i>et al.</i> , 2009)
	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
	<i>D. loddigesii</i>	Stem	(Ma <i>et al.</i> , 2019)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
4-Hydroxy-3,5,3'-trimethoxybibenzyl [17]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)
5-Hydroxy-3,4,3',4',5'-pentamethoxybibenzyl [18]	<i>D. secundum</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Isoamoenylin [19]	<i>D. amoenum</i>	Whole plant	(Majumder <i>et al.</i> , 1999)
Moniliformine [20]	<i>D. williamsonii</i>	Whole plant	(Yang <i>et al.</i> , 2018)
Moscatilin [21]	<i>D. amoenum</i>	Whole plant	(Majumder <i>et al.</i> , 1999)
3,3',4-Trihydroxybibenzyl [22]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
3,3',5-Trihydroxybibenzyl [23]	<i>D. cariniferum</i>	Whole plant	(Liu <i>et al.</i> , 2009a)
	<i>D. loddigesii</i>	Stem	(Ma <i>et al.</i> , 2019b)
3,5,4'-Trihydroxybibenzyl [24]	<i>D. gratiosissimum</i>	Stem	(Zhang <i>et al.</i> , 2008a)

Stilbenoids	Plant	Plant part	Reference
4,5,4'-Trihydroxy-3,3'-dimethoxybibenzyl [25]	<i>D. secundum</i>	Stem	(Sritularak, Duangrak and Likhitwitayawuid, 2011)
	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
	<i>D. parishii</i>	Whole plant	(Kongkatitham <i>et al.</i> , 2018)
	<i>D. loddigesii</i>	Stem	(Ma <i>et al.</i> , 2019b)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
Tristin [26]	<i>D. aphyllum</i>	Stem	(Yang <i>et al.</i> , 2015)
	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
	<i>D. densiflorum</i>	Stem	(Fan <i>et al.</i> , 2001)
	<i>D. gratiosissimum</i>	Stem	(Zhang <i>et al.</i> , 2008a)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. officinale</i>	Stem	(Zhao <i>et al.</i> , 2018)
	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
	<i>D. loddigesii</i>	Stem	(Ma <i>et al.</i> , 2019b)
Dendromonilaside E [27]	<i>D. moniliforme</i>	Stem	(Zhao <i>et al.</i> , 2003)
5,4'-Dihydroxy-3,4,3'-trimethoxybibenzyl [28]	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
4,3',4'-trihydroxy-3,5-dimethoxybibenzyl [29]	<i>D. parishii</i>	Whole plant	(Kongkatitham <i>et al.</i> , 2018)
Dendrophenol [30]	<i>D. candidum</i>	Stem	(Li <i>et al.</i> , 2008)
3,4-Dihydroxy-5,4'-dimethoxybibenzyl [31]	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
	<i>D. signatum</i>	Whole plant	(Mittraphab <i>et al.</i> , 2016)
	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
4,4'-Dihydroxy-3,5-dimethoxybibenzyl [32]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2008)
	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Loddigesiinol C [33]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
3-O-Methylgigantol [34]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2008)
	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)

Stilbenoids	Plant	Plant part	Reference
Dendrocandin A [35]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2008)
Dendrocandin B [36]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2008)
	<i>D. signatum</i>	Whole plant	(Mittraphab <i>et al.</i> , 2016)
Dendrocandin C [37]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2009a)
Dendrocandin D [38]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2009a)
Dendrocandin E [39]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2009a)
	<i>D. parishii</i>	Whole plant	(Kongkatitham <i>et al.</i> , 2018)
Dendrocandin F [40]	<i>D. candidum</i>	Stem	(Yan Li <i>et al.</i> , 2009b)
Dendrocandin G [41]	<i>D. candidum</i>	Stem	(Yan Li <i>et al.</i> , 2009b)
Dendrocandin H [42]	<i>D. candidum</i>	Stem	(Yan Li <i>et al.</i> , 2009b)
Dendrosinen A [43]	<i>D. sinense</i>	Whole plant	(X. J. Chen <i>et al.</i> , 2014)
Dendrosinen B [44]	<i>D. sinense</i>	Whole plant	(X.-J. Chen <i>et al.</i> , 2014)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Dendrosinen C [45]	<i>D. sinense</i>	Whole plant	(Chen <i>et al.</i> , 2014)
Dendroscabrol B [46]	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
Dendrosinen D [47]	<i>D. sinense</i>	Whole plant	(X. J. Chen <i>et al.</i> , 2014)
Dendrocandin I [48]	<i>D. candidum</i>	Stem	(Y. Li <i>et al.</i> , 2008)
	<i>D. signatum</i>	Whole plant	(Mittraphab <i>et al.</i> , 2016)
Densiflorol A [49]	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)
Longicornuol A [50]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Trigonopol A [51]	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
Trigonopol B [52]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
	<i>D. aphyllum</i>	Stem	(D. Yang <i>et al.</i> , 2015)
Crepidatuol A [53]	<i>D. crepidatum</i>	Stem	(C. B. Li <i>et al.</i> , 2013)
Crepidatuol B [54]	<i>D. crepidatum</i>	Stem	(C. B. Li <i>et al.</i> , 2013)
Loddigesinol D [55]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
Dencryol A [56]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Dencryol B [57]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Dengraol A [58]	<i>D. gratiosissimum</i>	Stem	(C. F. Zhang <i>et al.</i> , 2008)
Dengraol B [59]	<i>D. gratiosissimum</i>	Stem	(C. F. Zhang <i>et al.</i> , 2008)

Stilbenoids	Plant	Plant part	Reference
4-[2-(3-Hydroxyphenol)-1-methoxyethyl]-2,6-dimethoxyphenol [60]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Nobilin A [61]	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2006)
Nobilin B [62]	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2006)
Nobilin C [63]	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2006)
Nobilin D [64]	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2007)
Nobilin E [65]	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2007)
Dendrofalconerol A [66]	<i>D. falconeri</i>	Stem	(Sritularak & Likhitwitayawuid, 2009)
	<i>D. signatum</i>	Whole plant	(Mittraphab <i>et al.</i> , 2016)
	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
Dendrofalconerol B [67]	<i>D. falconeri</i>	Stem	(Sritularak & Likhitwitayawuid, 2009)
Dendrosignatol [68]	<i>D. signatum</i>	Whole plant	(Mittraphab <i>et al.</i> , 2016)
2,2'-Dihydroxy-3,3',4,4',7,7'-hexamethoxy-9,9',10,10'-tetrahydro-1,1'-biphenanthrene [69]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
2,2'-Dimethoxy-4,4',7,7'-tetrahydroxy-9,9',10,10'-tetrahydro-1,1'-biphenanthrene [70]	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)
Flavanthrin [71]	<i>D. aphyllum</i>	Whole plant	(Y. Chen <i>et al.</i> , 2008)
Phoyunnanin C [72]	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
Phoyunnanin E [73]	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
Amoenumin [74]	<i>D. amoenum</i>	Whole plant	(Veerraju <i>et al.</i> , 1989)
Crystalltone [75]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Dendropalpebrone [76]	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
Chrysotoxol A [77]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Chrysotoxol B [78]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)



Stilbenoids	Plant	Plant part	Reference
Confusarin [79]	<i>D. chrysseum</i>	Stem	(Ma <i>et al.</i> , 1998)
	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2008)
	<i>D. officinale</i>	Stem	(G. Y. Zhao <i>et al.</i> , 2018)
2,6-Dihydroxy-1,5,7-trimethoxyphenanthrene [80]	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
Dendrochrysanene [81]	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
Bulbophyllanthrin [82]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
5-Hydroxy-2,4-dimethoxyphenanthrene [83]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
3-Hydroxy-2,4,7-trimethoxyphenanthrene [84]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
Cypripedin [85]	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
Densiflorol B [86]	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
Denbinobin [87]	<i>D. moniliforme</i>	Stem	(Lin <i>et al.</i> , 2001)
	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
Fimbriatone [88]	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2008)
	<i>D. pulchellum</i>	Stem	(Chanvorachote <i>et al.</i> , 2013)
Dendroscabrol A [89]	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
Loddigesiinol B [90]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
Dendronone [91]	<i>D. cariniferum</i>	Whole plant	(Y. Chen <i>et al.</i> , 2008)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Ephemeranthoquinone [92]	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)

Stilbenoids	Plant	Plant part	Reference
5-Methoxy-7-hydroxy-9,10-dihydro-1,4-phenanthrenequinone [93]	<i>D. draconis</i>	Stem	(Sritularak, Anuwat, <i>et al.</i> , 2011)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
Moniliformin [94]	<i>D. moniliforme</i>	Stem	(Lin <i>et al.</i> , 2001)
Moscatin [95]	<i>D. aphyllum</i>	Whole plant	(Y. Chen <i>et al.</i> , 2008)
	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)
	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. polyanthum</i>	Stem	(Hu <i>et al.</i> , 2009)
	<i>D. rotundatum</i>	Whole plant	(Majumder & Pal, 1992)
Coelonin [96]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
	<i>D. aphyllum</i>	Whole plant	(Y. Chen <i>et al.</i> , 2008)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
9,10-Dihydromoscatin [97]	<i>D. polyanthum</i>	Stem	(Hu <i>et al.</i> , 2009)
9,10-Dihydrophenanthrene-2,4,7-triol [98]	<i>D. polyanthum</i>	Stem	(Hu <i>et al.</i> , 2009)
4,5-Dihydroxy-2,3-dimethoxy-9,10-dihydrophenanthrene [99]	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
	<i>D. sinense</i>	Whole plant	(Chen <i>et al.</i> , 2014)
4,5-Dihydroxy-2,6-dimethoxy-9,10-dihydrophenanthrene [100]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
4,5-Dihydroxy-3,7-dimethoxy-9,10-dihydrophenanthrene [101]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)


Stilbenoids	Plant	Plant part	Reference
4,5-Dihydroxy-2-methoxy-9,10-dihydrophenanthrene [102]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
Lusianthridin [103]	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010; H. K. Yang <i>et al.</i> , 2007)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
2,7-Dihydroxy-3,4,6-trimethoxy-9,10-dihydrophenanthrene [104]	<i>D. rotundatum</i>	Whole plant	(Majumder & Pal, 1992)
2,8-Dihydroxy-3,4,7-trimethoxy-9,10-dihydrophenanthrene [105]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
4,7-Dihydroxy-2,3,6-trimethoxy-9,10-dihydrophenanthrene [106]	<i>D. sinense</i>	Whole plant	(X. J. Chen <i>et al.</i> , 2013)
Ephemeranthol A [107]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010; H. K. Yang <i>et al.</i> , 2007)
	<i>D. officinale</i>	Stem	(G. Y. Zhao <i>et al.</i> , 2018)
	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Ephemeranthol C [108]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
Erianthridin [109]	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010; H. K. Yang <i>et al.</i> , 2007)

Stilbenoids	Plant	Plant part	Reference
	<i>D. plicatile</i>	Stem	(Yamaki & Honda, 1996)
Flavanthridin [110]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
Hircinol [111]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
	<i>D. aphyllum</i>	Stem	(D. Yang <i>et al.</i> , 2015)
	<i>D. draconis</i>	Stem	(Sritularak, Anuwat, <i>et al.</i> , 2011)
	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
3-Hydroxy-2,4,7-trimethoxy-9,10-dihydrophenanthrene [112]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
7-hydroxy-2,3,4-trimethoxy-9,10-dihydrophenanthrene [113]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
Dendroinfundin A [114]	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
Dendroinfundin B [115]	<i>D. infundibulum</i>	Whole plant	(Na Ranong <i>et al.</i> , 2019)
3,4-Dimethoxy-1-(methoxymethyl)-9,10-dihydrophenanthrene-2,7-diol [116]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
2-Hydroxy-4,7-dimethoxy-9,10-dihydrophenanthrene [117]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
7-Methoxy-9,10-dihydrophenanthrene-2,4,5-triol [118]	<i>D. draconis</i>	Stem	(Sritularak, Anuwat, <i>et al.</i> , 2011)
2,5,7-Trihydroxy-4-methoxy-9,10-dihydrophenanthrene [119]	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
Plicatol C [120]	<i>D. plicatile</i>	Stem	(Honda & Yamaki, 2000)

Stilbenoids	Plant	Plant part	Reference
Rotundatin [121]	<i>D. rotundatum</i>	Whole plant	(Majumder & Pal, 1992)
2,5-Dihydroxy-3,4-dimethoxyphenanthrene [122]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
2,5-Dihydroxy-4,9-dimethoxyphenanthrene [123]	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2008)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
2,8-Dihydroxy-3,4,7-trimethoxyphenanthrene [124]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
Epheranthol B [125]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Fimbriol B [126]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007);
Flavanthrinin [127]	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
	<i>D. nobile</i>	Stem	(X. Zhang, J. K. Xu, <i>et al.</i> , 2008)
	<i>D. venustum</i>	Whole plant	(Sukphan <i>et al.</i> , 2014)
	<i>D. parishii</i>	Whole plant	(Kongkatitham <i>et al.</i> , 2018a)
Loddigesiinol A [128]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
Nudol [129]	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
	<i>D. rotundatum</i>	Whole plant	(Majumder & Pal, 1992)
Plicatol A [130]	<i>D. nobile</i>	Stem	(H. K. Yang <i>et al.</i> , 2007)
	<i>D. plicatile</i>	Stem	(Honda & Yamaki, 2000)
Plicatol B [131]	<i>D. plicatile</i>	Stem	(Honda & Yamaki, 2000)
2,3,5-Trihydroxy-4,9-dimethoxyphenanthrene [132]	<i>D. nobile</i>	Stem	(Yang, Sung and Kim., 2007)
3,4,8-Trimethoxyphenanthrene-2,5-diol [133]	<i>D. nobile</i>	Stem	(Hwang <i>et al.</i> , 2010)
Aphyllone A [134]	<i>D. aphyllum</i>	Stem	(Yang <i>et al.</i> , 2015)

Stilbenoids	Plant	Plant part	Reference
(S)-2,4,5,9-Tetrahydroxy-9,10-dihydro phenanthrene [135]	<i>D. fimbriatum</i>	Stem	(Xu et al., 2014)
1,5,7-Trimethoxy phenanthren-2-ol [136]	<i>D. nobile</i>	Stem	(Kim et al., 2015)
1,5-Dihydroxy-3,4,7-trimethoxy-9,10-dihydrophenanthrene [137]	<i>D. moniliforme</i>	Whole plant	(Zhao et al., 2016)
2,5,9S-Trihydroxy-9,10-dihydro phenanthrene-4-O- $\beta$ -D-glucopyranoside [138]	<i>D. primulinum</i>	Whole plant	(Ye et al., 2016)
Loddigesiinol G [139]	<i>D. loddigesii</i>	Stem	(Lu et al., 2014)
Loddigesiinol H [140]	<i>D. loddigesii</i>	Stem	(Lu et al., 2014)
Loddigesiinol I [141]	<i>D. loddigesii</i>	Stem	(Lu et al., 2014)
Loddigesiinol J [142]	<i>D. loddigesii</i>	Stem	(Lu et al., 2014)
Dendrowillol A [143]	<i>D. williamsonii</i>	Whole plant	(M. Yang et al., 2018)
Dendrocandin P1 [144]	<i>D. officinale</i>	Stem	(G. Y. Zhao et al., 2018)
Dendrocandin P2 [145]	<i>D. officinale</i>	Stem	(G. Y. Zhao et al., 2018)
Orchinol [146]	<i>D. officinale</i>	Stem	(G. Y. Zhao et al., 2018)
2,4,7-Trihydroxy-9,10-dihydro-phenanthrene [147]	<i>D. officinale</i>	Stem	(G. Y. Zhao et al., 2018)
4-Methoxy-5,9R-dihydroxy-9,10-dihydro phenanthrene 2-O- $\beta$ -D-glucopyranoside [148]	<i>D. nobile</i>	Stem	(Zhou et al., 2017)
Dihydroresveratrol [149]	<i>D. aphyllum</i>	Stem	(D. Yang et al., 2015)
Aphyllone B [150]	<i>D. aphyllum</i>	Stem	(D. Yang et al., 2015)
Aphyllal C [151]	<i>D. aphyllum</i>	Stem	(D. Yang et al., 2015)
	<i>D. loddigesii</i>	Stem	(R. J. Ma et al., 2019)


Stilbenoids	Plant	Plant part	Reference
Aphyllal D [152]	<i>D. aphyllum</i>	Stem	(D. Yang <i>et al.</i> , 2015)
Aphyllal E [153]	<i>D. aphyllum</i>	Stem	(D. Yang <i>et al.</i> , 2015)
(-)-dendroparishioid [154]	<i>D. parishii</i>	Whole plant	(Kongkatitham <i>et al.</i> , 2018a)
( <i>R</i> )-4,5,4'-trihydroxy-3,3', $\alpha$ -trimethoxybibenzyl [155]	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)
Dendrofindlaphenol A [156]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
6"-de-O-methyldendrofindlaphenol A [157]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Dendrofindlaphenol B [158]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Dendrofindlaphenol C [159]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Dendrodevonin A [160]	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
Dendrodevonin B [161]	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)



	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
[1] Aloifol I	OMe	OH	OMe	OH	H	H
[2] Amoenylin	OMe	OH	OMe	H	OMe	H
[3] Batatasin	OMe	H	H	OH	H	OH
[4] Batatasin III	OH	H	OMe	H	H	OH
[5] Brittonin A	OMe	OMe	OMe	OMe	OMe	OMe
[6] Chrysotobibenzyl	OMe	OMe	OMe	OMe	OMe	H
[7] Chrysotoxine	OMe	OH	OMe	OMe	OMe	H
[8] Crepidatin	OMe	OMe	OMe	OMe	OH	H
[9] Cumulatin	OMe	OMe	OH	OH	OMe	OMe
[10] 4,5 Dihydroxy-3,3'-dimethoxybibenzyl	OH	OH	OMe	H	H	OMe
[11] 3,3'-Dihydroxy-4,5-dimethoxybibenzyl	OMe	OMe	OH	H	H	OH
[12] 3,4'-Dihydroxy-5-methoxybibenzyl	OH	H	OMe	H	OH	H
[13] 3,4'-Dihydroxy-5,5'-dimethoxydihydrostilbene	OH	H	OMe	OMe	OH	H


**Figure 3.** Structures of stilbenoids from *Dendrobium*






	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
[14] 4,5-Dihydroxy-3,3'-dimethoxybibenzyl	OMe	OH	OH	H	H	OMe
[15] Erianin	OMe	OMe	OMe	H	OMe	OH
[16] Gigantol	OMe	H	H	H	OH	OMe
[17] 4-Hydroxy-3,5,3'-trimethoxybibenzyl	OMe	OH	OMe	H	H	OMe
[18] 5-Hydroxy-3,4,3',4',5'-pentamethoxybibenzyl	OMe	OMe	OH	OMe	OMe	OMe
[19] Isoamoenylin	OMe	OMe	OMe	H	H	OH
[20] Moniliformine	OH	OH	OMe	H	OMe	H
[21] Moscatilin	OMe	OH	OMe	H	OH	OMe
[22] 3,3',4-Trihydroxybibenzyl	OH	OH	H	H	H	OH
[23] 3,3',5-Trihydroxybibenzyl	OH	H	OH	H	H	OH
[24] 3,5,4'-Trihydroxybibenzyl	OH	H	OH	H	OH	H

Figure 3. Structures of stilbenoids from *Dendrobium* (continued)




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
[25] 4,5,4'-Trihydroxy-3-3'-dimethoxybibenzyl	OMe	OH	OH	OH	OMe
[26] Tristin	OH	H	OH	OH	OMe
[27] Dendromonilside E	OGlc	OGlc	OMe	OMe	H
[28] 5,4'-Dihydroxy-3,4,3'-trimethoxybibenzyl	OH	OMe	OMe	OH	OMe
[29] 4,3',4'-trihydroxy-3,5-dimethoxybibenzyl	OMe	OH	OMe	OH	OH





	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>
[30] Dendrophenol	OMe	OH	OMe	OH	OH	H	H
[31] 3,4-Dihydroxy-5,4'-dimethoxybibenzyl	OH	OH	OMe	H	OMe	H	H
[32] 4,4'-Dihydroxy-3,5-dimethoxybibenzyl	OMe	OH	OMe	H	OH	H	H
[33] Loddigesiinol C	OMe	OH	OMe	H	OH	OMe	OMe
[34] 3-O-Methylgigantol	OMe	H	OH	OMe	OMe	H	H

**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)



[35] Dendrocandin A


[36] Dendrocandin B

[37] Dendrocandin C  $R_1 = H$   $R_2 = OMe$ [40] Dendrocandin F  $R = OMe$ [38] Dendrocandin D  $R_1 = H$   $R_2 = OEt$ [41] Dendrocandin G  $R = OH$ [39] Dendrocandin E  $R_1 = OH$   $R_2 = H$ 

[42] Dendrocandin H

[43] Dendrosinen A  $R_1 = OMe$   $R_2 = OH$ [44] Dendrosinen B  $R_1 = OH$   $R_2 = H$ 

**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)



**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)






Figure 3. Structures of stilbenoids from *Dendrobium* (continued)



**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)



R<sub>1</sub> R<sub>2</sub> R<sub>3</sub> R<sub>4</sub> R<sub>5</sub> R<sub>6</sub> R<sub>7</sub> R<sub>8</sub>


[69] 2,2'-Dihydroxy-3,3',4,4',7,7'-hexa methoxy-9,9',10,10'-tetrahydro-1,1'-biphenanthrene

OMe OH OMe OMe OH OMe OMe OMe

[70] 2,2'-Dimethoxy-4,4',7,7'-tetrahydroxy-9,9',10,10'-tetrahydro-1,1'-biphenanthrene

OH OMe H OH OMe H OH OH

[71] Flavanthrin OH OH H OMe OH H OMe OH



[72] Phoyunnanin C


[73] Phoyunnanin E




[74] Amoenumin

[75] Crystalltone

Figure 3. Structures of stilbenoids from *Dendrobium* (continued)




[76] Dendropalpebrone




[77] Chrysotoxol A R = H


[78] Chrysotoxol B R = OMe

[79] Confusarin R<sub>1</sub> = OMe R<sub>2</sub> = OH

[80] 2,6-Dihydroxy-1,5,7-trimethoxyphenanthrene

R<sub>1</sub> = OH R<sub>2</sub> = OMe

[81] Dendrochrysanene



[82] Bulbophyllanthrin

[83] 5-Hydroxy-2,4-dimethoxy phenanthrene

[84] 3-Hydroxy-2,4,7-trimethoxy phenanthrene

R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>
OH	OH	H	H
H	OH	H	H
OH	H	OMe	H

Figure 3. Structures of stilbenoids from *Dendrobium* (continued)








Figure 3. Structures of stilbenoids from *Dendrobium* (continued)




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
[96] Coelonin	OH	H	OMe	H	H	OH
[97] 9,10-Dihydromoscatin	H	H	OH	OMe	H	OH
[98] 9,10-Dihydrophenanthrene-2,4,7-triol	OH	H	OH	H	H	OH
[99] 4,5-Dihydroxy-2,3-dimethoxy-9,10-dihydrophenanthrene	OMe	OMe	OH	OH	H	H
[100] 4,5-Dihydroxy-2,6-dimethoxy-9,10-dihydrophenanthrene	OMe	H	OH	OH	OMe	H
[101] 4,5-Dihydroxy-3,7-dimethoxy-9,10-dihydrophenanthrene	H	OMe	OH	OH	H	OMe
[102] 4,5-Dihydroxy-2-methoxy-9,10-dihydrophenanthrene	OMe	H	OH	OH	H	H
[103] Lusianthridin	OMe	H	OH	H	H	OH

**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)



	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>
[104] 2,7-Dihydroxy-3,4,6-trimethoxy-9,10-dihydrophenanthrene		OH	OMe	OMe	H	OMe	OH
[105] 2,8-Dihydroxy-3,4,7-trimethoxy-9,10-dihydrophenanthrene		OH	OMe	OMe	H	H	OMe
[106] 4,7-Dihydroxy-2,3,6-trimethoxy-9,10-dihydrophenanthrene		OMe	OMe	OH	H	OMe	OH
[107] Ephemeranthol A		OH	H	H	OH	OMe	OMe
[108] Ephemeranthol C		OH	OH	OMe	OH	H	H
[109] Erianthridin		OH	OMe	OMe	H	H	OH
[110] Flavanthridin		OH	H	H	OMe	OH	OMe
[111] Hircinol		OH	H	OMe	OH	H	H
[112] 3-Hydroxy-2,4,7-trimethoxy-9,10-dihydro-phenanthrene		OMe	OH	OMe	H	H	OMe
[113] 7-hydroxy-2,3,4-trimethoxy-9,10-dihydrophenanthrene		OH	H	H	OMe	OMe	OMe
[114] Dendroinfundin A		OMe	OMe	OH	H	H	OMe
[115] Dendroinfundin B		OMe	OMe	OH	OH	H	H
[116] 3,4-Dimethoxy-1-(methoxymethyl)-9,10-dihydrophenanthrene-2,7-diol		OH	H	H	OMe	OMe	-CH <sub>2</sub> OMe

**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
[117] 2-Hydroxy-4,7-dimethoxy-9,10-dihydrophenanthrene	OMe	H	OMe	H	H
[118] 7-Methoxy-9,10-dihydrophenanthrene-2,4,5-triol	OH	OH	OMe	H	H
[119] 2,5,7-Trihydroxy-4-methoxy-9,10-dihydrophenanthrene	OMe	OH	OH	H	H
[120] Plicatol C	H	OMe	OH	H	OMe
[121] Rotundatin	H	OMe	OH	H	OH


  

	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>	R <sub>7</sub>
[122] 2,5-Dihydroxy-3,4-dimethoxyphenanthrene	OH	OMe	OMe	OH	H	H	H
[123] 2,5-Dihydroxy-4,9-dimethoxyphenanthrene	OH	H	OMe	OH	H	H	OMe
[124] 2,8-Dihydroxy-3,4,7-trimethoxyphenanthrene	OH	OMe	OMe	H	OMe	OH	H
[125] Epheranthol B	H	H	OMe	OH	OMe	H	H
[126] Fimbriol B	OH	OMe	OH	H	H	H	H
[127] Flavanthrinin	H	H	OMe	H	OH	H	H


**Figure 3.** Structures of stilbenoids from *Dendrobium* (continued)




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub>
[128] Loddigesinol A	H	OMe	H	H	OH	H
[129] Nudol	OMe	H	OH	H	H	H
[130] Plicatol A	H	OH	H	H	OMe	OMe
[131] Plicatol B	H	OH	H	H	H	H
[132] 2,3,5-Trihydroxy-4,9-dimethoxyphenanthrene	OH	OH	H	H	OMe	H
[133] 3,4,8-Trimethoxyphenanthrene-2,5-diol	OMe	OH	H	OMe	H	H




[134] Aphyllone A



[135] 2,4,5,9-Tetrahydroxy-9,10-dihydrophenanthrene




[136] 1,5,7-Trimethoxyphenanthren-2-ol




[137] 1,5-Dihydroxy-3,4,7-trimethoxy-9,10-dihydrophenanthrene


Figure 3. Structures of stilbenoids from *Dendrobium* (continued)




[138] 2,5,9 $S$ -Trihydroxy-9,10-dihydrophenanthrene  
-4- $O$ - $\beta$ -D-glucopyranoside



[139] Loddigesiinol G R = H




[140] Loddigesiinol H R = OH



[141] Loddigesiinol I

[142] Loddigesiinol J



[143] Dendrowillol A

Figure 3. Structures of stilbenoids from *Dendrobium* (continued)







Figure 3. Structures of stilbenoids from *Dendrobium* (continued)




[154] (-)-dendroparishiol


[155] (R)-4,5,4'-trihydroxy-3,3', $\alpha$ -trimethoxybibenzyl

[156] 6''-de-O-methyldendrofindlaphenol A R = Me


[157] 6''-de-O-methyldendrofindlaphenol A R = H




[158] Dendrofindlaphenol B



[159] Dendrofindlaphenol C



[160] Dendrodevonin A



[161] Dendrodevonin B

Figure 3. Structures of stilbenoids from *Dendrobium* (continued)



**Table 2.** Flavonoids from genus *Dendrobium*

Flavonoids	Plant	Plant part	Reference
(2S)-Homoeriodictyol [162]	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
Naringenin [163]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. densiflorum</i>	Stem	(C. Fan <i>et al.</i> , 2001)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
(2S)-Eriodictyol [164]	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
Vicenin-2 [165]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
Apigenin [166]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
	<i>D. williamsonii</i>	Whole plant	(Rungwichaniwat <i>et al.</i> , 2014)
5,6-Dihydroxy-4'-methoxyflavone [167]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Chrysoeriol [168]	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
Luteolin [169]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Whole plant	(Liu <i>et al.</i> , 2009)
	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
6-C-( $\alpha$ -Arabinopyranosyl)-8-C-[(2-O- $\alpha$ -rhamnopyranosyl)- $\beta$ -galactopyranosyl] apigenin [170]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)

Flavonoids	Plant	Plant part	Reference
6-C-( $\alpha$ -Arabinopyranosyl)-8-C- [(2-O- $\alpha$ -rhamnopyranosyl)- $\beta$ - glucopyranosyl] apigenin [171]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
6'''-Glucosyl-vitexin [172]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Isoschaftoside [173]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
Isoviolanthin [174]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
6-C-[(2-O- $\alpha$ -Rhamno pyranosyl)- $\beta$ -gluco pyranosyl]-8-C-( $\alpha$ - arabinopyranosyl) apigenin [175]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
6-C-( $\beta$ -Xylopyranosyl)-8-C- [(2-O- $\alpha$ -rhamnopyranosyl)- $\beta$ - gluco pyranosyl]apigenin [176]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
Kaempferol [177]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
Kaempferol-3-O- $\alpha$ -L- rhamnopyranoside [178]	<i>D. secundum</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Kaempferol-3,7-O-di- $\alpha$ -L- rhamnopyranoside [179]	<i>D. secundum</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Kaempferol-3-O- $\alpha$ -L- rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ - D-glucopyranoside [180]	<i>D. capillipes</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Kaempferol-3-O- $\alpha$ -L- rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ - D-xylopyranoside [181]	<i>D. capillipes</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Quercetin-3-O- $\alpha$ -L- rhamnopyranoside [182]	<i>D. secundum</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)
Quercetin-3-O- $\alpha$ -L- rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ - D-xylopyranoside [183]	<i>D. capillipes</i>	Stem	(Phechrmeekha <i>et al.</i> , 2012)

Flavonoids	Plant	Plant part	Reference
5-Hydroxy-3-methoxy-flavone-7-O-[ $\beta$ -D-aposyl-(1 $\rightarrow$ 6)]- $\beta$ -D-glucoside [184]	<i>D. devonianum</i>	Stem	(Sun <i>et al.</i> , 2014)
Isorhamnetin-3-O- $\beta$ -D-rutinoside [185]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2017)
(S)-5,5',7-trihydroxy-3',4'-dimethoxyflavanone [186]	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)






Figure 4. Structures of flavonoids from *Dendrobium*




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
[172] 6'''-Glucosyl-vitexin	H	H	-Glc
[173] Isoschaftoside	H	-Ara	-Glc
[174] Isoviolanthin	H	-Rha	-Glc
[175] 6-C-[(2-O-a-Rhamnopyranosyl)- $\beta$ -glucopyranosyl]-8-C-(a-arabinopyranosyl)apigenin	H	-Glc-Rha	-Ara
[176] 6-C-( $\beta$ -Xylopyranosyl)-8-C-[(2-O-a-rhamnopyranosyl)- $\beta$ -glucopyranosyl]apigenin	H	-Xyl	-Glc-Rha
[177] Kaempferol	OH	H	H



	R <sub>1</sub>	R <sub>2</sub>
[178] Kaempferol-3-O-a-L-rhamnopyranoside	O-Rha	OH
[179] Kaempferol-3,7-O-di-a-L-rhamnopyranoside	O-Rha	O-Rha
[180] Kaempferol-3-O-a-L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-glucopyranoside	O-Glc-Rha	OH
[181] Kaempferol-3-O-a-L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-xylopyranoside	O-Xyl-Rha	OH

**Figure 4.** Structures of flavonoids from *Dendrobium* (continued)




[182] Quercetin-3-O- $\alpha$ -L-rhamnopyranoside


R = O-Rha

[183] Quercetin-3-O- $\alpha$ -L-rhamnopyranosyl-(1 $\rightarrow$ 2)- $\beta$ -D-xylopyranoside

R = O-Xyl-Rha



[184] 5-Hydroxy-3-methoxy-flavone-7-O-[[ $\beta$ -D-aposyl-(1 $\rightarrow$ 6)]- $\beta$ -D-glucoside



[185] Isorhamnetin-3-O- $\beta$ -D-rutinoside [186] (S)-5,5',7-trihydroxy-3',4'-dimethoxyflavanone

Figure 4. Structures of flavonoids from *Dendrobium* (continued)


**Table 3.** Terpenoids from *Dendrobium*

Terpenoids	Plant	Plant part	Reference
Aduncin [187]	<i>D. aduncum</i>	Whole plant	(Gawell & Leander, 1976)
Amoenin [188]	<i>D. amoenum</i>	Whole plant	(Dahmen & Leander, 1978; Majumder <i>et al.</i> , 1999)
Amotin [189]	<i>D. amoenum</i>	Whole plant	(Dahmen & Leander, 1978; Majumder <i>et al.</i> , 1999)
asiatic acid [190]	<i>D. parishii</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
a-Dihydropicrotoxinin [191]	<i>D. moniliforme</i>	Stem	(Bi <i>et al.</i> , 2004)
Dendrobane A [192]	<i>D. nobile</i>	Stem	(X. Zhang, H. Gao, <i>et al.</i> , 2007)
Dendronobilin A [193]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin B [194]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin C [195]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin D [196]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin E [197]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin F [198]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin G [199]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin H [200]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
Dendronobilin I [201]	<i>D. nobile</i>	Stem	(Xue Zhang <i>et al.</i> , 2007)
	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendronobilin J [202]	<i>D. nobile</i>	Stem	(X. Zhang, H. Gao, <i>et al.</i> , 2007)
Dendronobilin K [203]	<i>D. nobile</i>	Stem	(X. Zhang, F. J. Tu, <i>et al.</i> , 2008)
Dendronobilin L [204]	<i>D. nobile</i>	Stem	(X. Zhang, F. J. Tu, <i>et al.</i> , 2008)
Dendronobilin M [205]	<i>D. nobile</i>	Stem	(X. Zhang, F. J. Tu, <i>et al.</i> , 2008)
Dendronobilin N [206]	<i>D. nobile</i>	Stem	(X. Zhang, F. J. Tu, <i>et al.</i> , 2008)
	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)


Terpenoids	Plant	Plant part	Reference
Dendrowardol A [207]	<i>D. wardianum</i>	Stem	(W. W. Fan <i>et al.</i> , 2013)
Dendrowardol B [208]	<i>D. wardianum</i>	Stem	(W. W. Fan <i>et al.</i> , 2013)
Dendrowardol C [209]	<i>D. wardianum</i>	Stem	(W. W. Fan <i>et al.</i> , 2013)
Corchoionoside C [210]	<i>D. polyanthum</i>	Stem	(Hu <i>et al.</i> , 2009)
Crystallinin [211]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
	<i>D. findlayanum</i>	Whole plant	(Qin <i>et al.</i> , 2011)
Findlayanin [212]	<i>D. findlayanum</i>	Whole plant	(Qin <i>et al.</i> , 2011)
3-Hydroxy-2-oxodendrobine [213]	<i>D. nobile</i>	Stem	(H. Wang <i>et al.</i> , 1985)
Dendrobine [214]	<i>D. nobile</i>	Stem	(H. Wang <i>et al.</i> , 1985)
	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
2-Hydroxydendrobine [215]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Findlayine A [216]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Findlayine B [217]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Findlayine C [218]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Findlayine D [219]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2018)
Dendromonilaside A [220]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
Dendromonilaside B [221]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
Dendromonilaside C [222]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
Dendromonilaside D [223]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
Dendronobiloside A [224]	<i>D. nobile</i>	Stem	(W. M. Zhao <i>et al.</i> , 2001); (Ye & Zhao, 2002)
Dendronobiloside B [225]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002; W. M. Zhao <i>et al.</i> , 2001)
Dendronobiloside C [226]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002; W. M. Zhao <i>et al.</i> , 2001)
Dendronobiloside D [227]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002; W. M. Zhao <i>et al.</i> , 2001)
Dendronobiloside E [228]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002; W. M. Zhao <i>et al.</i> , 2001)
Dendroside A [229]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
	<i>D. nobile</i>	Stem	(W. M. Zhao <i>et al.</i> , 2001)
		Stem	(Ye & Zhao, 2002)

Terpenoids	Plant	Plant part	Reference
	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendroside B [230]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)
Dendroside C [231]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)
Dendroside D [232]	<i>D. nobile</i>	Stem	(Ye & Zhao, 2002)
Dendroside E [233]	<i>D. nobile</i>	Stem	(Ye <i>et al.</i> , 2002)
Dendroside F [234]	<i>D. moniliforme</i>	Stem	(Ye <i>et al.</i> , 2002)
Dendroside G [235]	<i>D. nobile</i>	Stem	(Ye <i>et al.</i> , 2002)
Dendrowillin A [236]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2019)
Dendrowillin B [237]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2019)
(-)-Picrotin [238]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2019)
10 $\beta$ ,12,14-trihydroxyaromadendrane [239]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
10 $\beta$ ,13,14-trihydroxyaromadendrane [240]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendrofindlayanoside A [241]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendrofindlayanoside B [242]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendrofindlayanoside C [243]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
Dendrofindlayanobilin [244]	<i>D. findlayanum</i>	Stem	(D. Yang <i>et al.</i> , 2019)
(+)-(1R,2S,3R,4S,5R,6S,9R)-3,11,12-trihydroxypicrotoxane-2(15)-lactone [245]	<i>D. nobile</i>	Stem	(C. Ma <i>et al.</i> , 2019)
(-)-(1S,2R,3S,4R,5S,6R,9S,12R)-3,11,13-trihydroxypicrotoxane-2(15)-lactone [246]	<i>D. nobile</i>	Stem	(C. Ma <i>et al.</i> , 2019)
(+)-(1R,5R,6S,8R,9R)-8,12-dihydroxy-copacamphan-3-en-2-one [247]	<i>D. nobile</i>	Stem	(C. Ma <i>et al.</i> , 2019)
Dendroterpene A [248]	<i>D. nobile</i>	Stem	(P. Wang <i>et al.</i> , 2019)
Dendroterpene B [249]	<i>D. nobile</i>	Stem	(P. Wang <i>et al.</i> , 2019)
Dendroterpene C [250]	<i>D. nobile</i>	Stem	(P. Wang <i>et al.</i> , 2019)
Dendroterpene D [251]	<i>D. nobile</i>	Stem	(P. Wang <i>et al.</i> , 2019)







[187] Aduncin




[188] Amoenin




[189] Amotin




[190] Asiatic acid




[191] a-Dihydropicrotoxinin




[192] Dendrobane A




[193] Dendronobilin A




[194] Dendronobilin B




[195] Dendronobilin C




[196] Dendronobilin D



[197] Dendronobilin E




[198] Dendronobilin F




[199] Dendronobilin G


**Figure 5.** Structures of terpenoids from *Dendrobium*




[200] Dendronobilin H




[201] Dendronobilin I




[202] Dendronobilin J




[203] Dendronobilin K




[204] Dendronobilin L




[205] Dendronobilin M




[206] Dendronobilin N




[207] Dendrowardol A




[208] Dendrowardol B



[209] Dendrowardol C



[210] Corchoionoside C



[211] Crystallinin

Figure 5. Structures of terpenoids from *Dendrobium* (continued)







Figure 5. Structures of terpenoids from *Dendrobium* (continued)




[227] Dendronobiloside D



[228] Dendronobiloside E




[229] Dendroside A




[230] Dendroside B R = OGlc


[231] Dendroside C R = OH




[232] Dendroside D




[233] Dendroside E



[234] Dendroside F R = H





[236] Dendrowillin A R = OH




[238] (-)-Picrotin


[235] Dendroside G R = OH

[237] Dendrowillin B R = H

[239] 10 $\beta$ , 12, 14-trihydroxyaromadendrane[240] 10 $\beta$ , 13, 14-trihydroxyaromadendraneFigure 5. Structures of terpenoids from *Dendrobium* (continued)



**Figure 5.** Structures of terpenoids from *Dendrobium* (continued)



[250] Dendroterpene C     R = H

[251] Dendroterpene D     R = OH

**Figure 5.** Structures of terpenoids from *Dendrobium* (continued)

**Table 1** Miscellaneous compounds from *Dendrobium*

Categories and compounds	Plant	Plant part	Reference
<b>Aliphatic acid derivatives</b>			
Aliphatic acids [252]	<i>D. clavatum</i> var. <i>aurantiacum</i>	Stem	(S. J. Chang <i>et al.</i> , 2001)
Aliphatic alcohols [253]	<i>D. clavatum</i> var. <i>aurantiacum</i>	Stem	(S. J. Chang <i>et al.</i> , 2001)
Malic acid [254]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
Dimethyl malate [255]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
(-)-Shikimic acid [256]	<i>D. fuscescens</i>	Whole plant	(Talapatra <i>et al.</i> , 1989)
	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. pulchellum</i>	Stem	(Chanvorachote <i>et al.</i> , 2013)
Isopentyl butyrate [257]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)
Dendrodevonic acid A [258]	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
Dendrodevonic acid B [259]	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
<b>Benzoic acid derivatives and phenolic compounds</b>			
3-Hydroxy-2-methoxy-5,6-dimethylbenzoic acid [260]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Salicylic acid [261]	<i>D. huoshanense</i>	Aerial part	(C. C. Chang <i>et al.</i> , 2010)

Categories and compounds	Plant	Plant part	Reference
	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Vanilloside [262]	<i>D. denneanum</i>	Stem	(Pan <i>et al.</i> , 2012)
	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
Gallic acid [263]	<i>D. longicornu</i>	Whole plant	(J. T. Li <i>et al.</i> , 2009)
Syringic acid [264]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Vanillic acid [265]	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)
	<i>D. williamsonii</i>	Whole plant	(Rungwichaniwat <i>et al.</i> , 2014)
Antiarol [266]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Ethylhaematommate [267]	<i>D. longicornu</i>	Whole plant	(J. T. Li <i>et al.</i> , 2009)
<i>p</i> -Hydroxy benzaldehyde [268]	<i>D. devonianum</i>	Whole plant	(Sun <i>et al.</i> , 2014)
	<i>D. falconeri</i>	Stem	(Sritularak & Likhitwitayawuid, 2009)
	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Methyl $\beta$ -orsellinate [269]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Tachioside [270]	<i>D. denneanum</i>	Stem	(Pan <i>et al.</i> , 2012)
Alkyl 4'-hydroxy-trans-cinnamates [271]	<i>D. clavatum</i> var. <i>aurantiacum</i>	Stem	(S. J. Chang <i>et al.</i> , 2001)
Alkyl trans-ferulate [272]	<i>D. clavatum</i> var. <i>aurantiacum</i>	Stem	(S. J. Chang <i>et al.</i> , 2001)
	<i>D. scabrilingue</i>	Whole plant	(Sarakulwattana <i>et al.</i> , 2018)
Defuscin [273]	<i>D. fuscescens</i>	Whole plant	(Talapatra <i>et al.</i> , 1989)
	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
<i>n</i> -Octacosyl ferulate [274]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. moniliforme</i>	Stem	(Bi <i>et al.</i> , 2004)
<i>n</i> -Triacetyl <i>p</i> -hydroxy-cis-cinnamate [275]	<i>D. moniliforme</i>	Stem	(Bi <i>et al.</i> , 2004)

Categories and compounds	Plant	Plant part	Reference
Tetratriacontanyl- <i>trans-p</i> -coumarate [276]	<i>D. williamsonii</i>	Whole plant	(Rungwichaniwat <i>et al.</i> , 2014)
<i>p</i> -hydroxyphenethyl- <i>trans</i> -ferulate [277]	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)
<i>n</i> -Docosyl <i>trans</i> -ferulate [278]	<i>D. longicornu</i>	Whole plant	(J. T. Li <i>et al.</i> , 2009)
	<i>D. williamsonii</i>	Whole plant	(Rungwichaniwat <i>et al.</i> , 2014)
<i>trans</i> -Tetracosyl ferulate [279]	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
<i>cis</i> -Hexacosanoyl ferulate [280]	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
Ferulaldehyde [281]	<i>D. longicornu</i>	Whole plant	(J. T. Li <i>et al.</i> , 2009)
Ferulic acid [282]	<i>D. secundum</i>	Stem	(Sritularak, Duangrak, <i>et al.</i> , 2011)
2-( <i>p</i> -Hydroxyphenyl) ethyl <i>p</i> -coumarate [283]	<i>D. falconeri</i>	Stem	(Sritularak & Likhitwitayawuid, 2009)
Dihydroconiferyl dihydro- <i>p</i> -coumarate [284]	<i>D. formosum</i>	Whole plant	(Inthongkaew <i>et al.</i> , 2017)
	<i>D. loddigesii</i>	Stem	(R. J. Ma <i>et al.</i> , 2019)
	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
1-[4-( $\beta$ -D-Glucopyranosyloxy)-3,5-dimethoxyphenyl]-1-propanone [285]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
3-Hydroxy-1-(4-hydroxy-3,5-dimethoxyphenyl)-1-propanone [286]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)



Categories and compounds	Plant	Plant part	Reference
2-Hydroxy-3-(4-hydroxy-3-methoxyphenyl)-3-methoxypropyl-3-(4-hydroxyphenyl)-propanoate [287]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
Coniferyl alcohol [288]	<i>D. trigonopus</i>	Stem	(Hu <i>et al.</i> , 2008b)
( <i>E</i> )-Coniferyl aldehyde [289]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
Sinapicaldehyde [290]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
Decumbic acid A [291]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
Decumbic acid B [292]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
(–)-Decumbic acid [293]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
(+)-Dendrolactone [294]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
4-(3-Hydroxyphenyl)-2-butanone [295]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
3-Hydroxy-1(3-methoxy-4-hydroxyphenyl)-propan-1-one [296]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
3',4',5'-Trimethoxy cinnamyl acetate [297]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2016)
Alatusol A [298]	<i>D. hainanense</i>	Aerial part	(Y. Y. Zhang <i>et al.</i> , 2019)
<i>p</i> -Hydroxyphenyl propionic methyl ester [299]	<i>D. aphyllum</i>	Whole plant	(Yegao Chen <i>et al.</i> , 2008)
Phloretic acid [300]	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
Dihydroconiferyl alcohol [301]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Salidrosol [302]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Shashenoside I [303]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
Syringin [304]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)


Categories and compounds	Plant	Plant part	Reference
Tetracosyl( <i>Z</i> )- <i>p</i> -coumarate [305]	<i>D. falconeri</i>	Whole plant	(Sritularak & Likhitwitayawuid, 2009)
(7 <i>S</i> ,8 <i>R</i> )-Dehydrodiconiferyl alcohol 9'- $\beta$ -D-glucopyranoside [306]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2017)
Koaburaside [307]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2017)
Juniperoside [308]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2017)
Dehydrodiconiferyl alcohol-4- $\beta$ -D-glucoside [309]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2017)
(3 <i>R</i> ,3' <i>S</i> ,4 <i>R</i> ,4' <i>S</i> )-3,3',4,4'-Tetrahydro-6,6'-dimethoxy[3,3'-bi-2 <i>H</i> -benzopyran]-4,4'-diol [310]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
<i>threo</i> -7- <i>O</i> -ethyl-9- <i>O</i> -(4-hydroxyphenyl) propionyl-guaiacyl glycerol [311]	<i>D. loddigesii</i>	Stem	(Ma, R. J., <i>et al.</i> , 2019b)
<b>Coumarins</b>			
Ayapin [312]	<i>D. densiflorum</i>	Stem	(Fan <i>et al.</i> , 2001)
Coumarin [313]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. clavatum</i> var. <i>aurantiacum</i>	Stem	(S. J. Chang <i>et al.</i> , 2001)
Denthyrsin [314]	<i>D. thyrsiflorum</i>	Stem	(G. N. Zhang <i>et al.</i> , 2005)
Scoparone [315]	<i>D. densiflorum</i>	Stem	(Fan <i>et al.</i> , 2001)
	<i>D. thyrsiflorum</i>	Stem	(G. N. Zhang <i>et al.</i> , 2005)
	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
Scopoletin [316]	<i>D. densiflorum</i>	Stem	(Fan <i>et al.</i> , 2001)
<b>Lignans and neolignans</b>			
Episyngaresinol [317]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2008)

Categories and compounds	Plant	Plant part	Reference
Episingaresinol 4''-O- $\beta$ -D-glucopyranoside [318]	<i>D. moniliforme</i>	Stem	(C. S. Zhao <i>et al.</i> , 2003)
(-)-(7S,8R,7'E)-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-7,9'-bis-O- $\beta$ -D-glucopyranoside [319]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
Lyoniresinol [320]	<i>D. chrysanthum</i>	Stem	(Ye <i>et al.</i> , 2004)
(-)-Syringaresinol-4,4'-bis-O- $\beta$ -D-glucopyranoside [321]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
Syringaresinol-4-O-D-monoglucopyranoside [322]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(Xiong <i>et al.</i> , 2013)
Dendrocoumarin [323]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2018)
Itolide A [324]	<i>D. nobile</i>	Stem	(Zhou <i>et al.</i> , 2018)
(-)-Medioresinol [325]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2008)
(-)-Pinoresinol [326]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2008)
(+)-Pinoresinol [327]	<i>D. devonianum</i>	Stem	(Wu <i>et al.</i> , 2019)
Erythro-1-(4-O- $\beta$ -D-glucopyranosyl-3-methoxyphenyl)-2-[4-(3-hydroxypropyl)-2,6-dimethoxyphenoxy]-1,3-propanediol [328]	<i>D. longicornu</i>	Stem	(Hu <i>et al.</i> , 2008a)
Syringaresinol [329]	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2008)
	<i>D. secundum</i>	Stem	(Sritularak, Duangrak, <i>et al.</i> , 2011)
	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Acanthoside B [330]	<i>D. chrysanthum</i>	Stem	(Ye <i>et al.</i> , 2004)

Categories and compounds	Plant	Plant part	Reference
Liriodendrin [331]	<i>D. brymerianum</i>	Whole plant	(Chen <i>et al.</i> , 2014)
	<i>D. pulchellum</i>	Stem	(Chanvorachote <i>et al.</i> , 2013)
(-)-(8 <i>R</i> ,7' <i>E</i> )-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-9,9'-diol 4,9-bis- <i>O</i> - $\beta$ -D-glucopyranoside [332]	<i>D. auranticum</i> var. <i>denneanum</i>	Stem	(Li <i>et al.</i> , 2014)
(-)-(8 <i>S</i> ,7' <i>E</i> )-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-9,9'-diol 4,9-bis- <i>O</i> - $\beta$ -D-glucopyranoside [333]	<i>D. auranticum</i> var. <i>denneanum</i>	Stem	(Li <i>et al.</i> , 2014)
(-)-(8 <i>R</i> ,7' <i>E</i> )-4-Hydroxy-3,3',5,5',9'-penta methoxy-8,4'-oxyneolign-7'-ene-9-ol 4,9-bis- <i>O</i> - $\beta$ -D-glucopyranoside [334]	<i>D. auranticum</i> var. <i>denneanum</i>	Stem	(Li <i>et al.</i> , 2014)
<b>Fluorenones</b>			
Denchrysan B [335]	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
Denchrysan A [336]	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)
Dendroflorin [337]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
Dengibsin [338]	<i>D. aurantiacum</i> var. <i>denneanum</i>	Stem	(L. Yang, Wang, <i>et al.</i> , 2006)
	<i>D. chrysanthum</i>	Stem	(L. Yang, Qin, <i>et al.</i> , 2006)
	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)


Categories and compounds	Plant	Plant part	Reference
Nobilone [339]	<i>D. brymerianum</i>	Whole plant	(Klongkumnuankarn <i>et al.</i> , 2015)
	<i>D. nobile</i>	Stem	(X. Zhang <i>et al.</i> , 2007)
	<i>D. palpebrae</i>	Whole plant	(Kyokong <i>et al.</i> , 2019)
1,4,5-Trihydroxy-7-methoxy-9 <i>H</i> -fluoren-9-one [340]	<i>D. chrysotoxum</i>	Whole plant	(Y. P. Li <i>et al.</i> , 2009)
2,4,7-Trihydroxy-1,5-dimethoxy-9-fluorenone [341]	<i>D. chrysotoxum</i>	Stem	(H. Yang <i>et al.</i> , 2004)
<b>Others</b>			
3,6,9-Trihydroxy-3,4-dihydroanthracen-1-(2 <i>H</i> )-one [342]	<i>D. chrysotoxum</i>	Stem	(Hu <i>et al.</i> , 2012)
Palmarumycin JC2 [343]	<i>D. crystallinum</i>	Stem	(L. Wang <i>et al.</i> , 2009)
Dehydrovomifoliol [344]	<i>D. loddigesii</i>	Whole plant	(Ito <i>et al.</i> , 2010)
2,6-Dimethoxy Benzoquinone [345]	<i>D. chryseum</i>	Stem	(Ma <i>et al.</i> , 1998)
4-(2-Hydroxypropyl)- 2(5 <i>H</i> )-furanone [346]	<i>D. tortile</i>	Whole plant	(Limpanit <i>et al.</i> , 2016)
5,7-Dihydroxy chromen-4-one [347]	<i>D. ellipsophyllum</i>	Whole plant	(Tanagornmeatar <i>et al.</i> , 2014)
Balanophonin [348]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Ergosta-8(9),22-diene-3,5,6,7-tetraol [349]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Stigmast-4-en-3 $\alpha$ , 6 $\beta$ -diol [350]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
3 $\beta$ -Hydroxy-5 $\alpha$ ,8 $\alpha$ -epidioxyergosta-6,9,22-triene [351]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
$\beta$ -Sitosterol [352]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)
Daucosterol [353]	<i>D. williamsonii</i>	Whole plant	(M. Yang <i>et al.</i> , 2018)

Categories and compounds	Plant	Plant part	Reference
Anosmine [354]	<i>D. parishii</i>	Whole plant	(Hemscheidt & Spenser, 1991)
di- <i>p</i> -hydroxyphenylpropionic acid- <i>p</i> -coumaric acid lactone [355]	<i>D. chrysanthum</i>	Whole plant	(Cai <i>et al.</i> , 2018)
RF-3192C [356]	<i>D. scabrilingue</i>	Whole plant	(Sarakuwattana <i>et al.</i> , 2018)
Crepidatamine C [357]	<i>D. crepidatum</i>	Stem	(Xu <i>et al.</i> , 2019)
Crepidatamine D [358]	<i>D. crepidatum</i>	Stem	(Xu <i>et al.</i> , 2019)
Crepidine [359]	<i>D. crepidatum</i>	Stem	(Xu <i>et al.</i> , 2019)
Isocrepidamine [360]	<i>D. crepidatum</i>	Stem	(Xu <i>et al.</i> , 2019)
Crepidamine [361]	<i>D. crepidatum</i>	Stem	(Xu <i>et al.</i> , 2019)




[252] Aliphatic acids      R = COOH      n = 19-31

[253] Aliphatic alcohol      R = OH      n = 22-32




[254] Malic acid    R<sub>1</sub> = R<sub>2</sub> = OH


[255] Dimethyl malate    R<sub>1</sub> = R<sub>2</sub> = OMe



[256] (-)-Shikimic acid



[257] Isopentyl butyrate




[258] Dendrodevonic acid A


[259] Dendrodevonic acid B

R = H


R = acetyl




[260] 3-Hydroxy-2-methoxy-5,6-dimethylbenzoic acid



[261] Salicylic acid



[262] Vanilloside



[263] Gallic acid    R<sub>1</sub> = OH    R<sub>2</sub> = OH

[264] Syringic acid    R<sub>1</sub> = OMe    R<sub>2</sub> = OMe

[265] Vanillic acid    R<sub>1</sub> = H    R<sub>2</sub> = OMe

Figure 6. Structures of miscellaneous compounds from *Dendrobium*








Figure 6. Structures of miscellaneous compounds from *Dendrobium* (continued)




[277] *p*-hydroxyphenethyl-*trans*-ferulate[278] *n*-Docosyl *trans*-ferulate:  $R = \text{COOCH}_2(\text{CH}_2)_{20}\text{CH}_3$ [279] *trans*-Tetracosyl ferulate  $R = \text{COOCH}_2(\text{CH}_2)_{22}\text{CH}_3$ [280] *cis*-Hexacosanoyl ferulate  $R = \text{COOCH}_2(\text{CH}_2)_{24}\text{CH}_3$ [281] Ferulaldehyde  $R = \text{CHO}$ [282] Ferulic acid  $R = \text{COOH}$ [283] 2-(*p*-Hydroxyphenyl) ethyl *p*-coumarate[284] Dihydroconiferyl dihydro-*p*-coumarate[285] 1-[4-( $\beta$ -D-Glucopyranosyloxy)-3,5-dimethoxyphenyl]-1-propanone

[286] 3-Hydroxy-1-(4-hydroxy-3,5-dimethoxyphenyl)-1-propanone

Figure 6. Structures of miscellaneous compounds from *Dendrobium* (continued)




[287] 2-Hydroxy-3-(4-hydroxy-3-methoxyphenyl)-3-methoxypropyl-3-(4-hydroxyphenyl)propanoate



[288] Coniferyl alcohol

[289] (*E*)-Coniferyl aldehyde

[290] Sinapicaldehyde



[291] Decumbic acid A

[292] Decumbic acid B


[293] (-)-Decumbic acid

[294] (+)-Dendrolactone




[295] 4-(3-Hydroxyphenyl)-2-butanone

[296] 3-Hydroxy-1-(3-methoxy-4-hydroxyphenyl)-propan-1-one




[297] 3',4',5'-Trimethoxy cinnamyl acetate






[298] Alatusol A


**Figure 6.** Structures of miscellaneous compounds from *Dendrobium* (continued)




**Figure 6.** Structures of miscellaneous compounds from *Dendrobium* (continued)

[309] Dehydrodiconiferyl alcohol-4- $\beta$ -D-glucoside[310] (3*R*,3'*S*,4*R*,4'*S*)-3,3',4,4'-Tetrahydro-6,6'-dimethoxy[3,3'-  
bi-2*H*-benzopyran]-4,4'-diol[311] *threo*-7-*O*-ethyl-9-*O*-(4-hydroxyphenyl) propionyl-guaiacyl glycerol


[312] Ayapin



[313] Coumarin




[314] Denthyrsin




[315] Scoparone R = OMe

[316] Scopoletin R = OH




[317] Episingaresinol R = H


[318] Episingaresinol 4''-*O*- $\beta$ -D-glucopyranoside  
R =  $\beta$ -D-glucoseFigure 6. Structures of miscellaneous compounds from *Dendrobium* (continued)



[319] (-)-(7*S*,8*R*,7'*E*)-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-7,9'-bis-*O*- $\beta$ -D-glucopyranoside



[320] Lyoniresinol




[321] (-)-Syringaresinol-4,4'-bis-*O*- $\beta$ -D-glucopyranoside

R = Glc

[322] Syringaresinol-4-*O*-D-monoglucopyranoside


R = H



[323] Dendrocoumarin

$R_1 = H$

$R_2 = OH$



[325] (-)-Medioresinol

R = OMe

[324] Itolide A


$R_1 = OH$

$R_2 = H$



[326] (-)-Pinoresinol


R = H

**Figure 6.** Structures of miscellaneous compounds from *Dendrobium* (continued)



[327] (+)-Pinoresinol


[328] Erythro-1-(4-O- $\beta$ -D-glucopyranosyl-3-methoxyphenyl)-2-[4-(3-hydroxypropyl)-2,6-dimethoxyphenoxy]-1,3-propanediolFigure 6. Structures of miscellaneous compounds from *Dendrobium* (continued)




[332] (-)-(8*R*,7'*E*)-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-9,9'-diol 4,9-bis-O- $\beta$ -D-glucopyranoside  
R = OH; 8*R*

[333] (-)-(8*S*,7'*E*)-4-Hydroxy-3,3',5,5'-tetramethoxy-8,4'-oxyneolign-7'-ene-9,9'-diol 4,9-bis-O- $\beta$ -D-glucopyranoside  
R = OH; 8*S*

[334] (-)-(8*R*,7'*E*)-4-Hydroxy-3,3',5,5',9'-pentamethoxy-8,4'-oxyneolign-7'-ene-9-ol 4,9-bis-O- $\beta$ -D-glucopyranoside  
R = OMe; 8*R*




[335] Denchrysan B




	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>
[336] Denchrysan A	H	OH	OH	OMe	OH
[337] Dendroflorin	OH	H	OH	OMe	OH
[338] Dengibsin	H	OH	OMe	OH	H
[339] Nobilone	H	OH	H	OMe	OH
[340] 1,4,5-Trihydroxy-7-methoxy-9 <i>H</i> -fluoren-9-one	OH	H	OH	OH	OMe
[341] 2,4,7-trihydroxy-1,5-dimethoxy-9-fluorenone	OMe	OH	OH	OMe	OH

Figure 6. Structures of miscellaneous compounds from *Dendrobium* (continued)



**Figure 6.** Structures of miscellaneous compounds from *Dendrobium* (continued)





**Figure 6.** Structures of miscellaneous compounds from *Dendrobium* (continued)

## 2. Alpha-glucosidase inhibitory activity of *Dendrobium* species

One of the largest genera of Orchidaceae is the *Dendrobium* genus. Thousands of species of *Dendrobium* can be found all over the world. Dozens of species, such as *D. officinale*, *D. nobile*, *D. huoshanense*, and *D. chrysanthum*, are grown in China. Among all of these *Dendrobium* species, *D. officinale* is one of the most widely known due to its functions in Traditional Chinese Medicine (TCM), like tonifying the stomach, promoting fluid, nourishing yin, and clearing heat. The fresh stem of *D. officinale* can be orally consumed directly and used as a soup stock or tea too. On the other hand, modern pharmacology research has shown some beneficial bioactivities of *D. officinale*, like anti-oxidant, anti-tumor, hypoglycemic activities as well as gastrointestinal regulatory functions (Zhou, C., Xie, Z., et. al., 2018).


Alpha-glucosidase inhibitors (AGIs) are a class of Oral Hypoglycemic Agents (OHAs) for the treatment of DM II. The mechanism of AGIs is to inhibit binding the active centers of  $\alpha$ -glucosidase enzyme with oligosaccharides. The treatment of DM II patients can use AGIs as drugs that inhibit the absorption of carbohydrates from the gut. Because of their nitrogen component, AGIs can block the enzymatic reaction of  $\alpha$ -glucosidase (Joshi *et. al.*, 2015). Acarbose, miglitol and voglibose are drugs that include AGIs (Patel *et al.*, 2012). Among the AGIs, acarbose has the most robust evidence base for the treatment of DM II. It is the most studied drug with proven efficacy in lowering insulin dose in postprandial hyperglycemia (PPHG) and improving insulin resistance/secretion and dyslipidemia in patients with DM II (Joshi *et al.*, 2015).

The inhibitory effect of  $\alpha$ -glucosidase in type 2 diabetes can be used to absorb postprandial glucose.  $\alpha$ -Glucosidase has a role in carbohydrates into glucose conversion. By inhibiting  $\alpha$ -glucosidase, blood glucose levels can be returned within the normal range (Van de Laar, 2008).


In previous reports, several compounds from *Dendrobium* spp. have been found to possess  $\alpha$ -glucosidase effect. Table 5 showed examples of  $\alpha$ -glucosidase inhibitors from *Dendrobium* spp.

**Table 5.** Examples of  $\alpha$ -glucosidase inhibitors from *Dendrobium* spp.


Structure name	Plant	Plant part	Parts
<b>Flavonoids</b>			
5-Hydroxy-3-methoxy-flavone-7-O-[ $\beta$ -D-apiosyl-(1 $\rightarrow$ 6)]- $\beta$ -D-glucoside [363]	<i>D. devonianum</i>	(Sun <i>et al.</i> , 2014)	whole plant
<b>Bisbibenzyls</b>			
Dendrofalconerol A [364]	<i>D. tortile</i>	(Limpanit <i>et al.</i> , 2016)	whole plant
<b>Phenanthrene Quinones</b>			
5-Methoxy-7-hydroxy-9,10-dihydro-1,4-phenanthrenequinone [365]	<i>D. formosum</i>	(Inthongkaew <i>et al.</i> , 2017)	stem
<b>Bibenzyl derivatives</b>			
Loddigesiinol J [366]	<i>D. loddigesii</i>	(Lu <i>et al.</i> , 2014)	stem
3,4-Dihydroxy-5,4'-dimethoxybibenzyl [367]	<i>D. tortile</i>	(Limpanit <i>et al.</i> , 2016)	whole plant




[363] 5-Hydroxy-3-methoxy-flavone-7-O-  
[β-D-apiosyl-(1→6)]-β-D-glucoside




[364] Dendrofalconerol A



[365] 5-Methoxy-7-hydroxy-9,10-  
dihydro-1,4-phenanthrenequinone



[366] Loddigesinol J



[367] 3,4-Dihydroxy-5,4'-dimethoxybibenzyl

Figure 7. Structures of α-glucosidase inhibitors from *Dendrobium* spp.

## CHAPTER III

### EXPERIMENTAL

#### 1. Source of plant materials

##### 1.1 *Dendrobium braianense* ( เอื้องคำปือกลาว )

*Dendrobium braianense* samples were bought from Chatuchak market, Bangkok, in March 2018. This plant was identified by Mr. Yanyong Punpreuk, Department of Agriculture, Bangkok. Voucher specimens (BS-DB-032561) has been saved at the herbarium of the Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University.

##### 1.2 *Dendrobium kentrophyllum* ( เอื้องกำพลาใหญ่ )

*Dendrobium kentrophyllum* samples were purchased from an orchid farm, Suphanburi, in August 2018. They were identified by Mr. Yanyong Punpreuk, Department of Agriculture, Bangkok. Voucher specimens (BS-DK-072561) have been saved at the herbarium of the Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University.

#### 2. General Techniques

##### 2.1 Analytical thin-layer chromatography (TLC)

###### 2.1.1 Normal phase thin-layer chromatography

Technique	One-dimension ascending
Absorbent	Silica gel 60 F254 precoated plate (E. Merck)
Temperature	Laboratory temperature (30-35 °C)
Detection	1. Ultraviolet light at wavelengths of 254 and 365 nm.

2. Spraying with anisaldehyde reagent (p-anisaldehyde 15 g in ethanol 250 mL and concentrated sulfuric acid 2.5 mL) and heating at 105 °C for 10 minutes.

### 2.1.2 Reverse phase thin-layer chromatography

<b>Technique</b>	One-dimension ascending
<b>Absorbent</b>	RP C-18 precoated on aluminum sheet (Anal Tech)
<b>Temperature</b>	Laboratory temperature (30-35 °C)
<b>Detection</b>	Ultraviolet light at wavelengths of 254 and 365 nm.

## 2.2 Column chromatography (CC)

### 2.2.1 Vacuum liquid chromatography (VLC)

<b>Adsorbent</b>	Silica gel 60 (No. 107734), size 0.063-0.200 mm (E. Merck)
<b>Packing method</b>	Dry packing
<b>Sample loading</b>	The sample was dissolved in a small volume of organic solvent, triturated with a slight amount of the adsorbent, dried, and then gradually put on the top of the column.
<b>Detection</b>	Each fraction was studied by TLC under UV light at the wavelengths of 254 and 365 nm.

### 2.2.2 Flash column chromatography (FCC), normal phase

<b>Adsorbent</b>	Silica gel 60 (No. 109385), size 0.040-0.063 mm (E. Merck)
<b>Packing method</b>	Wet packing
<b>Sample loading</b>	The sample was dissolved in a minor volume of organic solvent, triturated with a small amount of the adsorbent, dried, and then gradually put on the top of the column.
<b>Detection</b>	Fractions were studied as described in section 2.2.1

### 2.2.3 Flash column chromatography (FCC), reverse phase

<b>Adsorbent</b>	C-18 (No. 113900), size 40-63 µm (E. Merck)
<b>Packing method</b>	Wet packing

**Sample loading** The sample was dissolved with a little volume of organic solvent, and then gradually loaded on the top of the column.

**Detection** Fractions were studied as described in section 2.2.1

#### 2.2.4 Gel filtration chromatography

**Gel filter** Sephadex LH-20, particle size 25-100  $\mu\text{m}$  (GE Healthcare)

**Packing method** The gel filter was suspended in an appropriate solvent, left standing for about 24 hours, and then poured into the column and left to set tightly.

**Sample loading** It was liquefied in a little volume of the eluent and then gradually dispensed on the top of the column.

**Detection** Fractions were studied as described in section 2.2.1.

#### 2.2.5 Semi-preparative high pressure liquid chromatography (HPLC)

**Column** COSMOSIL 5C18-AR-II (10ID x 250 mm)

**Flow rate** 3 mL/min

**Mobile phase** Isocratic 50% methanol in water

**Sample preparation** It was liquefied in a little volume of the eluent and filtered through Millipore filter paper before it was injected.

**Injection volume** 1 mL

**Pump** LC-8A (Shimadzu)

**Detector** SPD-10A UV-Vis Detector (Shimadzu)

**Recorder** C-R6A Chromatopac (Shimadzu)

**Temperature** Room temperature

## 2.3 Spectroscopy

### 2.3.1 Mass spectra

Mass spectra (MS) were recorded using a Bruker micro TOF mass spectrometer (Department of Chemistry, Faculty of Science, Mahidol University).

### 2.3.2 Proton and carbon-13 nuclear magnetic resonance ( $^1\text{H}$ and $^{13}\text{C}$ -NMR) spectra

$^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectra were recorded utilizing a Bruker Avance DPX-300 FT-NMR spectrometer (Faculty of Pharmaceutical Sciences, Chulalongkorn University).  $^1\text{H}$  NMR (500 MHz) and  $^{13}\text{C}$  NMR (125 MHz) spectra were recorded on a Bruker Avance III HD 500 NMR spectrometer (Scientific and Technology Research Equipment Center, Chulalongkorn University).

Solvents for NMR spectra constituted deuterated acetone (acetone- $d_6$ ), deuterated dimethyl sulfoxide (DMSO- $d_6$ ), or deuterated chloroform ( $\text{CDCl}_3$ ). Chemical shifts were described in ppm scale using the chemical shift of the solvent as the reference signal.

## 2.4 Solvents

All organic solvents used throughout this work were of commercial grade and were redistilled before being used.

## 3. Extraction and isolation


### 3.1 Extraction and isolation of compounds from *Dendrobium braianense*

#### 3.1.1 Extraction

The dried whole plant of *Dendrobium braianense* (2 kg) was ground into a powder then mixed with MeOH solvent at room temperature. After three days the liquid extracts were taken and filtered to the flask. Then the extracts were evaporated to remove the solvent with a rotary evaporator at  $40^\circ\text{C}$  to get concentrated MeOH extract. This process was repeated 3 times with the same



volume of solvent. Finally, the extracts were weighed and stored in the fridge till their usage in the different tests (**Scheme 1**).



**Scheme 1.** Extraction steps for *Dendrobium braianense*

### 3.1.2 Isolation


#### 3.1.2.1 Isolation of compound DBra1 (Chrysotoxine) and compound DBra2 (Moscatilin)

Ethyl acetate extract (92 g) was initially divided using vacuum liquid chromatography (silica gel, hexane-EtOAc) to give five fractions (A to E). Fraction B was isolated with Sephadex LH-20 with acetone to give five fractions (B1 – B5). Fraction B2 was divided by silica gel column (EtOAc: Hexane) gradient to give five fractions B2.1 – B2.5 (**Scheme 2**).

Fraction B2.3 after drying gave compound DBra1 (97 mg) as a white powder. Compound DBra1 was later identified as chrysotoxine. Fraction B2.4 after drying gave compound DBra2 (10 mg) as brown amorphous. Compound DBra2 was then known to be moscatilin.

#### 3.1.2.2 Isolation of compound DBra3 (Gigantol)

Fraction B3 (2.38 g) was subjected to column chromatography (CC) on a silica gel column (hexane-EtOAc) (**Scheme 2**). Three subfractions (B3.1 to B3.3) were obtained. Fraction B3.2 gave brown amorphous after being left standing at room temperature overnight. The precipitates were collected, washed with EtOAc, and dried to give Compound DBra3 (170 mg). This compound was identified as gigantol.




**Scheme 2.** Isolation of compounds from ethyl acetate extract of *Dendrobium braianense*

### 3.2 Extraction and isolation of compounds from *Dendrobium kentrophyllum*

#### 3.2.1 Extraction

The dried whole plant of *Dendrobium kentrophyllum* (3.8 kg) was ground into powder then macerated with MeOH solvent at room temperature. After three days the liquid extracts were taken and filtered to the flask. Then the extracts were evaporated to remove the solvent with a rotary evaporator at 40°C to get concentrate MeOH extract. This process was repeated 3 times with the same volume of solvent. Finally, the extracts were weighed and stored in the fridge till their usage in the different tests (**Scheme 3**).



**Scheme 3.** Extraction steps for *Dendrobium kentrophyllum*


### 3.2.2 Isolation

#### 3.2.2.1 Isolation of compound DK 1 (Kaempferol) and compound DK 2 (Quercetin)


The ethyl acetate extract (72 g) was separated by vacuum-liquid chromatography (silica gel, hexane-acetone gradient) to give 7 fractions (A-H). Fraction G (27 g) was fractionated by column chromatography over silica gel (acetone-hexane, gradient) to give 11 fractions (G1-G11) (**Scheme 4**). Compound DK 1 (302 mg) was obtained from fractions G7 and G8 as a yellow powder and identified as kaempferol. Fraction G10 (1 g) was further divided by column chromatography (silica gel, toluene) to yield compound DK 2 (245 mg) as a yellow powder and identified as quercetin.

#### 3.2.2.2 Isolation of compound DK3 (Rutin)

Fraction H (2 g) was purified by column chromatography (C-18, methanol-water, gradient) to give 4 fractions (H1 – H4) (**Scheme 5**). Compound DK3 (84 mg) was obtained from H2 as a yellow powder and identified as rutin.



**Scheme 4.** Isolation of compounds from EtOAc extract of *Dendrobium kentrophyllum*



**Scheme 5.** Isolation of compounds from fraction H of *Dendrobium kentrophyllum*

#### 4. Physical and spectral data of isolated compounds

##### 4.1 Compound DBra 1 (Chrysotoxine)

Compound **DBra 1** was gained as a white powder (97 mg, 0.00485% based on dried weight of plant). It was soluble in acetone.

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in acetone- $d_6$ ; **Table 7**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 7**

##### 4.2 Compound DBra 2 (Moscatilin)

Compound **DBra 2** was gained as brown amorphous (10 mg, 0.0005% based on the dried weight of the plant). It was soluble in acetone.

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in acetone- $d_6$ ; **Table 8**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 8**

##### 4.3 Compound DBra 3 (Gigantol)

Compound **DBra 3** was gained as white crystals (170 mg, 0.0085% based on dried weight of root). It was soluble in acetone.

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in acetone- $d_6$ ; **Table 9**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 9**

##### 4.4 Compound DK 1 (Kaempferol)

Compound **DK 3** was gained as a yellow powder (302 mg, 0.00795% based on dried weight of plant). It was soluble in acetone.

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in acetone- $d_6$ ; **Table 10**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 10**

#### 4.5 Compound DK 2 (Quercetin)

Compound **DK 2** was gained as a yellow powder (245 mg, 0.00645% based on dried weight of plant). It was soluble in acetone.

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in acetone- $d_6$ ; **Table 11**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 11**

#### 4.6 Compound DK 3 (Rutin)

Compound **DK 3** was gained as a yellow powder (84 mg, 0.00221% based on dried weight of plant). It was soluble in acetone- $d_6$ .

$^1\text{H}$  NMR  $\delta$  ppm, 300 MHz, in DMSO- $d_6$ ; **Table 12**

$^{13}\text{C}$  NMR  $\delta$  ppm, 75 MHz, in acetone- $d_6$ ; **Table 12**





## 5. Assay for $\alpha$ -glucosidase inhibitory activity

The *in vitro*  $\alpha$ -glucosidase inhibition assay was performed by spectrophotometric method, involving hydrolysis of substrate *p*-nitrophenyl  $\alpha$ -D-glucopyranoside (Srianta *et al.*, 2013).

In this research,  $\alpha$ -glucosidase inhibitory activity was assessed using the spectrophotometric measurement of the amount of *p*-nitrophenol (pNP) discharged from the hydrolytic reaction of *p*-nitrophenyl- $\alpha$ -D-glucopyranoside (p-NPG) catalyzed by  $\alpha$ -glucosidase enzyme. p-NPG is a synthetic substrate representing the  $\alpha$ -linked terminal glucose of polysaccharides. The experiment was conducted at microscale *in vitro* in a 96-well plate following established protocols (Inthongkaew *et al.*, 2017).

### 5.1 Materials and instruments

- *p*-Nitrophenyl- $\alpha$ -D-glucopyranoside (p-NPG) (Sigma-Aldrich, USA)
- $\alpha$ -Glucosidase enzyme (Sigma-Aldrich, USA)
- Na<sub>2</sub>CO<sub>3</sub> (Sigma-Aldrich, USA)
- DMSO (Sigma-Aldrich, USA)
- Acarbose (Sigma-Aldrich, USA)
- Vortex mixer (Vortex-Genie2, Scientific Industries)
- Incubator (BM500, Memmert)
- Microplate reader (Perkin Elmer, BMG LABTECH)

## 5.2 Determination of $\alpha$ -glucosidase inhibitory activity

Prepare the sample 1 mg in 1 ml of 50% DMSO. Briefly, 10  $\mu$ L of the test samples and 40  $\mu$ L of  $\alpha$ -glucosidase solution (0.1 U/ml) were mixed in 96 well microplates. Incubate 96-well plate at 37°C for 10 minutes. After pre-incubation, 50  $\mu$ L of 20 Mm *p*-nitrophenol- $\alpha$ -D-glucopyranoside (p-NPG) in 50 M phosphate buffer (pH 6.8) was added to each well and incubated at 37 °C for 20 minutes. Finally, Na<sub>2</sub>CO<sub>3</sub> (100  $\mu$ L, 1 Mm) was added to each well to stop the reaction. The mixture was measured with a microplate reader at 405 nm. The  $\alpha$ -glucosidase inhibitory activity was shown to be the percentage of inhibition and was calculated as follows :

$$\% \text{ Inhibition} = \frac{(Neg - Neg') - \left(\frac{S}{A} - \frac{S'}{A'}\right)}{(Neg - Neg')}$$

Neg' = Buffer 40 $\mu$ L + 50% DMSO 10 $\mu$ L	Neg = Enzyme 40 $\mu$ L + 50% DMSO 10 $\mu$ L
A' = Buffer 40 $\mu$ L + Acarbose 10 $\mu$ L	A = Enzyme 40 $\mu$ L + Acarbose 10 $\mu$ L
S' = Buffer 40 $\mu$ L + Sample 10 $\mu$ L	S = Enzyme 40 $\mu$ L + Sample 10 $\mu$ L
Neg = Negative Control	S = Sample
A = Acarbose	' = blank ( no enzyme)

## CHAPTER IV

### RESULTS AND DISCUSSION

In this research, the extracts were prepared from *Dendrobium braianense* and *D. kentrophyllum*. They were found to have significant  $\alpha$ -glucosidase inhibitory potential and thus were subjected to further research to identify the active principles. This chapter is divided into four main sections. Each section describes the results and discussion on the phytochemical and biological studies of each plant.

#### 1. Preliminary biological activity evaluation of *D. braianense* and *D. kentrophyllum*

*Dendrobium braianense* crude extract was prepared by maceration of the dried powdered whole plants with methanol. The MeOH extract was evaluated for  $\alpha$ -glucosidase inhibitory activity and found to show 64.7% inhibition at a concentration of 50  $\mu$ g/ml.

The MeOH extract was divided using ethyl acetate, butanol, and water to give ethyl acetate and butanol extracts. These extracts were then assessed for their  $\alpha$ -glucosidase inhibitory property. The ethyl acetate extract was found to show a strong  $\alpha$ -glucosidase inhibitory effect with 82.1% inhibition at a concentration of 100  $\mu$ g/ml, whereas the butanol extract exhibited 68.6% inhibition (**Table 6**). Therefore the ethyl acetate extract was chosen for further chemical research.

*Dendrobium kentrophyllum* crude extract was prepared by maceration of the dried powdered whole plants with methanol. The MeOH extract was assessed for  $\alpha$ -glucosidase inhibitory activity and had 69.7 % inhibition at a concentration of 50  $\mu$ g/ml. The MeOH extract was divided using ethyl acetate, butanol, and water to give ethyl acetate and butanol extracts. These extracts were then assessed for their  $\alpha$ -glucosidase inhibitory property. The ethyl acetate extract was found to show a strong  $\alpha$ -glucosidase inhibitory effect with 76.4% inhibition at a concentration of 50  $\mu$ g/ml,

whereas the butanol extract exhibited 12.74 % inhibition (**Table 6**). Therefore the ethyl acetate extract was chosen for further chemical research.

**Table 6.**  $\alpha$ -glucosidase inhibitory activity of crude extracts from *D. braianense* and *D.*

*kentrophyllum*

<i>Dendrobium</i> spp.	Extracts (100 $\mu$ g/ml)	% Inhibition
<i>D. braianense</i>	Methanol Extract	64.7
	Ethyl acetate Extract	82.1
	Butanol Extract	68.6
<i>D. kentrophyllum</i>	Methanol Extract	69.7
	Ethyl acetate Extract	76.4
	Butanol Extract	12.7
Acarbose (positive control)		74.8

## 2. Structure determination of isolated compounds from *D. braianense*.

### 2.1 Identification of compound DBra1 (Chrysotoxine)

Compound DBra1 was detached as a white powder. Its HR-ESI-MS (**Appendix 1.1**) showed an  $[M+Na]^+$  ion at  $m/z$  341.1389 (calcd. For  $C_{18}H_{22}O_5Na$ , 341.1364) and the molecular formula  $C_{18}H_{22}O_5$ . The  $^1H$  NMR spectrum (**Appendix 1.2** and **Table 7**) showed a characteristic signals of bibenzyl for five aromatic protons [ $\delta$  6.49 (2H, br s, H-2, H-6), 6.71 (1H, dd,  $J = 8.1, 1.8$  Hz, H-6'), 6.82 (1H, br s, H-2'), 6.83 (1H, d,  $J = 8.1$  Hz, H-5')], four methoxy groups at  $\delta$  3.77 (9H, s) and 3.76 (3H, s), and four methylene protons at  $\delta$  2.82 (4H, br s, H<sub>2</sub>- $\alpha$  and H<sub>2</sub>- $\alpha'$ ). It was confirmed by the presence of twelve aromatic carbons and two methylene carbons at  $\delta$  38.0 (C- $\alpha$ ) and  $\delta$  (C- $\alpha'$ ) in  $^{13}C$  NMR and HSQC spectrum (**Appendix 1.3** and **1.4**). The position of H-2 and H-6 were assigned based on their HMBC correlations with C- $\alpha$  (**Appendix 1.7**). The assignments of H-2' and H-6' were based on their HMBC correlations with C- $\alpha'$ . The substitution of two symmetric methoxys at C-3 and C-5 was confirmed by their NOESY correlations with H-2 and H-6 (**Appendix 1.9**). The other two methoxys were located at C-3' and C-4', as shown by their NOESY interactions with H-2' and H-5', respectively. Compound DBra1 was known to be chrysotoxine based on the above spectral data. Its  $^1H$  and  $^{13}C$  NMR properties are in line with previously reported values (**Table 7**) (Ono *et al.*, 1995).




Figure 8. Chrysotoxine

**Table 7.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of compound DBra1 in acetone- $d_6$  and  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) of chrysotoxine in  $\text{CDCl}_3$


Position	Compound DBra1		Chrysotoxine <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$
1	-	132.2	-	132.8
2, 6	6.49 (s)	106.0	6.36 (s)	105.2
3, 5	-	147.6	-	146.8
4	-	134.1	-	132.8
$\alpha$	2.82 (br s)	38.0	2.83 (s)	38.3
$\alpha'$	2.82 (br s)	37.6	2.83 (s)	37.8
1'	-	134.7	-	134.3
2'	6.82 (br s)	112.7	6.66 (d, 1.8)	111.9
3'	-	147.8	-	147.2
4'	-	149.3	-	148.7
5'	6.83 (d, 8.1)	112.0	6.79 (d, 8.1)	111.2
6'	6.71 (dd, 8.1, 1.8)	120.4	6.70 (dd, 1.8, 8.1)	120.4
MeO-3, MeO-5	3.77 (s)	55.7	3.84 (s)*	56.2
MeO-3'	3.77 (s)	55.3*	3.85 (s)*	55.9*
MeO-4'	3.76 (s)	55.1*	3.84 (s)*	55.8*

<sup>a</sup> (Ono *et al.*, 1995) \* Value in the same column are interchangeable.

## 2.2 Identification of compound DBra2 (Moscatilin)

Compound DBra2 was gained as a brown amorphous solid. The HR-ESI-MS of this compound showed an  $[M+Na]^+$  ion at  $m/z$  327.1216 (calcd. for  $C_{17}H_{20}O_5Na$ , 327.1208), corresponding to the molecular formula  $C_{17}H_{20}O_5$  (**Appendix 2.1**). The  $^1H$  and  $^{13}C$  NMR spectrum (**Appendix 2.2, 2.3** and **Table 8**) of compound DBra2 were similar to compound DBra1, except for the substitution of a hydroxyl instead of a methoxyl at C-4'. The  $^1H$  NMR revealed the signals of five aromatic protons at  $\delta$  6.49 (2H, br s, H-2, H-6), 6.65 (1H, dd,  $J = 8.1, 1.5$ , H-6'), 6.72 (1H, d,  $J = 8.1$  Hz, H-5'), 6.79 (1H, d,  $J = 1.5$  Hz, H-2'), three methoxyls at 3.77 (6H, s, MeO-3,5) and 3.79 (3H, s, MeO-3'), and two methylene protons at  $\delta$  2.79 (4H, br s, H<sub>2</sub>- $\alpha$  and H<sub>2</sub>- $\alpha'$ ). The  $^{13}C$  NMR spectrum (**Appendix 2.3**) exhibited 14 carbon signals of 17 carbons.

Compound DBra2 was identified as moscatilin based on the above spectral evidence and by comparing it with prior reported data (**Table 8**) (Klongkumnuankarn *et al.*, 2015).



**Figure 9.** Moscatilin

**Table 8.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of compound DBra2 in acetone- $d_6$  and  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) of moscatilin in acetone- $d_6$

Position	Compound DBra2		Moscatilin <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$
1	-	132.3	-	133.1
2, 6	6.49 (br s)	106.0	6.48 (s)	106.7
3, 5	-	147.6	-	148.3
4	-	134.1	-	134.8
$\alpha$	2.79 (br s)	38.2	2.78 (m)	38.3
$\alpha'$	2.79 (br s)	37.6	2.78 (m)	38.8
1'	-	133.3	-	134.1
2'	6.79 (d, 1.8)	112.1	6.78 (d, 2.0)	112.9
3'	-	147.2	-	147.9
4'	-	144.7	-	145.3
5'	6.72 (d, 7.8)	114.7	6.75 (d, 8.0)	115.4
6'	6.65 (dd, 8.1, 1.8)	120.8	6.64 (dd, 8.0, 2.0)	121.6
MeO-3, MeO-5	3.77 (s)	55.7	3.75 (s)	56.5
MeO-3'	3.79 (s)	55.3	3.76 (s)	56.1


<sup>a</sup> (Klongkumnuankarn *et al.*, 2015)



### 2.3 Identification of compound DBra3 (Gigantol)

Compound DBra3, a brown amorphous solid, possesses a molecular formula  $C_{16}H_{18}O_4$ , as suggested by an  $[M+Na]^+$  ion at  $m/z$  297.1113 (calcd. For  $C_{16}H_{18}O_4Na$ , 297.1102) in the HR-ESI-MS (**Table 9**). The  $^1H$  NMR (**Appendix 3.2, Table 9**) showed signals for two methylene protons at  $\delta$  2.76 (4H, br s,  $H_2-\alpha$  and  $H_2-\alpha'$ ). The  $^1H$  NMR also displayed six aromatic proton signals at  $\delta$  6.26 (1H, br d,  $J = 2.1$  Hz, H-4), 6.31 (1H, br s, H-2), 6.33 (1H, br s, H-6), 6.66 (1H, dd,  $J = 8.1, 1.5$  Hz, H-6'), 6.74 (1H, d,  $J = 8.1$  Hz, H-5') and 6.80 (1H, d,  $J = 1.5$  Hz, H-2'), and two methoxyl protons at  $\delta$  3.70 (3H, s, MeO-3) and 3.79 (3H, s, MeO-3'). The  $^{13}C$  (**Appendix 3.3**) exhibited 16 carbon signals, corresponding to six quaternary carbons, six methines, two methylenes, and two methoxyls (**Appendix 3.3**).

This compound was identified as gigantol based on above spectral evidence. Its  $^1H$  and  $^{13}C$  NMR data are in line with previously reported data (**Table 9**) (Klongkumnuankarn *et al.*, 2015).



**Figure 10.** Gigantol

**Table 9.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of compound DBra3 in acetone- $d_6$  and  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) of gigantol in acetone- $d_6$

Position	Compound DBra3		Gigantol <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$
1	-	144.6	-	145.4
2	6.31 (br s)	108.1	6.22 (t, 2.0)	108.8
3	-	160.9	-	159.1
4	6.26 (t, 2.1)	98.9	6.28 (t, 2.0)	99.6
5	-	158.4	-	161.7
6	6.33 (br s)	105.4	6.30 (t, 2.0)	106.2
$\alpha$	2.76 (s)	38.2	2.78 (m)	39.0
$\alpha'$	2.76 (s)	37.1	2.78 (m)	37.9
1'	-	133.3	-	134.0
2'	6.80 (d, 1.5)	112.0	6.79 (d, 1.5)	115.4
3'	-	147.2	-	147.9
4'	-	144.3	-	145.1
5'	6.74 (d, 8.1)	114.7	6.69 (d, 8.0)	112.8
6'	6.66 (dd, 8.1, 1.5)	120.7	6.64 (dd, 8.0, 1.5)	121.5
MeO-3	3.70 (3H, s)	54.4	3.69 (s)	55.2
MeO-3'	3.79 (3H, s)	55.3	3.78 (s)	56.0

<sup>a</sup> (Klongkumnuankarn *et al.*, 2015)

### 3. Structure determination of isolated compounds from *D. kentrophyllum*

#### 3.1 Identification of compound DK1 (Kaempferol)

Compound DK1 was gained as a yellow powder. The molecular formula of  $C_{15}H_{10}O_6$  was analyzed from its  $[M+H]^+$  ion at 287.0545 (calcd. for  $C_{15}H_{11}O_6$  287.0555) in the HR-APCI-MS. The  $^1H$  and  $^{13}C$  NMR spectra (**Appendix 4.1 & 4.2**) of this compound exhibited the characteristics of flavonoid nucleus by a sharp singlet proton signal at  $\delta$  12.17 (chelated hydroxyl) and the carbon signal at  $\delta$  175.7 (C-4). The  $^1H$  NMR showed two doublets at  $\delta$  6.26 (1H, d,  $J = 1.8$  Hz) and  $\delta$  6.52 (1H, d,  $J = 1.8$  Hz) assignable to H-6 and H-8 of ring A, respectively. The assignment of H-6 was based on the HMBC of H-6 with C-5 ( $\delta$  161.4) (**Appendix 4.6**). It also showed a pair of doublets, 2H each at  $\delta$  7.02 (2H, d,  $J = 9.0$  Hz, H-3' and H-5') and 8.14 (2H, d,  $J = 9.0$  Hz, H-2' and H-6'), suggesting the *para*-hydroxyl substitution of ring B. The  $^{13}C$  NMR and HSQC spectra (**Appendix 4.2 & 4.4**) displayed 13 signals corresponding to 15 carbons including nine quaternary carbons and six methines. By comparing the above spectroscopic data with reported values (Lin *et al.*, 2016). Thus, compound DK1 was known to be kaempferol (**Table 10**).




Figure 11. Kaempferol


**Table 10.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of DK1 in acetone- $d_6$  and  $^1\text{H}$  NMR (600 MHz) and  $^{13}\text{C}$  NMR (150 MHz) of kaempferol in DMSO- $d_6$

Position	Compound DK1		Kaempferol <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$
2	-	146.1	-	146.8
3	-	135.7	-	135.6
4	-	175.7	-	175.9
5	-	161.4	-	160.7
6	6.26 (d, 1.8)	98.3	6.19 (d, 1.8)	98.2
7	-	164.1	-	164.9
8	6.52 (d, 1.8)	93.6	6.44 (d, 1.8)	93.5
9	-	156.9	-	156.2
10	-	103.3	-	103.0
1'	-	122.4	-	121.7
2'	8.14 (d, 9.0)	129.6	8.04 (d, 9.0)	130.5
3'	7.02 (d, 9.0)	115.4	6.92 (d, 9.0)	115.4
4'	-	159.3	-	159.2
5'	7.02 (d, 9.0)	115.4	6.92 (d, 9.0)	115.4
6'	8.14 (d, 9.0)	129.6	8.04 (d, 9.0)	130.5
HO-5	12.17 (s)	-	-	-

<sup>a</sup> Lin LJ, Huang XB, Lv ZC. Isolation and identification of flavonoids components from *Pteris vittata* L. 2016; 5: 1649.

### 3.2 Identification of compound DK2 (Quercetin)

Compound DK2 was detached as a yellow powder. The molecular formula was  $C_{15}H_{10}O_7$  by HR-APCI-MS of its  $[M+H]^+$  ion at 303.0492 (calcd. for  $C_{15}H_{11}O_7$  303.0504). The  $^1H$  and  $^{13}C$  NMR data were the same as those of **1**, except for the presence of a hydroxyl group at C-3' of ring B. This was confirmed by the presence of the ABM spin system of  $^1H$  NMR spectrum at  $\delta$  7.00 (1H, d,  $J = 8.4$  Hz, H-5'),  $\delta$  7.69 (1H, dd,  $J = 8.4, 2.1$  Hz, H-6') and 7.82 (1H, d,  $J = 2.1$  Hz, H-2'). The  $^1H$  NMR also exhibited two doublets of H-6 (1H,  $J = 1.8$  Hz) and H-8 (1H,  $J = 1.8$  Hz), and a sharp singlet of 5-OH at  $\delta$  12.17. The  $^{13}C$  NMR (Appendix 5.2) displayed 15 signals corresponding to ten quaternary carbons and five methines. Compound DK2 was identified to be quercetin by comparing with prior reported data (**Table 11**) (Lin *et al.*, 2016).



**Figure 12.** Quercetin

**Table 11.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of compound DK2 in acetone- $d_6$  and  $^1\text{H}$  NMR (600 MHz) and  $^{13}\text{C}$  NMR (150 MHz) of quercetin in DMSO- $d_6$

Position	Compound DK2		Quercetin <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$
2	-	146.1	-	146.7
3	-	135.9	-	135.6
4	-	175.7	-	175.8
5	-	161.4	-	160.4
6	6.26 (d, 1.8)	98.3	6.19 (d, 1.8)	98.1
7	-	164.1	-	163.8
8	6.51 (d, 1.8)	93.6	6.40 (d, 2.4)	93.3
9	-	156.9	-	156.1
10	-	103.2	-	102.9
1'	-	122.8	-	121.8
2'	7.82 (d, 2.1)	114.8	7.67 (d, 2.4)	114.9
3'	-	144.9	-	144.9
4'	-	147.5	-	147.6
5'	7.00 (d, 8.4)	115.3	6.88 (d, 8.4)	115.5
6'	7.69 (dd, 8.4, 2.1)	120.6	7.54 (dd, 8.4, 2.4)	119.9
HO-5	12.17 (s)	-	-	-

<sup>a</sup> Lin LJ, Huang XB, Lv ZC. Isolation and identification of flavonoids components from *Pteris vittata* L. 2016; 5: 1649.

### 3.3 Identification of compound DK3 (Rutin)

Compound DK3, a yellow powder, showed its  $[M+Na]^+$  ion at 633.1436 (calcd. for  $C_{27}H_{30}O_{16}Na$  633.1431) in the HR-ESI-MS, suggesting a molecular formula  $C_{27}H_{30}O_{16}$ . The  $^1H$  NMR spectrum (**Appendix 6.1**) showed the presence of the ABM splitting system of ring B at  $\delta$  7.52 (1H, br s, H-2'),  $\delta$  7.54 (1H, br d,  $J = 8.7$  Hz, H-6') and  $\delta$  6.82 (1H, d,  $J = 8.7$  Hz, H-5') and two broad singlets of H-6 and H-8 at  $\delta$  6.18 and  $\delta$  6.37. In addition, the appearance of two anomeric proton signals at  $\delta$  5.33 (1H, d,  $J = 6.9$  Hz, H-1'') and  $\delta$  4.37 (1H, s, H-1'''), which revealed HSQC correlations with the carbon at  $\delta$  101.6 (C-1'') and  $\delta$  101.2 (C-1'''), respectively, suggest that this compound should be a diglycoside of compound DK2. The  $^{13}C$  NMR and HSQC spectra (**Appendix 6.5**) exhibited 12 carbon signals of sugar carbons and 15 carbon signals of the quercetin nucleus. The sugars were identified as rutinose ( $\alpha$ -rhamnopyranosyl- $\beta$ -glucopyranose) by comparison with reported values (Sintayehu *et al.*, 2012). The connection of glucose and rhamnose was maintained by the HMBC correlations of C-1''' with H-6'' (**Appendix 6.8**). The rutinose was located at C-3 of quercetin based on the HMBC correlations of C-3 ( $\delta$  133.7) with H-1''. Compound DK3 was determined as quercetin-3-O-rutinoside or rutin by comparing with prior reported data (**Table 9**) (Sintayehu *et al.*, 2012).




Figure 13. Rutin

**Table 12.**  $^1\text{H}$  NMR (300 MHz) and  $^{13}\text{C}$  NMR (75 MHz) spectral data of compound DK3 in acetone- $d_6$  and  $^1\text{H}$  NMR (400 MHz) and  $^{13}\text{C}$  NMR (100 MHz) of rutin in DMSO- $d_6$

Position	Compound DK3		Rutin <sup>a</sup>	
	$\delta_{\text{H}}$ (mult., $J$ in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (mult., $J$ in	$\delta_{\text{C}}$
2	-	156.9	-	157.1
3	-	133.7	-	133.7
4	-	177.8	-	177.8
5	-	161.7	-	161.6
6	6.18 (br s)	99.1	6.20 (d, 1.6)	99.1
7	-	164.6	-	164.5
8	6.37 (br s)	94.0	6.39 (d, 1.6)	94.1
9	-	157.0	-	156.9
10	-	104.4	-	104.4
1'	-	121.6	-	121.6
2'	7.52 (br s)	116.7	7.52 (s)	115.7
3'	-	145.2	-	145.2
4'	-	148.9	-	148.8
5'	6.82 (d, 8.7)	115.7	6.85 (d, 8.4)	116.7
6'	7.54 (br d, 8.7)	122.0	7.55 (d, 8.4)	122.1
HO-5	12.59 (s)	-	12.56 (s)	-
1''	5.33 (d, 6.9)	101.6	5.33 (d)	101.6
2''	3.22-3.28 (m)	74.5	3.32-3.81	74.5
3''	3.22-3.28 (m)	76.9	3.32-3.81	76.9
4''	3.22-3.28 (m)	71.0	3.32-3.81	71.1
5''	3.22-3.28 (m)	76.4	3.32-3.81	76.3
6''	3.68, 3.29 (m)	67.4	3.32-3.81	67.4
1'''	4.37 (s)	101.2	4.38 (s)	101.2
2'''	3.22-3.28 (m)	70.8	3.32-3.81	70.8
3'''	3.03-3.09 (m)	70.4	3.32-3.81	70.4
4'''	3.03-3.09 (m)	72.3	3.32-3.81	72.3
5'''	3.22-3.28 (m)	68.7	3.32-3.81	68.7
6'''	0.98 (d, 6.0)	18.2	1.11 (d, 6.0)	18.2

<sup>a</sup> Sintayehu B, Asres K, Raghavendra Y. Radical scavenging activities of the leaf extracts and a flavonoid glycoside isolated from *Cineraria abyssinica* Sch. Bip. Exa. Rich. Journal of Applied Pharmaceutical Science. 2012; 2: 44-49.



#### 4. Biological activity of isolated compounds of *Dendrobium braianense* and *Dendrobium kentrophyllum*

In this study, the isolated compounds were assessed for  $\alpha$ -glucosidase inhibitory activity (Table 13). Chrysotoxine and moscatilin did not possess  $\alpha$ -glucosidase inhibitory activity (28.71% and 14.44 % inhibition at 100  $\mu$ g/ml), in line with prior research (Inthongkaew, *et al.*, 2013). Gigantol exhibited a stronger inhibitory effect ( $IC_{50}$  349  $\mu$ M) than acarbose ( $IC_{50}$  532.4  $\mu$ M). These findings show that those compounds have  $\alpha$ -glucosidase inhibitory activity compared with earlier reported values. Gigantol ( $IC_{50}$  79.87  $\mu$ M) than acarbose ( $IC_{50}$  724.74  $\mu$ M) (San, H. T., *et al.*, 2011). Kaempferol ( $IC_{50}$  42.59  $\mu$ M) and Quercetin ( $IC_{50}$  30,67  $\mu$ M), than acarbose ( $IC_{50}$  177,5  $\mu$ M) (Habtemariam, S., 2011). Rutin did not possess  $\alpha$ -glucosidase inhibitory activity which is in agreement with a previous report (Habtemariam, S., 2011). Kaempferol and quercetin exhibited strong inhibitory effects with  $IC_{50}$  of 87.5 and 109.1  $\mu$ M compared with acarbose. These findings were in agreement with previously reported values (Habtemariam, S., 2011).

**Table 13.**  $\alpha$ -Glucosidase inhibitory activity of *D. braianense* and *D. kentrophyllum*

compounds	$IC_{50}$ ( $\mu$ M)
DBra 1 (Chrysotoxine)	NA
DBra 2 (Moscatilin)	NA
DBra 3 (Gigantol)	349.0 $\pm$ 9.3
DK 1 (Kaempferol)	87.5 $\pm$ 6.2
DK 2 (Quercetin)	109.1 $\pm$ 7.6
DK 3 (Rutin)	NA
Acarbose	532.4 $\pm$ 19.6

## CHAPTER V

### CONCLUSION



The ethyl acetate extracts prepared from the whole plant of *Dendrobium braianense* and *D. kentrophyllum* exhibited  $\alpha$ -glucosidase inhibitory activity 82.1% and 76.4%, respectively, at a concentration of 100  $\mu\text{g/mL}$ . Phytochemical research of the ethyl acetate extract of *D. braianense* led to the isolation of three bibenzyls, which included chrysotoxine, moscatilin, and gigantol. Chromatographic separation of the ethyl acetate extract of *Dendrobium kentrophyllum* led to the isolation of three flavonoids, namely kaempferol, quercetin, and rutin. The structures of these isolates were determined utilizing the analysis of their  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, and MS data and compared with the prior reported values. All the isolated compounds were assessed for their  $\alpha$ -glucosidase inhibitory activity. Gigantol, kaempferol, and quercetin were found to exhibit strong  $\alpha$ -glucosidase inhibitory activity with  $\text{IC}_{50}$  349.0, 87.5, and 109.1  $\mu\text{M}$ , respectively, in comparison with positive control acarbose ( $\text{IC}_{50}$  532.4  $\mu\text{M}$ ). These findings show that those compounds have  $\alpha$ -glucosidase inhibitory activity compared with earlier reported values. Gigantol ( $\text{IC}_{50}$  79.87  $\mu\text{M}$ ) than acarbose ( $\text{IC}_{50}$  724.74  $\mu\text{M}$ ) (San, H. T., *et al.*, 2011). Kaempferol ( $\text{IC}_{50}$  42.59  $\mu\text{M}$ ) and Quercetin ( $\text{IC}_{50}$  30,67  $\mu\text{M}$ ), than acarbose ( $\text{IC}_{50}$  177,5  $\mu\text{M}$ ) (Habtemariam, S., 2011).

Chrysotoxine, moscatilin, and rutin were devoid of activity. Finally, the writer suggest for further experiments to broaden scientific information.


## APPENDICES

## Appendix 1: The spectral data of compound DBra 1


## 1.1 Mass spectrum of compound DBra 1

1.2  $^1\text{H}$ -NMR (300 MHz) spectrum of compound DBra 1 (Acetone- $d_6$ )


### 1.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 1 (Acetone- $d_6$ )




### 1.4 HSQC spectrum of compound DBra 1 (Acetone- $d_6$ ) expansion $[\delta\text{H } 2.82 - 6.83 \text{ ppm}, \delta\text{C } 37.6 - 120.4]$



1.5 HSQC spectrum of compound DBra 1 (Acetone- $d_6$ )  
expansion [ $\delta_H$  6.49 – 6.83 ppm,  $\delta_C$  106.0 – 120.4]




1.6 HMBC spectrum of compound DBra 1 (Acetone- $d_6$ )  
expansion [ $\delta_H$  2.82 – 6.83 ppm,  $\delta_C$  37.6 – 149.3 ppm]






### 1.7 HMBC spectrum of compound DBra 1 (Acetone- $d_6$ )



expansion [ $\delta_H$  2.82 – 6.83 ppm,  $\delta_C$  106.0 – 149.3 ppm]



### 1.8 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ )


expansion [ $\delta_H$  2.82 – 3.77 ppm (F1),  $\delta_H$  6.49 – 6.83 ppm (F2)]




1.9 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ )expansion [ $\delta_H$  2.82 – 6.83 ppm]1.10 NOESY spectrum of compound DBra 1 (Acetone- $d_6$ )expansion [ $\delta_H$  6.49 – 6.83 ppm]

## Appendix 2: The spectral data of compound DBra 2

### 2.1 Mass spectrum of compound DBra 2




### 2.2 <sup>1</sup>H-NMR (300 MHz) spectrum of compound DBra 2 (Acetone-d<sub>6</sub>)






### 2.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 2 (Acetone- $d_6$ )




## Appendix 3: The spectral data of compound DBra 3


### 3.1 Mass spectrum of compound DBra 3



### 3.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DBra 3 (Acetone- $d_6$ )




### 3.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DBra 3 (Acetone- $d_6$ )




## Appendix 4: The spectral data of compound DK 1


### 4.1. Mass spectrum of compound DK 1



### 4.2 $^1\text{H}$ -NMR (300 MHz) spectrum of compound DK 1 (Acetone- $d_6$ )






#### 4.3 $^{13}\text{C}$ -NMR (300 MHz) spectrum of compound DK 1 (Acetone- $d_6$ )



#### 4.4 HSQC NMR spectrum of compound DK 1 (Acetone- $d_6$ )


expansion [ $\delta\text{H}$  6.26 – 8.14 ppm,  $\delta\text{C}$  93.6 – 129.6 ppm]




4.5 HMBC NMR spectrum of compound DK 1 (Acetone- $d_6$ )expansion [ $\delta_H$  6.26 – 12.17 ppm,  $\delta_C$  93.6 – 164.1 ppm]4.6 HMBC NMR spectrum of compound DK 1 (Acetone- $d_6$ )expansion [ $\delta_H$  6.26 – 12.17 ppm,  $\delta_C$  93.6 – 164.1 ppm]

## Appendix 5: The spectral data of compound DK 2


## 5.1 Mass spectrum of compound DK 2



### 5.2 $^1\text{H}$ -NMR (500 MHz) spectrum of compound DK 2 (Acetone- $d_6$ )





### 5.3 $^{13}\text{C}$ -NMR (75 MHz) spectrum of compound DK 2 (Acetone- $d_6$ )






## Appendix 6: The spectral data of compound DK 3

## 6.1 Mass spectrum of compound DK 3

6.2  $^1\text{H}$ -NMR (300 MHz) spectrum of compound DK 3 ( $\text{DMSO}-d_6$ )




### 6.3 $^{13}\text{C}$ -NMR (300 MHz) spectrum of compound DK 3 (DMSO- $d_6$ )




### 6.4 HSQC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta\text{H}$  0.98 – 4.37 ppm,  $\delta\text{C}$  67.4 – 76.9 ppm]




### 6.5 HSQC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta_H$  4.37 – 7.54 ppm,  $\delta_C$  94.0 – 122.0 ppm]




### 6.6 HSQC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta_H$  67.4 – 76.9 ppm]




### 6.6 HMBC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta\text{H}$  0.98 – 12.59 ppm,  $\delta\text{C}$  18.2 – 177.8 ppm]



### 6.7 HMBC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta\text{H}$  3.22 – 12.59 ppm,  $\delta\text{C}$  94.0 – 177.8 ppm]




### 6.8 HMBC spectrum of compound DK 3 (Acetone- $d_6$ )


expansion [ $\delta\text{H}$  0.98 – 4.37 ppm,  $\delta\text{C}$  67.4 – 76.9 ppm]



### 6.9 HMBC spectrum of compound DK 3 (Acetone- $d_6$ )

expansion [ $\delta\text{H}$  0.98 – 7.54 ppm]



6.10 HMBC spectrum of compound DK 3 (Acetone- $d_6$ )expansion [ $\delta$ H 3.03 – 5.33 ppm]

## REFERENCES



จุฬาลงกรณ์มหาวิทยาลัย  
**CHULALONGKORN UNIVERSITY**

## REFERENCES

- Ahamad, J., Naquvi, K. J., Mir, S. R., Ali, M., & Shuaib, M. (2011). Review on role of natural Alpha-Glucosidase inhibitors for Management of Diabetes Mellitus. International Journal of Biomedical Research, 2(6), 374-380.
- Bi, Z. M., Wang, Z. T., & Xu, L. S. (2004). Chemical constituents of *Dendrobium moniliforme*. Acta Botanica Sinica, 46(1), 124-126.
- Bulpitt, C. J., Li, Y., Bulpitt, P. F., & Wang, J. (2007). The use of orchids in Chinese medicine. Journal of the Royal Society of Medicine, 100(12), 558-563.
- Cai, J., Yang, C., Chen, T., & Zhao, L. (2018). Detection of new phenylpropanoids from *Dendrobium chrysanthum*. Natural Product Research, 32(13), 1600-1604. doi:10.1080/14786419.2017.1385006
- Chang, C. C., Ku, A. F., Tseng, Y. Y., Yang, W. B., Fang, J. M., & Wong, C. H. (2010). 6,8-di-C-glycosyl flavonoids from *Dendrobium huoshanense*. Journal of Natural Products, 73(2), 229-232. doi:10.1021/np900252f
- Chang, S. J., Lin, T. H., & Chen, C. C. (2001). Constituents from the stems of *Dendrobium clavatum* var. *aurantiacum*. Journal of Chinese Medicine, 12(3), 211-218.
- Chanvorachote, P., Kowitdamrong, A., Ruanghirun, T., Sritularak, B., Mungmee, C., & Likhitwitayawuid, K. (2013). Anti-metastatic activities of bibenzyls from *Dendrobium pulchellum*. Natural product communications, 8, 115-118.
- Chen, C. C., Wu, L. G., Ko, F. N., & Teng, C. M. (1994). Antiplatelet aggregation principles of *Dendrobium loddigesii*. Journal of Natural Products, 57(9), 1271-1274. doi:10.1021/np50111a014
- Chen, X. J., Mei, W. L., Cai, C. H., Guo, Z. K., Song, X. Q., & Dai, H. F. (2014). Four new bibenzyl derivatives from *Dendrobium sinense*. Phytochemistry Letters, 9, 107-112. doi:10.1016/j.phytol.2014.04.012
- Chen, X. J., Mei, W. L., Zuo, W. J., Zeng, Y. B., Guo, Z. K., Song, X. Q., & Dai, H. F. (2013). A new antibacterial phenanthrenequinone from *Dendrobium sinense*. Journal of Asian Natural Products Research, 15(1), 67-70. doi:10.1080/10286020.2012.740473

- Chen, Y., Li, J., Wang, L., & Liu, Y. (2008). Aromatic compounds from *Dendrobium aphyllum*. Biochemical Systematics and Ecology, 36(5-6), 458-460. doi:10.1016/j.bse.2007.11.004
- Chen, Y., Liu, Y., Jiang, J., Zhang, Y., & Yin, B. (2008). Dendronone, a new phenanthrenequinone from *Dendrobium cariniferum*. Food Chemistry, 111(1), 11-12. doi:10.1016/j.foodchem.2008.03.017
- Chen, Y. G., Yu, H., & Liu, Y. (2014). Chemical constituents from *Dendrobium brymerianum* Rchb. f. Biochemical Systematics and Ecology, 57, 175-177. doi:10.1016/j.bse.2014.08.027
- Choudhury, H., Pandey, M., Hua, C. K., Mun, C. S., Jing, J. K., Kong, L., Yee, T. S. (2018). An update on natural compounds in the remedy of diabetes mellitus: A systematic review. Journal of traditional and complementary medicine, 8(3), 361-376.
- Chuakul, W. (2002). Ethnomedical uses of Thai orchidaceous plants. Mahidol Univ J Pharm Sci, 29(3-4), 41-45.
- Dahmen, J., & Leander, K. (1978). Amotin and amoenin, two sesquiterpenes of the picrotoxane group from *Dendrobium amoenum*. Phytochemistry, 17(11), 1949-1952. doi:10.1016/S0031-9422(00)88740-3
- Dong, H.-Q., Li, M., Zhu, F., Liu, F.-L., & Huang, J.-B. (2012). Inhibitory potential of trilobatin from *Lithocarpus polystachyus* Rehd against  $\alpha$ -glucosidase and  $\alpha$ -amylase linked to type 2 diabetes. Food Chemistry, 130(2), 261-266.
- Fan, C., Wang, W., Wang, Y., Qin, G., & Zhao, W. (2001). Chemical constituents from *Dendrobium densiflorum*. Phytochemistry, 57(8), 1255-1258. doi:10.1016/s0031-9422(01)00168-6
- Fan, W. W., Xu, F. Q., Dong, F. W., Li, X. N., Li, Y., Liu, Y. Q., Hu, J. M. (2013). Dendrowardol C, a novel sesquiterpenoid from *Dendrobium wardianum* Warner. Natural Products and Bioprospecting, 3(3), 89-92. doi:10.1007/s13659-013-0024-9
- Fatimah, R. N. (2015). Diabetes melitus tipe 2. Jurnal Majority, 4(5).



- Gawell, L., & Leander, K. (1976). The constitution of aduncin, a sesquiterpene related to picrotoxinin, found in *Dendrobium aduncum*. Phytochemistry, 15(12), 1991-1992. doi:10.1016/s0031-9422(00)88869-x
- Gong, C. Y., Lu, B., Yang, L., Wang, L., & Ji, L.-L. (2013). Bibenzyl from *Dendrobium* inhibits angiogenesis and its underlying mechanism. Acta pharmaceutica Sinica, 48(3), 337-342.
- Habtemariam, S. (2011).  $\alpha$ -Glucosidase inhibitory activity of kaempferol-3-O-rutinoside. Natural Product Communications, 6(2), 1934578X1100600211
- He, L., Su, Q., Bai, L., Li, M., Liu, J., Liu, X., Shi, J. (2020). Recent research progress on natural small molecule bibenzyls and its derivatives in *Dendrobium* species. European Journal of Medicinal Chemistry, 204, 112530.
- Hemscheidt, T., & Spenser, I. D. (1991). Biosynthesis of anosmine, an imidazole alkaloid of the orchid *Dendrobium parishii*. Journal of the Chemical Society, Chemical Communications (7), 494. doi:10.1039/c39910000494
- Honda, C., & Yamaki, M. (2000). Phenanthrenes from *Dendrobium plicatile*. Phytochemistry, 53(8), 987-990. doi:10.1016/S0031-9422(99)00497-5
- Hossain, M. M. (2011). Therapeutic orchids: traditional uses and recent advances—an overview. Fitoterapia, 82(2), 102-140.
- Hu, J. M., Chen, J. J., Yu, H., Zhao, Y. X., & Zhou, J. (2008a). Five new compounds from *Dendrobium longicornu*. Planta Medica, 74(5), 535-539. doi:10.1055/s-2008-1074492
- Hu, J. M., Chen, J. J., Yu, H., Zhao, Y. X., & Zhou, J. (2008b). Two novel bibenzyls from *Dendrobium trigonopus*. Journal of Asian Natural Products Research, 10(7-8), 653-657. doi:10.1080/10286020802133605
- Hu, J. M., Fan, W. W., Dong, F. W., Miao, Z. H., & Zhou, J. (2012). Chemical components of *Dendrobium chrysotoxum*. Chinese Journal of Chemistry, 30(6), 1327-1330. doi:10.1002/cjoc.201100670
- Hu, J. M., Zhao, Y. X., Miao, Z. H., & Zhou, J. (2009). Chemical components of *Dendrobium polyanthum*. Bulletin of the Korean Chemical Society, 30(9), 2098-2100. doi:10.5012/bkcs.2009.30.9.2098

- Hwang, J. S., Lee, S. A., Hong, S. S., Han, X. H., Lee, C., Kang, S. J., Hwang, B. Y. (2010). Phenanthrenes from *Dendrobium nobile* and their inhibition of the LPS-induced production of nitric oxide in macrophage RAW 264.7 cells. Bioorganic & Medicinal Chemistry Letters, 20(12), 3785-3787. doi:10.1016/j.bmcl.2010.04.054
- Inthongkaew, P., Chatsumpun, N., Supasuteekul, C., Kitisripanya, T., Putalun, W., Likhitwitayawuid, K., & Sritularak, B. (2017).  $\alpha$ -Glucosidase and pancreatic lipase inhibitory activities and glucose uptake stimulatory effect of phenolic compounds from *Dendrobium formosum*. Revista brasileira de farmacognosia, 27(4), 480-487.
- Ito, M., Matsuzaki, K., Wang, J., Daikonya, A., Wang, N. L., Yao, X. S., & Kitanaka, S. (2010). New phenanthrenes and stilbenes from *Dendrobium loddigesii*. Chemical and Pharmaceutical Bulletin, 58(5), 628-633. doi:10.1248/cpb.58.628.
- Joshi, S. R., Standl, E., Tong, N., Shah, P., Kalra, S., & Rathod, R. (2015). Therapeutic potential of  $\alpha$ -glucosidase inhibitors in type 2 diabetes mellitus: an evidence-based review. Expert opinion on pharmacotherapy, 16(13), 1959-1981.
- Kashima, Y., Yamaki, H., Suzuki, T., & Miyazawa, M. (2013). Structure-activity relationships of bergenin derivatives effect on  $\alpha$ -glucosidase inhibition. Journal of enzyme inhibition and medicinal chemistry, 28(6), 1162-1170.
- Khuraijam, J. S., & Roy, R. K. (2016). *Dendrobium kentrophyllum* (Orchidaceae)—a new record for the orchid flora of Manipur, India. Richardiana, 17, 273-276.
- Kim, J. H., Oh, S. Y., Han, S. B., Uddin, G. M., Kim, C. Y., & Lee, J. K. (2015). Anti-inflammatory effects of *Dendrobium nobile* derived phenanthrenes in LPS-stimulated murine macrophages. Archives of Pharmacal Research, 38(6), 1117-1126. doi:10.1007/s12272-014-0511-5
- Klongkumnuankarn, P., Busaranon, K., Chanvorachote, P., Sritularak, B., Jongbunprasert, V., & Likhitwitayawuid, K. (2015). Cytotoxic and antimigratory activities of phenolic compounds from *Dendrobium brymerianum*. Evidence-

- based Complementary and Alternative Medicine, 2015, 1-9.  
doi:10.1155/2015/350410
- Kongkatitham, V., Muangnoi, C., Kyokong, N., Thaweesest, W., Likhitwitayawuid, K., Rojsitthisak, P., & Sritularak, B. (2018). Anti-oxidant and anti-inflammatory effects of new bibenzyl derivatives from *Dendrobium parishii* in hydrogen peroxide and lipopolysaccharide treated RAW264.7 cells. Phytochemistry Letters, 24, 31-38. doi:10.1016/j.phytol.2018.01.006
- Kumar, S., & Pandey, A. K. (2013). Chemistry and biological activities of flavonoids: an overview. The scientific world journal, 2013.
- Kyokong, N., Muangnoi, C., Thaweesest, W., Kongkatitham, V., Likhitwitayawuid, K., Rojsitthisak, P., & Sritularak, B. (2019). A new phenanthrene dimer from *Dendrobium palpebrae*. Journal of Asian Natural Products Research, 21(4), 391-397. doi:10.1080/10286020.2018.1429416
- Li, C. B., Wang, C., Fan, W. W., Dong, F. W., Xu, F. Q., Wan, Q. L., Zhou, J. (2013). Chemical components of *Dendrobium crepidatum* and their neurite outgrowth enhancing activities. Natural Products and Bioprospecting, 3(2), 70-73. doi:10.1007/s13659-012-0103-3
- Li, J. T., Yin, B. L., Liu, Y., Wang, L. Q., & Chen, Y. G. (2009). Mono-aromatic constituents of *Dendrobium longicornu*. Chemistry of Natural Compounds, 45(2), 234-236. doi:10.1007/s10600-009-9293-2
- Li, X.-h., Guo, L., Yang, L., Peng, C., He, C.-j., Zhou, Q.-m., Zhang, T.-m. (2014). Three new neolignan glucosides from the stems of *Dendrobium aurantiacum* var. *denneanum*. Phytochemistry Letters, 9, 37-40.
- Li, Y., Wang, C. L., Guo, S. X., Yang, J. S., & Xiao, P. G. (2008). Two new compounds from *Dendrobium candidum*. Chemical and Pharmaceutical Bulletin, 56(10), 1477-1479. doi:10.1248/cpb.56.1477
- Li, Y. P., Qing, C., Fang, T. T., Liu, Y., & Chen, Y. G. (2009). Chemical constituents of *Dendrobium chrysotoxum*. Chemistry of Natural Compounds, 45(3), 414-416. doi:10.1007/s10600-009-9329-7
- Li, J. T., Yin, B. L., Liu, Y., Wang, L. Q. and Chen, Y. G. (2009a). Mono-aromatic constituents of *Dendrobium longicornu*. Chemistry of Natural Compounds

45(2), 234-236.

- Li, Y., Wang, C. L., Wang, Y. J., Guo, S. X., Yang, J. S., Chen, X. M. and Xiao, P. G. (2009b). Three new bibenzyl derivatives from *Dendrobium candidum*. Chemical and Pharmaceutical Bulletin (Tokyo) 57(2), 218-219.
- Limpanit, R., Chuanasa, T., Likhitwitayawuid, K., Jongbunprasert, V., & Sritularak, B. (2016).  $\alpha$ -Glucosidase inhibitors from *Dendrobium tortile*. Records of Natural Products, 10, 609-616.
- Lin, L. J., Huang, X. B., & Lv, Z. C. (2016). Isolation and identification of flavonoids components from *Pteris vittata* L. SpringerPlus, 5(1), 1-3.
- Lin, T. H., Chang, S. J., Chen, C. C., Wang, J. P., & Tsao, L. T. (2001). Two phenanthraquinones from *Dendrobium moniliforme*. Journal of Natural Products, 64(8), 1084-1086. doi:10.1021/np010016i
- Liu, Y., Jiang, J. H., Yin, B. L. and Chen, Y. G. (2009a). Chemical constituents of *Dendrobium cariniferum*. Chemistry of Natural Compounds 45(2), 237-238.
- Liu, Y., Jiang, J. H., Zhang, Y., & Chen, Y. G. (2009b). Chemical constituents of *Dendrobium aurantiacum* var. *denneanum*. Chemistry of Natural Compounds, 45(4), 525-527. doi:10.1007/s10600-009-9393-z
- Lu, Y., Kuang, M., Hu, G. P., Wu, R. B., Wang, J., Liu, L., & Lin, Y. C. (2014). Loddigesiinols G-J:  $\alpha$ -glucosidase inhibitors from *Dendrobium loddigesii*. Molecules, 19(6), 8544-8555. doi:10.3390/molecules19068544
- Ma, G. X., Wang, T. S., Yin, L., Pan, Y., Xu, G. J., & Xu, L. S. (1998). Studies on chemical constituents of *Dendrobium chryseum*. Journal of Chinese Pharmaceutical Sciences, 7(1), 52-54.
- Ma, C., Meng, C. W., Zhou, Q. M., Peng, C., Liu, F., Zhang, J. W., Xiong, L. (2019a). New sesquiterpenoids from the stems of *Dendrobium nobile* and their neuroprotective activities. Fitoterapia, 138, 104351. doi:10.1016/j.fitote.2019.104351
- Ma, R. J., Yang, L., Bai, X., Li, J. Y., Yuan, M. Y., Wang, Y. Q., Zhou, J. (2019b). Phenolic Constituents with Antioxidative, Tyrosinase Inhibitory and Anti-aging Activities

- from *Dendrobium loddigesii* Rolfe. Natural Products and Bioprospecting, 9(5), 329-336. doi:10.1007/s13659-019-00219-y
- Majumder, P. L., & Chatterjee, S. (1989). Crepidatin, a bibenzyl derivative from the orchid *Dendrobium crepidatum*. Phytochemistry, 28(7), 1986-1989. doi:10.1016/S0031-9422(00)97904-4
- Majumder, P. L., Guha, S., & Sen, S. (1999). Bibenzyl derivatives from the orchid *Dendrobium amoenum*. Phytochemistry, 52(7), 1365-1369. doi:10.1016/S0031-9422(99)00370-2
- Majumder, P. L., & Pal, S. (1992). Rotundatin, a new 9,10-dihydrophenanthrene derivative from *Dendrobium rotundatum*. Phytochemistry, 31(9), 3225-3228. doi:10.1016/0031-9422(92)83480-M
- Majumder, P. L., & Pal, S. (1993). Cumulatin and tristin, two bibenzyl derivatives from the orchids *Dendrobium cumulatum* and *Bulbophyllum triste*. Phytochemistry, 32(6), 1561-1565. doi:10.1016/0031-9422(93)85180-Y
- Majumder, P. L., & Sen, R. C. (1987). Moscatilin, a bibenzyl derivative from the orchid *Dendrobium moscatum*. Phytochemistry, 26(7), 2121-2124. doi:10.1016/S0031-9422(00)81777-X
- Mittraphab, A., Muangnoi, C., Likhitwitayawuid, K., Rojsitthisak, P., & Sritularak, B. (2016). A new bibenzyl-phenanthrene derivative from *Dendrobium signatum* and its cytotoxic activity. Natural Product Communications, 11(5), 657-659. doi:10.1177/1934578X1601100526
- Miyazawa, M., Shimamura, H., Nakamura, S., Sugiura, W., Kosaka, H., & Kameoka, H. (1999). Moscatilin from *Dendrobium nobile*, a naturally occurring bibenzyl compound with potential antimutagenic activity. Journal of Agricultural and Food Chemistry, 47(5), 2163-2167. doi:10.1021/jf970930a
- Mukherjee, S., & Jagtap, S. (2020). Use of Orchids in treating Diabetes and related diseases. The Journal of Phytopharmacology, 9(2) : 130-138.
- Na Ranong, S., Likhitwitayawuid, K., Mekboonsonglarp, W., & Sritularak, B. (2019). New dihydrophenanthrenes from *Dendrobium infundibulum*. Natural product research, 33(3), 420-426.

- Nkobole, N., Houghton, P. J., Hussein, A., & Lall, N. (2011). Antidiabetic activity of *Terminalia sericea* constituents. Natural product communications, 6(11), 1934578X1100601106.
- Ono M, Ito Y, Masuoka C, Koga H, Nohara T. (1995) Antioxidative constituents from *Dendrobii* Herba (stems of *Dendrobium* spp.). Food Science and Technology International, 2, 115-120.
- Pan, H. M., Chen, B., Li, F., & Wang, M. K. (2012). Chemical Constituents of *Dendrobium denneanum*. Chinese Journal of Applied Environmental Biology, 18(3), 378-380. doi:10.3724/sp.j.1145.2012.00378
- Patel, D., Kumar, R., Laloo, D., & Hemalatha, S. (2012). Diabetes mellitus: an overview on its pharmacological aspects and reported medicinal plants having antidiabetic activity. Asian Pacific Journal of Tropical Biomedicine, 2(5), 411-420.
- Phechrmeekha, T., Sritularak, B., & Likhitwitayawuid, K. (2012). New phenolic compounds from *Dendrobium capillipes* and *Dendrobium secundum*. Journal of Asian Natural Products Research, 14(8), 748-754. doi:10.1080/10286020.2012.689979
- Plants of the World Online. (2017). *Dendrobium braianense* Gagnep. Retrieved from <http://plantsoftheworldonline.org/taxon/urn:lsid:ipni.org:names:626961-1>
- Qi, L.-W., Liu, E.-H., Chu, C., Peng, Y.-B., Cai, H.-X., & Li, P. (2010). Anti-diabetic agents from natural products—an update from 2004 to 2009. Current Topics in Medicinal Chemistry, 10(4), 434-457.
- Qin, X. D., Qu, Y., Ning, L., Liu, J. K., & Fan, S. K. (2011). A new picrotoxane-type sesquiterpene from *Dendrobium findlayanum*. Journal of Asian Natural Products Research, 13(11), 1047-1050. doi:10.1080/10286020.2011.620952
- Rungwichaniwat, P., Sritularak, B., & Likhitwitayawuid, K. (2014). Chemical constituents of *Dendrobium williamsonii*. Pharmacognosy Journal, 6(3), 36-41. doi:10.5530/pj.2014.3.6
- San, H. T., Boonsongcheep, P., Putalun, W., Mekboonsonglarp, W., Sritularak, B., & Likhitwitayawuid, K. (2020).  $\alpha$ -Glucosidase inhibitory and glucose uptake

- stimulatory effects of phenolic compounds from *Dendrobium christyanum*. Natural Product Communications, 15(3), 1934578X20913453.
- Sarakulwattana, C., Mekboonsonglarp, W., Likhitwitayawuid, K., Rojsitthisak, P., & Sritularak, B. (2018). New bisbibenzyl and phenanthrene derivatives from *Dendrobium scabrilingue* and their alpha-glucosidase inhibitory activity. Natural Product Research, 1-8. doi:10.1080/14786419.2018.1527839
- Sears, B., & Perry, M. (2015). The role of fatty acids in insulin resistance. Lipids in health and disease, 14(1), 1-9.
- Sintayehu B, Asres K, Raghavendra Y. Radical scavenging activities of the leaf extracts and a flavonoid glycoside isolated from *Cineraria abyssinica* Sch. Bip. Exa. Rich. Journal of Applied Pharmaceutical Science. 2012; 2: 44-49.
- Srianta, I., Kusumawati, N., Nugerahani, I., Artanti, N., & Xu, G. R. (2013). In vitro [alpha]-glucosidase inhibitory activity of *Monascus*-fermented durian seed extracts. International Food Research Journal, 20(2), 533.
- Sritularak, B., Anuwat, M., & Likhitwitayawuid, K. (2011). A new phenanthrenequinone from *Dendrobium draconis*. Journal of Asian Natural Products Research, 13(3), 251-255. doi:10.1080/10286020.2010.546354
- Sritularak, B., Duangrak, N., & Likhitwitayawuid, K. (2011). A new bibenzyl from *Dendrobium secundum*. Zeitschrift fur Naturforschung. C, Journal of Biosciences, 66(5-6), 205-208. doi:10.1515/znc-2011-5-602
- Sritularak, B., & Likhitwitayawuid, K. (2009). New bisbibenzyls from *Dendrobium falconeri*. Helvetica Chimica Acta, 92(4), 740-744. doi:10.1002/hlca.200800391
- Stumvoll, M., Goldstein, B. J., & Van Haeften, T. W. (2005). Type 2 diabetes: principles of pathogenesis and therapy. The Lancet, 365(9467), 1333-1346.
- Sukphan, P., Sritularak, B., Mekboonsonglarp, W., Lipipun, V., & Likhitwitayawuid, K. (2014). Chemical constituents of *Dendrobium venustum* and their antimalarial and anti-herpetic properties. Natural Product Communications, 9(6), 825-827.
- Sun, J., Zhang, F., Yang, M., Zhang, J., Chen, L., Zhan, R., Chen, Y. (2014). Isolation of alpha-glucosidase inhibitors including a new flavonol glycoside from

- Dendrobium devonianum*. Natural Product Research, 28(21), 1900-1905. doi:10.1080/14786419.2014.955495
- Talapatra, B., Das, A. K., & Talapatra, S. K. (1989). Defuscin, a new phenolic ester from *Dendrobium fuscescens*: Conformation of shikimic acid. Phytochemistry, 28(1), 290-292.
- Tanagornmeatar, K., Chaotham, C., Sritularak, B., Likhitwitayawuid, K., & Chanvorachote, P. (2014). Cytotoxic and anti-metastatic activities of phenolic compounds from *Dendrobium ellipsophyllum*. Anticancer Research, 34(11), 6573-6580. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/25368260>
- The Plant List. (2018). IOSPE Photos. *Dendrobium braianense* Gagnep. Retrieved from <http://www.orchidspecies.com/denbraianense.htm>
- Tsopmo, A., Awah, F. M., & Kuete, V. (2013). Lignans and stilbenes from African medicinal plants. In Medicinal plant research in Africa. Elsevier (pp. 435-478).
- Van de Laar, F. A. (2008). Alpha-glucosidase inhibitors in the early treatment of type 2 diabetes. Vascular health and risk management, 4(6), 1189.
- Veerraju, P., Rao, N. S. P., Rao, L. J., Rao, K. V. J., & Rao, P. R. M. (1989). Amoenumin, A 9, 10-dihydro-5H-phenanthro-(4, 5-b, c, d)-pyran from *Dendrobium amoenum*. Phytochemistry, 28(3), 950-951. doi:10.1016/0031-9422(89)80154-2
- Wang, H., Zhao, T., & Che, C.-T. (1985). Dendrobine and 3-hydroxy-2-oxodendrobine from *Dendrobium nobile*. Journal of Natural Products, 48(5), 796-801.
- Wang, L., Zhang, C. F., Wang, Z. T., Zhang, M., & Xu, L. S. (2009). Five new compounds from *Dendrobium crystallinum*. Journal of Asian Natural Products Research, 11(11), 903-911. doi:10.1080/10286020903128942
- Wang, P., Chen, X., Wang, H., Huang, S., Cai, C., Yuan, J., Dai, H. (2019). Four new picrotoxane-type sesquiterpenes from *Dendrobium nobile* Lindl. Frontiers in Chemistry, 7, 812. doi:10.3389/fchem.2019.00812
- Wu, L., Lu, Y., Ding, Y., Zhao, J., Xu, H., & Chou, G. (2019). Four new compounds from *Dendrobium devonianum*. Natural Product Research, 33(15), 2160-2168. doi:10.1080/14786419.2018.1490900



- Xiong, L., Cao, Z. X., Peng, C., Li, X. H., Xie, X. F., Zhang, T. M., Guo, L. (2013). Phenolic glucosides from *Dendrobium aurantiacum* var. *denneanum* and their bioactivities. Molecules, 18(6), 6153-6160. doi:10.3390/molecules18066153
- Xu, Fan, W.-W., Zi, C.-T., Dong, F.-W., Yang, D., Zhou, J., & Hu, J.-M. (2017). Four new glycosides from the stems of *Dendrobium fimbriatum* Hook. Natural product research, 31(7), 797-801.
- Xu, F. Q., Xu, F. C., Hou, B., Fan, W. W., Zi, C. T., Li, Y., Hu, J. M. (2014). Cytotoxic bibenzyl dimers from the stems of *Dendrobium fimbriatum* Hook. Bioorganic and Medicinal Chemistry Letters, 24(22), 5268-5273. doi:10.1016/j.bmcl.2014.09.052
- Xu, X., Li, Z., Yang, R., Zhou, H., Bai, Y., Yu, M., Li, B. (2019). Crepidatumines C and D, two new indolizidine alkaloids from *Dendrobium crepidatum* Lindl. ex Paxt. Molecules, 24(17). doi:10.3390/molecules24173071
- Yamaki, M., & Honda, C. (1996). The stilbenoids from *Dendrobium plicate*. Phytochemistry, 43(1), 207-208.
- Yang, D., Cheng, Z.-Q., Zi, C.-T., Yang, L., Dong, F.-W., Hu, J.-M., & Zhou, J. (2019). Four new sesquiterpene derivatives from *Dendrobium findlayanum*. Chinese Journal of Natural Medicines, 17(12), 900-905. doi:10.1016/s1875-5364(19)30110-4
- Yang, D., Cheng, Z. Q., Yang, L., Hou, B., Yang, J., Li, X. N., Hu, J. M. (2018). Seco-Dendrobine-Type Alkaloids and Bioactive Phenolics from *Dendrobium findlayanum*. The Natural Products Journal, 81(2), 227-235. doi:10.1021/acs.jnatprod.7b00150
- Yang, D., Liu, L. Y., Cheng, Z. Q., Xu, F. Q., Fan, W. W., Zi, C. T., Hu, J. M. (2015). Five new phenolic compounds from *Dendrobium aphyllum*. Fitoterapia, 100, 11-18. doi:10.1016/j.fitote.2014.11.004
- Yang, H., Chou, G. X., Wang, Z. T., Guo, Y. W., Hu, Z. B., & Xu, L. S. (2004). Two new compounds from *Dendrobium chrysotoxum*. Helvetica Chimica Acta, 87(2), 394-399. doi:10.1002/hlca.200490037

- Yang, H. K., Sung, S. H., & Kim, Y. C. (2007). Antifibrotic phenanthrenes of *Dendrobium nobile* stems. Journal of Natural Products, 70(12), 1925–1929. doi:10.1021/np070423f
- Yang, L., Qin, L. H., Bligh, S. W., Bashall, A., Zhang, C. F., Zhang, M., Xu, L. S. (2006). A new phenanthrene with a spirolactone from *Dendrobium chrysanthum* and its anti-inflammatory activities. Bioorganic and Medicinal Chemistry, 14(10), 3496–3501. doi:10.1016/j.bmc.2006.01.004
- Yang, L., Wang, Z. T., & Xu, L. S. (2006). Phenols and a triterpene from *Dendrobium aurantiacum* var. *denneanum* (Orchidaceae). Biochemical Systematics and Ecology, 34(8), 658–660. doi:10.1016/j.bse.2006.03.003
- Yang, M., Chen, L. J., Zhang, Y., & Chen, Y. G. (2019). Two new picrotoxane-type sesquiterpenoid lactones from *Dendrobium williamsonii*. Journal of Asian Natural Products Research, 21(2), 129–133. doi:10.1080/10286020.2017.1394294
- Yang, M., Zhang, Y., Chen, L., & Chen, Y. (2018). A new (propylphenyl) bibenzyl derivative from *Dendrobium williamsonii*. Natural Product Research, 32(14), 1–7. doi:10.1080/14786419.2017.1396599
- Ye, Q. H., Mei, Y., Yang, P. M., Cheng, L., & Kong, D. Y. (2016). A New 9,10-dihydrophenanthrene glycoside from *Dendrobium primulinum*. Chemistry of Natural Compounds, 52(3), 381–383. doi:10.1007/s10600-016-1653-0
- Ye, Q. H., Qin, G. W., & Zhao, W. M. (2002). Immunomodulatory sesquiterpene glycosides from *Dendrobium nobile*. Phytochemistry, 61(8), 885–890. doi:10.1016/s0031-9422(02)00484-3
- Ye, Q. H., & Zhao, W. M. (2002). New alloaromadendrane, cadinene and cyclocopacamphane type sesquiterpene derivatives and bibenzyls from *Dendrobium nobile*. Planta Medica, 68(8), 723–729. doi:10.1055/s-2002-33786
- Ye, Q. H., Zhao, W. M., & Qin, G. W. (2004). Lignans from *Dendrobium chrysanthum*. Journal of Asian Natural Products Research, 6(1), 39–43. doi:10.1080/1028602031000119808

- Zhang, C. F., Wang, M., Wang, L., linuma, M., Zhang, M., Xu, L. S., & Wang, Z. T. (2008). Chemical constituents of *Dendrobium grastiosissimum* and their cytotoxic activities. Indian Journal of Chemistry, 47B, 952-956.
- Zhang, G. N., Zhong, L. Y., Bligh, S. W., Guo, Y. L., Zhang, C. F., Zhang, M., Xu, L. S. (2005). Bi-bicyclic and bi-tricyclic compounds from *Dendrobium thyrsiflorum*. Phytochemistry, 66(10), 1113-1120. doi:10.1016/j.phytochem.2005.04.001
- Zhang, X., Gao, H., Han, H. Y., Liu, H., Wang, N. L., Yao, X.-S., & Wang, Z. (2007). Sesquiterpenes from *Dendrobium nobile*. Chinese Traditional and Herbal Drugs, 38, 1771-1774.
- Zhang, X., Gao, H., Wang, N. L., & Yao, X. S. (2006). Three new bibenzyl derivatives from *Dendrobium nobile*. Journal of Asian Natural Products Research, 8(1-2), 113-118. doi:10.1080/10286020500480654
- Zhang, X., Liu, H.-W., Gao, H., Han, H.-Y., Wang, N.-L., Wu, H.-M., Wang, Z. (2007). Nine New Sesquiterpenes from *Dendrobium nobile*. Helvetica Chimica Acta, 90(12), 2386-2394. doi:10.1002/hlca.200790245
- Zhang, X., Tu, F. J., Yu, H. Y., Wang, N. L., Wang, Z., & Yao, X. S. (2008). Copacamphane, picrotoxane and cyclocopacamphane sesquiterpenes from *Dendrobium nobile*. Chemical & Pharmaceutical Bulletin, 56(6), 854-857. doi:10.1248/cpb.56.854
- Zhang, X., Xu, J. K., Wang, J., Wang, N. L., Kurihara, H., Kitanaka, S., & Yao, X. S. (2007). Bioactive bibenzyl derivatives and fluorenones from *Dendrobium nobile*. The Natural Products Journal, 70(1), 24-28. doi:10.1021/np060449r
- Zhang, X., Xu, J. K., Wang, N. L., Kurihara, H., & Yao, X. S. (2008). Antioxidant phenanthrenes and lignans from *Dendrobium nobile*. Journal of Chinese Pharmaceutical Sciences, 17, 314-318.
- Zhang, Y. Y., Wang, P., Song, X. Q., Zuo, W. J., Wang, H., Chen, L. L., Dai, H. F. (2019). Chemical constituents from *Dendrobium hainanense*. Journal of Asian Natural Products Research, 21(9), 873-880. doi:10.1080/10286020.2018.1475476
- Zhao, C. S., Liu, Q. F., Halaweish, F., Shao, B. P., Ye, Y. Q., & Zhao, W. M. (2003). Copacamphane, picrotoxane, and alloaromadendrane sesquiterpene

- glycosides and phenolic glycosides from *Dendrobium moniliforme*. Journal of Natural Products, 66(8), 1140-1143. doi:10.1021/np0301801
- Zhou, C., Xie, Z., Lei, Z., Huang, Y., & Wei, G. (2018). Simultaneous identification and determination of flavonoids in *Dendrobium officinale*. Chemistry Central Journal, 12(1), 1-9.
- Zhao, N., Yang, G., Zhang, Y., Chen, L., & Chen, Y. (2016). A new 9, 10-dihydrophenanthrene from *Dendrobium moniliforme*. Natural product research, 30(2), 174-179.
- Zhao, G. Y., Deng, B. W., Zhang, C. Y., Cui, Y. D., Bi, J. Y., & Zhang, G. G. (2018). New phenanthrene and 9, 10-dihydrophenanthrene derivatives from the stems of *Dendrobium officinale* with their cytotoxic activities. Journal of Natural Medicines, 72(1), 246-251. doi:10.1007/s11418-017-1141-2
- Zhao, N., Yang, G., Zhang, Y., Chen, L., & Chen, Y. (2016). A new 9,10-dihydrophenanthrene from *Dendrobium moniliforme*. Natural Product Research, 30(2), 174-179. doi:10.1080/14786419.2015.1046379
- Zhao, W. M., Ye, Q. H., Tan, X. J., Jiang, H. L., Li, X. Y., Chen, K. X., & Kinghorn, A. D. (2001). Three new sesquiterpene glycosides from *Dendrobium nobile* with immunomodulatory activity. Journal of Natural Products, 64(9), 1196-1200. doi:10.1021/np0102612
- Zhou, X. M., Zhang, B., Chen, G. Y., Han, C. R., Jiang, K. C., Luo, M. Y., Lin, S. D. (2018). Dendrocoumarin: a new benzocoumarin derivative from the stem of *Dendrobium nobile*. Natural Product Research, 32(20), 1-4. doi:10.1080/14786419.2017.1419241
- Zhou, X. M., Zheng, C. J., Wu, J. T., Chen, G. Y., Chen, J., & Sun, C. G. (2016). Five new lactone derivatives from the stems of *Dendrobium nobile*. Fitoterapia, 115, 96-100. doi:10.1016/j.fitote.2016.10.002
- Zhou, X. M., Zheng, C. J., Wu, J. T., Chen, G. Y., Zhang, B., & Sun, C. G. (2017). A new phenolic glycoside from the stem of *Dendrobium nobile*. Natural Product Research, 31(9), 1042-1046. doi:10.1080/14786419.2016.1266352
- Ziegler, J., & Facchini, P. J. (2008). Alkaloid biosynthesis: metabolism and trafficking. Annu. Rev. Plant Biol., 59, 735-769.

## VITA

NAME	Mia Indria Permatasari
DATE OF BIRTH	15 February 1992
PLACE OF BIRTH	Ungaran, Indonesia
INSTITUTIONS ATTENDED	Faculty of Pharmaceutical Sciences, Chulalongkorn University
HOME ADDRESS	St. Kakap Raya No. 1A RT 12 RW 05, Sebantengan, Ungaran, Central Java, Indonesia
PUBLICATION	Mia Indria Permatasari, Kittisak Likhitwitayawuid, Boonchoo Sritularak. CHEMICAL CONSTITUENTS OF <i>Dendrobium braianense</i> AND THEIR ALPHA-GLUCOSIDASE INHIBITORY ACTIVITY. The 46th International Congress on Science, Technology and Technology-based Innovation. 2020; 988.
AWARD RECEIVED	ASEAN SCHOLARSHIP, Chulalongkorn University