

1-1-1981

ทดสอบภูมิคุ้มกันของท่อน ไทยเภสัชสาร

อัจฉรา อภิศวรรณกุล

Follow this and additional works at: <https://digital.car.chula.ac.th/tjps>



Part of the [Pharmacology Commons](#)

Recommended Citation

อภิศวรรณกุล, อัจฉรา (1981) "ทดสอบภูมิคุ้มกันของท่อน ไทยเภสัชสาร," *The Thai Journal of Pharmaceutical Sciences*: Vol. 6: Iss. 2, Article 10.

DOI: <https://doi.org/10.56808/3027-7922.1664>

Available at: <https://digital.car.chula.ac.th/tjps/vol6/iss2/10>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Pharmaceutical Sciences by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.



ทดสอบภูมิคุ้มกันของท่าน

PHARM QUIZ

ไทยเภสัชสาร

VOL 6 เล่ม 2

PHARM QUIZ

1. THE BROMSULPHALEIN (BSP.) EXCRETION TEST IS A MEASURE OF
- A. LIVER FUNCTION
 - B. THYROID FUNCTION
 - C. PANCREATIC FUNCTION
 - D. KIDNEY FUNCTION
 - E. CIRCULATION TIME

2. WHICH OF THE FOLLOWING TESTS IS SPECIFIC FOR THE MEASUREMENT OF GLUCOSE
- A. BENEDICT'S SOLUTION
 - B. ACETEST[®] TABLETS
 - C. CLINITEST[®] TABLETS
 - D. TES-TAPE[®]
 - E. KETOSTIX[®]

3. WHICH OF THE FOLLOWING AGENTS IS EFFECTIVE AGAINST

PENICILLINASE-PRODUCING STAPHYLOCOCCI

- A. SODIUM METHICILLIN
- B. PENICILLIN .
- C. PHENETHICILLIN
- D. POTASSIUM HETACILLIN
- E. AMOXICILLIN

4. A PHARMACIST RECEIVES A PRESCRIPTION ORDER FOR PHENYBUTAZONE (BUTAZOLIDIN[®]). HE SHOULD CONSULT WITH THE PHYSICIAN IF THE MEDICATION RECORD INDICATES THAT THE PATIENT.

- A. HAS GOUT
- B. HAS ARTHRITIS
- C. IS HYPERTENSIVE
- D. HAS INSOMNIA
- E. HAS A PEPTIC ULCER

5. IBUPROFEN HAS A P^{K_a} OF 5.5 IF THE pH OF THE PATIENT'S URINE IS 7.5, WHAT WOULD THE RATIO OF DISSOCIATED TO UNDISSOCIATED DRUG BE

- A. 2:1
- B. 100:1
- C. 20:1
- D. 1:2
- E. 1:100

6. THE SPLITTING OFF OF THE SURFACE OF A TABLET IS CALLED.

- A. PICKING
- B. CAPPING
- C. EROSION
- D. CRACKING
- E. A COMPRESSION FRACTURE

7. AN EXAMPLE OF A COUGH MIXTURE THAT CONTAINS ONLY THE EXPECTORANT GLYCERYL GUAIACOLATE AS THE ACTIVE INGREDIENT IS.

- A. CORICIDIN COUGH FORMULA[®]
- B. IPSATOL[®] SYRUP
- C. ROBITUSSIN[®]
- D. SILENCE IS GOLDEN[®] SYRUP
- E. TRIND[®]

8. PARENTERALLY ADMINISTERED ELECTROLYTES ARE USUALLY ORDERED IN.

- A. EQUIVALENTS
- B. MILLIEQUIVALENTS (mEq)
- C. MILLIGRAMS PERCENT (mg%)
- D. MILLIMOLES
- E. MICROGRAMS (mcg)

9. MYXEDEMA IS A _____ STATE.

- A. HYPOTHYROID
- B. HYPOPARATHYROID
- C. HYPERTHYROID
- D. HYPERPARATHYROID
- E. NORMAL PARATHYROID

10. THE BIOLOGICAL HALF-LIFE ($t_{1/2}$) OF A DRUG IS THE TIME.

A. IN WHICH THE POTENCY OF A DRUG IN SOLUTION FALLS ONE-HALF OF ITS INITIAL VALUE.

B. REQUIRED FOR THE BLOOD CONCENTRATION TO BE REDUCED BY 50%

C. REQUIRED FOR 50% OF THE DRUG TO BE ABSORBED FROM THE GI-TRACT

D. REQUIRED FOR THE PHARMACOLOGIC EFFECT TO DECREASE

TO ONE-HALF OF ITS PEAK.

E. REQUIRED FOR ONE-HALF
OF THE TOTAL DOSE TO BE
EXCRETED IN THE URINE

B	10	B	5
A	9	E	4
B	8	A	3
C	7	D	2
B	6	A	1

๑๒๒

อัจฉรา อู่สุวรรณกุล