

3-1-2004

EVALUATION OF THE AFLATOXIN B1 ADSORPTION CAPACITY OF BENTONITE USING AN IN VITRO METHOD MIMICKING MONOGASTRIC GASTRO-INTESTINAL TRACT CONDITIONS

Piyarat Chansiripornchai

Johanna Fink-Gremmels

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Chansiripornchai, Piyarat and Fink-Gremmels, Johanna (2004) "EVALUATION OF THE AFLATOXIN B1 ADSORPTION CAPACITY OF BENTONITE USING AN IN VITRO METHOD MIMICKING MONOGASTRIC GASTRO-INTESTINAL TRACT CONDITIONS," *The Thai Journal of Veterinary Medicine*: Vol. 34: Iss. 1, Article 5.

Available at: <https://digital.car.chula.ac.th/tjvm/vol34/iss1/5>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

EVALUATION OF THE AFLATOXIN B1 ADSORPTION CAPACITY OF BENTONITE USING AN *IN VITRO* METHOD MIMICKING MONOGASTRIC GASTRO-INTESTINAL TRACT CONDITIONS

Piyarat Chansiripornchai^{1*} Johanna Fink-Gremmels²

Abstract

Piyarat Chansiripornchai^{1*} Johanna Fink-Gremmels²

EVALUATION OF THE AFLATOXIN B1 ADSORPTION CAPACITY OF BENTONITE USING AN *IN VITRO* METHOD MIMICKING MONOGASTRIC GASTRO-INTESTINAL TRACT CONDITIONS

Aflatoxins are considered as unavoidable contaminants of food and animal feeds, since their production cannot be prevented successfully using current agricultural practices. The use of aflatoxin adsorbents as feed supplements is one of the most promising approaches to reduce aflatoxicoses in animals. Since there are numerous adsorbents marketed, an effective and practical method for the rapid evaluation of adsorption capacity of individual products is needed. In this study, we used an *in vitro* model designed to mimic the temperature, pH and the time of passage through the stomach and the intestinal tract of a monogastric animal, for the evaluation of aflatoxin B1 adsorption capacity of different mineral clays. The results indicate that 6 out of the 7 tested products had a high binding capacity (ranging from 98.97-100%), under the experimental conditions used. The data also demonstrate that the method allows us to differentiate those compounds with high binding capacity from those with a lesser binding capacity. Such an *in vitro* method seems useful for the rapid screening of different substances and for the pre-selection of adsorbent products that may be used as feed supplements.

Keywords : adsorbent, aflatoxin B₁, bentonite, detoxification, *in vitro*

¹Department of Veterinary Pharmacology, Faculty of Veterinary Science, Chulalongkorn University, Henri-Dunant Rd., Bangkok, 10330, Thailand

²Department of Veterinary Pharmacology, Pharmacy and Toxicology, Faculty of Veterinary Science, Utrecht University, P.O. Box 80.176.3508 TD Utrecht, The Netherlands

*Corresponding author

¹ภาควิชาเภสัชวิทยา คณะสัตวแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ถนนอังรีดูนังต์ กรุงเทพฯ 10330

²ภาควิชาเภสัชวิทยา เภสัชศาสตร์ และพิษวิทยา คณะสัตวแพทยศาสตร์ มหาวิทยาลัย อูเทรخت ตู้ไปรษณีย์ 80.176.3508 อูเทรخت ราชอาณาจักรเนเธอร์แลนด์

*ผู้รับผิดชอบบทความ

บทคัดย่อ

ปิยะรัตน์ จันทร์ศิริพรชัย^{1*} โยธันนา ฟิงค์-เครมเมิลส์²

การประเมินประสิทธิภาพของเบนโทไนด์ในการดูดซับอะฟลาท็อกซินบี 1 โดยวิธีจำลองสถานะของทางเดินอาหารแบบกระเพาะเดี่ยวภายนอกร่างกาย

อะฟลาท็อกซินเป็นสารพิษจากเชื้อราที่มักพบปนเปื้อนในอาหารของมนุษย์และสัตว์เนื่องจากกระบวนการทางเกษตรกรรม ในปัจจุบันยังไม่สามารถป้องกันการสร้างสารพิษชนิดนี้จากเชื้อราเมื่อมีการปนเปื้อนในอาหารได้ การใช้สารดูดซับอะฟลาท็อกซินผสมลงในอาหารสัตว์เป็นวิธีการหนึ่งที่ยอมรับใช้ในการลดความเป็นพิษจากอะฟลาท็อกซินในสัตว์ เนื่องจากมีการผลิตสารดูดซับออกจำหน่ายในท้องตลาดอย่างมากมาย ดังนั้นจึงมีความจำเป็นที่จะต้องมียุทธวิธีที่มีประสิทธิภาพและสะดวกในทางปฏิบัติเพื่อใช้ประเมินความสามารถในการดูดซับสารพิษของสารดูดซับแต่ละชนิด การศึกษาครั้งนี้ได้จำลองสถานะทางเดินอาหารของสัตว์กระเพาะเดี่ยวทั้งอุณหภูมิต่ำ ความเป็นกรดต่ำ และเวลาในการผ่านของอาหารจากกระเพาะอาหารเข้าสู่ลำไส้ เพื่อประเมินประสิทธิภาพของเบนโทไนด์ในการดูดซับอะฟลาท็อกซินบี 1 จากผลการทดลองพบว่าสารดูดซับ 6 ใน 7 ชนิดมีประสิทธิภาพในการดูดซับอะฟลาท็อกซินบี 1 ได้ดีมาก (98.97-100%) ผลการศึกษายังแสดงให้เห็นว่าสามารถจำแนกสารดูดซับอะฟลาท็อกซินที่มีประสิทธิภาพสูงออกจากสารดูดซับที่มีประสิทธิภาพต่ำได้โดยใช้วิธีการที่ประยุกต์ขึ้นนี้ ดังนั้นการใช้วิธีจำลองสถานะของทางเดินอาหารแบบกระเพาะเดี่ยวภายนอกร่างกายเพื่อประเมินประสิทธิภาพของสารดูดซับอะฟลาท็อกซินจึงเป็นวิธีการที่มีประโยชน์ในการทดสอบประสิทธิภาพของสารดูดซับอะฟลาท็อกซินเบื้องต้นได้อย่างรวดเร็ว เพื่อนำไปประกอบการพิจารณานำสารดูดซับชนิดนั้นๆ มาใช้เป็นสารเติมเพื่อลดพิษของอะฟลาท็อกซินในอาหารสัตว์

คำสำคัญ : สารดูดซับ อะฟลาท็อกซินบี 1 เบนโทไนด์ การลดความเป็นพิษ ภายนอกในร่างกาย

Introduction

Some field and storage fungi produce a variety of secondary metabolites, mycotoxins, that exert adverse effects on human and animal health. These mycotoxins can induce organ lesions and may be mutagenic, carcinogenic, immunosuppressive as well as interfering with hormonal function (Karlovsky, 1999). Among the group of mycotoxins, which represents more than 300 structurally diverse compounds, the aflatoxins have been the most intensely studied, since their discovery in the 1960s (Asao et al., 1965; Brechbuhler et al., 1967). The occurrence of these toxins in food and feed materials has caused not only adverse health effects in animals and humans, but has also resulted in economic loss, for exporting countries, due to non-marketable products (Basappa and Shantha, 1996).

Aflatoxin B₁ (AFB₁) and various aflatoxin analogs are commonly produced by *Aspergillus flavus* and *A. parasiticus* molds that occur in the field and during the storage of grains, cereals and nuts. Natural aflatoxins are AFB₁, AFB₂, AFG₁ and AFG₂. However, following hepatic biotransformation the metabolites M₁, M₂, M_{2a}, MP₁, P₁, H₁ and Q₁ have been identified, whereas aflatoxicol I and II originate from bacterial metabolism (Phillips, 1999). AFB₁ is the most potent mycotoxins and can be both acutely toxic and carcinogenic for many animal species (Wogan and Newberne, 1967; Hendricks, 1994).

At present, aflatoxins are considered unavoidable contaminants of food, since they cannot be prevented or eliminated by current agricultural practices (CAST, 1989). Because aflatoxin is heat stable, food and feed processing

procedures do not reduce the toxin concentration. Thus, aflatoxins are found as a contaminant of corn, peanuts, peanut butter, breakfast cereals, cornmeal, cottonseed, tortillas, animal feeds and the dairy products (including ice cream and cheese) derived from milk of toxin-exposed dairy cattle. In developed countries, grains and oil seeds that are contaminated with aflatoxins at concentrations higher than 20 ppb are condemned (Phillips, 1999). However, in lesser-developed countries with limited supplies of food, it is neither feasible, nor practical, to condemn and destroy all contaminated grains. Therefore, safe and practical chemical intervention methods for the detoxification of food and animal feeds in terms of chemoprevention against aflatoxicosis are critical needs.

In animal production, the use of mycotoxin adsorbents as feed supplements is one of the most promising approaches towards the reduction of mycotoxicoses in farm animals and towards a minimalization of carry-over of these mycotoxins from contaminated feeds into animal products (Ramos et al., 1996; Huwig et al., 2001). Different substances, including phyllosilicate minerals, zeolites, activated charcoal, synthetic resins and yeast cell-wall-derived products, have been used for the adsorption of aflatoxins *in vitro* and *in vivo* as described in numerous reports (Devegowda et al., 1996; Galvano et al., 1996; Kececi et al., 1998; Miazzo et al., 2000; Raju et al., 2000; Rosa et al., 2001). Since there are many adsorbents on the market, it is necessary to establish an effective and practical method to evaluate the aflatoxin and other mycotoxins adsorption capacity of individual adsorbents.

Our objective in this study was to determine the adsorption capacity of bentonite for AFB₁ using an *in vitro* method designed to mimic the conditions of a monogastric animal, which can also be used as a predictive value for humans.

Materials and methods

Materials. Seven mineral clays were obtained from commercial companies in the Netherlands. The composition of the mineral clays were labeled as

bentonites originating from different natural resources. AFB₁ standard was purchased from Sigma (St. Louis, CA, USA). All the solvents used in chromatography were of HPLC grade and all other reagents were of analytical grade.

Assessment of pH-dependent adsorption. A pH-dependent adsorption method was used as previously described by Sabater (Sabater, 2003). Seven bentonites named: bent.1, bent.2, bent.3, bent.4, bent.5, bent.6 and bent.7 were suspended in a phosphate buffer solution (CaCl₂ x H₂O, 1.2 mM; KCl, 2.7 mM; KH₂PO₄, 1.5 mM; MgCl₂ x 6H₂O, 1.1 mM; NaCl, 138 mM; Na₂HPO₄ x 2H₂O, 8.1 mM; pH = 7.4) to produce a final concentration of 2.5 mg of product / ml. AFB₁ standard was added to this suspension at a final concentration of 0.17 mg/l (170 ppb). Negative controls containing the AFB₁ solution without adsorbent were also included. The mixtures were adjusted to a pH of 2.5 and incubated at 37°C for 1 hr, with constant agitation, to simulate the gastric conditions of a monogastric animal. After taking an aliquot for analysis, the pH of the suspensions was further adjusted to 8 and the suspension was incubated at 37°C for 3 hr, using constant agitation to simulate the conditions of intestinal passage in a monogastric animal. A second aliquot was then taken and both aliquots were immediately filtered (Mini-GF; Sartorius, Gottingen, Germany) to separate the binder. 1.5 ml of each filtrated aliquot was extracted with 8.5 ml of chloroform. The water phase was discarded and the chloroform was evaporated. The final residue was re-dissolved in 250 µl of mobile phase from which 50 µl were injected into the HPLC system.

AFB₁ detection by HPLC. A Spherisorb ODS-2, 5 µm column (4.6x 250 mm, Chrompack, The Netherlands) was connected to two high precision pumps (Gynkotekmodel 300) set at a flow rate of 0.8 ml/min and controlled by Chromeleon-GynkotekHPLC software (Softron). Fluorescence detection was performed using a FP 920 fluorescence detector (Jasco, Japan) set at 365 nm

excitation wavelength and 420 nm emission wavelength. Mobile phase consisted of acetonitrile/methanol/water (20:20:60 v:v:v) with 58.44 mg of NaCl /l.

Statistical analysis. The significance of the differences between the control and the treatment groups was determined by one-way ANOVA. Statistical significance was accepted at $p < 0.05$.

Results

An *in vitro* system designed to mimic the temperature, pH and time of passage through the stomach and the intestinal tract of a monogastric animal was applied to evaluate the AFB₁ binding capacity of 7 bentonites. AFB₁ adsorption, after the gastric (pH 2.5) and the intestinal (pH 8) phases, was assessed by HPLC determination of the unbound AFB₁ in the incubation fluid. Table 1 shows near to 100% adsorption of AFB₁ for the bent.1 to bent.6 products tested at a concentration of 2.5 mg/ml. The bent.5 showed 100% adsorption of AFB₁ in both the pH 2.5 and the pH 8 buffer aliquots (Fig.1). In contrast, bent.7 could adsorb only 10% of AFB₁. Comparable results were found in both the pH 2.5 and the pH 8 phases (Table 2). According to the results, the best mineral clays for the pH 2.5 were bent.5, 6, 2, 4, 3 and 1, and for the pH 8 were bent.5, 2, 6, 1, 3 and 4. The product bent.7 was not a good adsorbent for AFB₁ in this study.

Discussion

Aflatoxins are considered as unavoidable contaminants of various feed and food commodities. The best procedure to prevent the negative effects of mycotoxins is to minimize the mycotoxin production itself, e.g. by harvesting the grain at maturity and by maintaining low moisture and low temperature during storage. This is difficult to carry out in countries with a warm and humid climate (Huwig et al., 2001). In order to avoid mycotoxicosis, several strategies have been investigated including biological, chemical and physical methods (Basappa and Shantha, 1996). Physical methods focus on the removal of mycotoxins by different adsorbents, added

to mycotoxin-contaminated diets (Ramos et al., 1996). Reduction of the bioavailability of aflatoxins in animals by the addition of dietary clay minerals has received considerable attention since the first reports using a processed montmorillonite clay (Phillips et al., 1987, 1988). The montmorillonites and the bentonites belong to a group of natural aluminosilicates (Ramos and Hernandez, 1997). The mechanism of action of these mineral clays is to bind irreversibly with the toxin in aqueous media, thereby, preventing absorption of the toxin across the intestinal wall (Schell et al., 1993). Now, new adsorbents for aflatoxins and other mycotoxins are released onto the market rapidly and thus it is necessary to determine the quality of individual adsorbent products. We therefore developed a practical and simple method to evaluate, *in vitro*, the aflatoxin binding capacity of mineral clays. Mycotoxin adsorbents are usually added to the feed in a concentration of 1 to 5 kg/t, which corresponds to a concentration of 1 to 5 mg/ml (Sabater, 2003). The concentration of AFB₁ used in this study (170 ppb) was much higher than the acceptable concentration of AFB₁ in animal feeds established by the US Food and Drug Administration (Basappa and Shantha, 1996). For performing AFB₁ binding assay under realistic conditions, we applied these concentrations of adsorbent and AFB₁ in our *in vitro* system.

According to our results, 6 out of 7 tested products appear to bind AFB₁ efficiently in both the pH 2.5 (98.97-100%) and the pH 8 (96.58-98.75%) phases. However, there was a different binding capacity among the tested adsorbents when tested at different pH. From the results, the binding capacity of the bent.3, 4 and 6 decreased in the pH 8 phase ($p < 0.05$). These data confirm the suitability of the applied method for the rapid screening of new products. Our results are consistent with those of Sabater (2003), who was the first to report the usefulness of such *in vitro* models under conditions of pH, temperature and time, comparable to those of a monogastric gastrointestinal tract, for evaluation of deoxynivalenol (DON) and zearalenone (ZEN) binding capacity of the adsorbents. The obvious

Table 1. AFB₁ *in vitro* adsorption at pH 2.5 assessed by HPLC for different mineral clays, added at 2.5 mg/ml¹

Product	AFB ₁ concentration (ng)	Adsorption (%)
Control	50	-
bent.1	0.52±0.01*	98.97±0.02
bent.2	0.06±0.09*	99.89±0.19
bent.3	0.29±0.17*	99.43±0.33
bent.4	0.16±0.14*	99.69±0.27
bent.5	0*	100
bent.6	0*	100
bent.7	44.31±2.81	10.77±5.68

¹Values are the mean±SD of three independent experiments.

*Significantly different from the control ($p<0.05$, one way ANOVA)

Table 2. AFB₁ *in vitro* adsorption at pH 8 assessed by HPLC for different mineral clays, added at 2.5 mg/ml¹

Product	AFB ₁ concentration (ng)	Adsorption (%)
Control	50	-
bent.1	0.83±0.83*	98.34±1.67
bent.2	0.63±0.56*	98.75±1.11
bent.3	1.06±0.10*	97.89±0.21
bent.4	1.71±0.21*	96.58±0.43
bent.5	0*	100
bent.6	0.72±0.27*	98.55±0.55
bent.7	46.43±1.39	6.48±2.81

¹ Values are the mean+SD of three independent experiments.

* Significantly different from the control ($p<0.05$, one way ANOVA)

advantage of the *in vitro* model is the ability to do rapid screening of different substances, enabling a pre-selection of reliable adsorbent products (Sabater, 2003).

Since *in vitro* studies are not always predictive of *in vivo* results, it is important to emphasize that the mycotoxins binding capacity of any new adsorbent should also be confirmed by *in vivo* experiments, before releasing the product onto the market.

Acknowledgment

The authors thank Maurice Selman, Roel Maas and Lilian de Nijs-Tjon for their kind support with the HPLC analyses.




Figure 1. HPLC-chromatograms showing the detection of AFB₁ after incubation with different adsorbents at pH 2.5 and pH 8. The arrows show the position of the AFB₁ peak. These chromatograms also demonstrate that no fluorescent AFB₁-breakdown products were formed.

References

- Asao, T., Buchi, G., Kader, M.M.A., Chang, S.B., Wick, E.L. and Wogan, G.N. 1965. The structures of aflatoxin B1 and G1. *J. Am. Chem. Soc.* 87: 882-886.
- Basappa, S.C. and Shantha, T. 1996. Methods for detoxification of aflatoxins in food and feeds-A critical appraisal. *J. Food Sci. Technol.* 33(2) : 95-107.
- Brechbuhler, S., Buchi, G. and Milne, G. 1967. The absolute configurations of the aflatoxins. *J. Org. Chem.* 32: 2641-2642.
- CAST (Council for Agricultural Science and Technology) 1989. Task Force Report No. 116. K. Niyo ed., Ames, IA, pp. 1-91.
- Devegowda, G., Aravind, B.I.R. and Morton, P. 1996. *Saccharomyces cerevisiae* mannan-oligosaccharides to counteract aflatoxins in broilers. *Proc. Aust. Poult. Sci. Symp.* 8: 103-106.
- Galvano, F., Pietri, A., Bertuzzi, T., Fusconi, G., Galvano, M., Piva, A. and Piva, G. 1996. Reduction of carry-over of aflatoxin from cow feed to milk by addition of activated carbons. *J. Food Prot.* 59: 551-554.

- Hendricks, J.D. 1994. Carcinogenicity of aflatoxins in nonmammalian organisms. In: *The Toxicology of Aflatoxins: Human Health, Veterinary, and Agricultural Significance*. D.L. Eaton and J.D. Groopman, eds., Academic Press, New York, pp. 103-136.
- Huwig, A., Freimund, S., Kappeli, O. and Dutler, H. 2001. Mycotoxin detoxification of animals feed by different adsorbents. *Toxicol. Letters*. 122: 179-188.
- Karlovsky, P. 1999. Biological detoxification of fungal toxins and its use in plant breeding, feed and food production. *Nat. Toxins* 7: 1-23.
- Kececi, T., Oguz, H., Kurtoglu, V. and Demet, O. 1998. Effect of polyvinylpoly- pyrrolidone, synthetic zeolite and bentonite on serum biochemical and haematological characters of broiler chickens during aflatoxicosis. *Br. Poult. Sci.* 39: 452-458.
- Miazzo, R.D., Rosa, C.A.R., Carvalho, E.C.Q., Magnoli, C., Chiacchiera, S.M., Palacio, G., Saenz, M., Kikot, A., Basaldella, E. and Dalcerro, M. 2000. Efficacy of synthetic zeolites to reduce the toxicity of aflatoxin in broiler chicks. *Poultry Sci.* 79: 1-6.
- Phillips, T.D. 1999. Dietary clay in the chemoprevention of aflatoxin-induced disease. *Toxicol. Sci.* 52(Suppl): 118-126.
- Phillips, T.D., Kubena, L.F., Harvey, R.B., Taylor, D.R. and Heidelbaugh, N.D. 1987 Mycotoxin hazards in agriculture: New approach to control. *J. Am. Vet. Med. Assoc.* 190: 1617.
- Phillips, T.D., Kubena, L.F., Harvey, R.B., Taylor, D.R. and Heidelbaugh, N.D. 1988. Hydrated sodium calcium aluminosilicate: A high affinity sorbent for aflatoxin. *Poultry Sci.* 67: 243-247.
- Raju, M.V.L.N. and Devegowda, G. 2000. Influence of esterified β -glucomannan on performance and organ morphology, serum biochemistry and haematology in broilers exposed to individual and combined mycotoxicosis (aflatoxin, ochratoxin and T-2 toxin). *Br. Poult. Sci.* 41: 640-650.
- Ramos, A.J., Fink-Gremmels, J. and Hernandez, E. 1996. Prevention of toxic effects of mycotoxins by means of non-nutritive adsorbent compounds. *J. Food Prot.* 59: 631-641.
- Ramos, A.J. and Hernandez, E. 1997. Prevention of aflatoxicosis in farm animals by means of hydrated sodium calcium aluminosilicate addition to feed stuffs: a review. *Anim. Feed Sci. Tech.* 65: 197-206.
- Rosa, C.A.R., Miazzo, R., Magnoli, C., Salvano, M., Chiacchiera, S.M., Ferrero, S., Saenz, M., Carvalho, E.C.Q. and Dalcerro, A. 2001. Evaluation of the efficacy of bentonite from the South Argentina to ameliorate the toxic effects of aflatoxins in broilers. *Poultry Sci.* 80: 139-144.
- Sabater, V.M. 2003. Assessment and intervention of food- and feed-borne mycotoxicoses. Ph.D. Thesis, Department of Veterinary Pharmacology, Pharmacy and Toxicology, Utrecht University, Utrecht, The Netherlands.
- Schell, T.C., Lindemann, M.D., Kornegay, E.T., Blodgett, D.J. and Doerr, J.A. 1993. Effectiveness of different types of clay for reducing the detrimental effects of aflatoxin-contaminated diets on performance and serum profiles of weanling pigs. *J. Anim. Sci.* 71: 1226-1231.
- Wogan, G.N. and Newberne, P.M. 1967. Dose-response characteristics of aflatoxin B₁ carcinogenesis in rats. *Cancer Res.* 27: 2370-2376.