

1-1-2012

THE CHARACTERIZATION OF A NOVEL BACTERIOCIN WITH ANTIMRSA ACTIVITY PRODUCED BY BACILLUS SP. STRAIN WARY9-1M

Siripom Kaewklom

Ratchaneewan Aunpad

Follow this and additional works at: <https://digital.car.chula.ac.th/tjps>



Part of the [Pharmacology Commons](#)

Recommended Citation

Kaewklom, Siripom and Aunpad, Ratchaneewan (2012) "THE CHARACTERIZATION OF A NOVEL BACTERIOCIN WITH ANTIMRSA ACTIVITY PRODUCED BY BACILLUS SP. STRAIN WARY9-1M," *The Thai Journal of Pharmaceutical Sciences*: Vol. 36: Iss. 0, Article 24.

Available at: <https://digital.car.chula.ac.th/tjps/vol36/iss0/24>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Pharmaceutical Sciences by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

THE CHARACTERIZATION OF A NOVEL BACTERIOCIN WITH ANTI-MRSA ACTIVITY PRODUCED BY *BACILLUS* SP. STRAIN WARY9-1M

Siriporn Kaewklom, and Ratchaneewan Aunpad*

^oGraduate Program in Biomedical Sciences, Faculty of Allied Health Sciences, Thammasat University, Klonglaung, Pathumthai 12121, Thailand

^{*}Corresponding author; Tel: +662-986-9213-9 ext. 7269; E-mail: Aunpad@yahoo.com

KEYWORDS: Bacteriocin, anti-MRSA, *Bacillus* sp., WARY9-1M

INTRODUCTION

The emergence and dissemination of antibiotic resistance pathogenic bacteria such as methicillin resistant *Staphylococcus aureus* (MRSA) becomes an increasing serious problem in the public health worldwide (Schmitz *et al.*, 1998). Moreover, MRSA strains tend to accumulate additional new antibiotic resistance such as mupirocin (Krishnan *et al.*, 2002). New strategies for controlling MRSA and multiresistant staphylococci are urgently needed. Many substances used in pharmaceutical industries have been isolated from microorganisms. It is promising to use antibacterial compound known as bacteriocins to replace currently used antibiotics. Bacteriocins are ribosomally synthesized antimicrobial protein or peptide produced by bacteria that usually inhibit closely related species (Klaenhammer, 1988). Current potential use of bacteriocins are in food industry as natural and safe food preservatives while less research has been conducted on the therapeutic applications as antimicrobial agent (Gray *et al.*, 2006). Bacteriocin production is common among many gram-positive bacteria (Pattnaik *et al.*, 2005) especially soil bacteria in the genus *Bacillus*. Like LAB, the genus *Bacillus* includes a variety of industrially species which has been granted Generally Recognized as Safe (GRSA) status by the Food and Drug Administration, USA (Cherif *et al.*, 2003). In a previous research, strain WARY9-1M showing high antimicrobial activity against MRSA was isolated (Aunpad *et al.*, 2007). The objective of this study is to characterize the bacteriocin from bacteriocin-producing bacteria (BAC) strain WARY9-1M. This strain is not hazard and might be suitable for using as an alternative source of peptide antibiotic for control of many important antibiotic resistant pathogenic bacteria in the future.

MATERIALS AND METHODS

Bacterial strain identification

The isolated strain WARY9-1M was identified according to its 16S rRNA gene sequence (Brosius *et al.*, 1978). The nucleotide sequence was compared with genbank nucleotide database using Blastn search (<http://www.ncbi.nlm.nih.gov/blastn>) in order to identify the strain.

Determination of bacteriocin activity

The antibacterial activity of bacteriocin was detected by an agar-well diffusion method (Tagg and Mac-Given, 1971) and bacteriocin activity (AU/ml) was determined by the serial dilution method (Jansen and Hirschmann, 1994). The assay for each sample was done in triplicate.

Preparation of partially purified bacteriocin (PPB)

A 200 ml TSB medium was inoculated with 1% (10^6 CFU/ml) of an overnight culture of strain WARY9-1M. The cultures were incubated at 37°C for 16-18 h with shaking. Following cultivation, cell-free culture supernatant was obtained through centrifugation at 8,000 x g for 20 min (Sorvall Biofuge, Mandel Scientific, Canada) followed by sterile filtration. Ammonium sulfate (103.2 g) was added to the supernatant while stirring to reach 80% saturation and left overnight at 4°C. The sample was centrifuged at 8000 x g for 50 min. Then the supernatant was discarded and the precipitate was dissolved in 5 ml of sterile distilled water and dialyzed against 1.5 l of sterile distilled water for 16-18 hour. The active supernatant was designated as *partially purified bacteriocin or PPB*.

Spectrum of inhibitory activity

The PPB was used to assess the antibacterial activity against a total of 21 selected Gram-positive and Gram-negative test bacteria by the agar-well diffusion method (Tagg and Mac-Given, 1971). Equal volume of sterile distilled water was used as control solution. The appearance of the inhibition zone was determined after 18 h of incubation.

Enzyme sensitivity, heat and pH stability

The PPB was treated at 37°C for 1 h with 1 mg/ml final concentration of the following enzymes: trypsin, α -chymotrypsin and proteinase K (Sigma-Aldrich, USA). After incubation, the reaction mixtures were boiled for 10 min to inactivate the enzymes and the residual antibacterial activity was measured by agar-well diffusion. Thermal stability of bacteriocin was investigated by determination of the residual antibacterial activity after incubation of PPB at different temperature ranging from 40-100°C for 30 and 60 min, and at 121°C for 15 min. To investigate the effect of pH, antibacterial activity was measured

following the pH adjustment of the bacteriocin with 0.1 N NaOH or 0.1 N HCl and incubation at 4°C for 1 h.

PCR detection of antimicrobial peptide encoding gene

The genomic DNA of strain WARY9-1M was extracted with E.Z.N.A Bacterial DNA kit (Omega Biotek, USA). Six pairs of primer for amplification of genes involved in synthesis of antimicrobial peptides, lichenysin A, B, C, surfactin, subtilosin and iturin were used in this study. PCR amplified products after agarose gel electrophoresis were purified by E.Z.N.A. Gel Extraction Kit (Omega Biotek, USA) and ligated with TA cloning vector (Vivantis, Malaysia). The sequences were determined using M13F forward primer. The nucleotide sequence was compared with GenBank nucleotide database using Blastn search (<http://www.ncbi.nlm.nih.gov/blastn>).

Molecular weight determination

The molecular weight of bacteriocin from isolated strain WARY9-1M were determined by the Tris-Tricine SDS-PAGE with 5% stacking gel and 16% separating gel (Schagger & von Jagow, 1987). The first half of the gel (protein gel) was stained with PageBlue Protein staining solution (Fermentas, USA) whereas another half of the gel (activity gel) was washed in sterile distilled water for 30 min and overlaid with TSA (0.8% agar) seeded with 1% (v/v) MRSA and incubated at 37°C for 16-18 h. The formation of clear halo or inhibition zone was observed and compared with protein gel.

RESULTS

The isolated strain WARY9-1M was identified as *Bacillus amyloliquifaciens* with 100% identity according to their partial 16S rRNA gene sequence. The antimicrobial activity spectrum of partially purified bacteriocin (PPB) from strain WARY9-1M against 17 test microorganisms was examined by agar diffusion method. The PPB obtained from strain WARY9-1M showed broad antimicrobial activity against both gram-negative and gram-positive test microorganisms. Based on the size of the clear zone, the most sensitive strains were MRSA and *S. aureus*. The complete inactivation was observed after treatment with all proteolytic enzymes (trypsin, α -chymotrypsin and proteinase K), a key criterion for bacteriocin characterization (Table 1). Temperature stability experiment revealed that PPB was completely stable at high temperature up to 100°C for 60 min (Table 1). With regard to pH sensitivity, antibacterial activity of PPB was maintained at high level within the pH range of 3.0-9.0 (Table 1). PCR was used to analyze for genes involved in the biosynthesis of antimicrobial substances reported to be produced by *Bacillus* sp. PCR analysis of the *B. amyloliquifaciens* strain WARY9-1M showed that the strain exhibited potential for the functional gene encoding malonyl CoA transacylase (*ituD*) and lichenysin synthetase C (*lchAC*) (Figure 1). Sequence of 1,203 bp fragments showed high similarity (99% identity) to malonyl CoA transacylase of *Bacillus amyloliquifaciens* subsp. *plantarum* and only point mutations were observed. The translated nucleotide sequence of *ituD* showed FabD S-malonyltransferase conserved domain at N-terminal part. The molecular weight of bacteriocin from isolated *B. amyloliquifaciens* strain WARY9-1M was determined by Tris-Tricine SDS-PAGE analysis of PPB. As shown in Figure 2, a single protein band with clear halo revealed a bacteriocin activity. The band had an apparent molecular mass of 3.14 kDa.




Figure 1. PCR products profiles of strain WARY9-1M. Lane 1: 100 bp DNA marker, Lane 2 and 3: PCR products obtained from primers specific to iturin A gene and lichenysin synthetase C gene, respectively.

DISCUSSION

The number of methicillin resistant *Staphylococcus aureus* or MRSA infections has been increasing and become a serious problem in public health worldwide. Novel antibacterial agents are urgently needed to combat this drug resistant problem. The use of bacteriocin as an alternative agent to overcome the problem is promising (Papagianni 2003). The isolated strain WARY9-1M showing high anti-MRSA was identified as *Bacillus amyloliquefaciens*. It is well documented that bacteriocin or bacteriocin-like production is common among different *Bacillus* species. Most of them can inhibit only gram positive bacteria and less effective against gram negative strains. The partially purified bacteriocin (PPB) prepared from strain WARY9-1M was active against both gram positive and gram negative bacteria under investigation with high antibacterial activity against MRSA and *S. aureus*. The sensitivity of PPB to proteinase K, trypsin and α -chymotrypsin suggests the proteinaceous nature of this antimicrobial substance. Therefore, this isolated bacterium can be regarded as bacteriocin-producing bacteria or BAC. The bacteriocin from strain WARY9-1M was heat stable as evidenced by its ability to reserve the activity at 100°C for 60 min. Moreover, it was stable within a wide range of pH (3-9). The heat stable property was also observed in two bacteriocins, *i.e.*, entomocin 9 and AMS T6-5, produced by *B. thuringiensis* HD9 (Cherif *et al.*, 2003) and *B. licheniformis* T6-5 (Korenblum *et al.*, 2005), respectively. It was shown by PCR that strain WARY9-1M harbors genes for producing iturin A and lichenysin C. Iturin A and lichenysin C are bacterial cyclic lipopeptides with molecular mass of 1.058 and 1.000 kDa, respectively (Benitez *et al.*, 2010). The molecular weight of bacteriocin from strain WARY9-1M as determined by Tris-Tricine SDS-PAGE analysis was 3.14 kDa. The molecular mass of subtilosin reported to be produced by *B. amyloliquefaciens* was 3.399 (Marx *et al.*, 2001). However, the strain WARY9-1M did not harbor the gene encoding for subtilosin as analyzed by PCR. This bacteriocin produced by strain WARY9-1M might be a novel bacteriocin.

Table 1 Effect of enzymes, temperature and pH on PPB of strain WARY9-1M

Treatments and conditions	Residual activity (%)
None (control)	100
Enzyme treatment	
Trypsin	0
α -chymotrypsin	20
Proteinase K	20
Temperature	
40° C, 30 min	100
40° C, 60 min	100
60° C, 30 min	100
60° C, 60 min	100
80° C, 30 min	100
80° C, 60 min	100
100° C, 30 min	100
100° C, 60 min	100
121° C, 15 min	0
pH	
3.0	80
4.0	90
5.0	100
6.0	100
7.0	100
8.0	100
9.0	100

CONCLUSION

In conclusion, *B. amyloliquefaciens* strain WARY9-1 M produced bacteriocin with high anti-MRSA activity. This strain is non-pathogenic and derived from the nature. The biochemical properties such as thermal stability and wide range pH stability are remarkable. The bacteriocin produced by this microorganism might be used as an alternative source for controlling MRSA in the future.




Figure 2. Tris-Tricine SDS PAGE analysis of PPB prepared from *B. amyloliquefaciens* strain WARY9-1M (A) Coomassie brilliant blue stained gel (B) The activity gel shows the clear zone (arrow) after overlaid with TSA (0.8% agar) seeded with MRSA and incubated overnight. Lane 1: Peptide molecular weight marker; lane 2 and 3: PPB

ACKNOWLEDGMENTS

This work was supported by grants from Higher Education Research Promotion and National Research University Project of Thailand, Office of the Higher Education Commission and National Research Council of Thailand.

REFERENCES

- Aunpad R, Na-Bangchang K, Pipatsatitpong D (2007). Bacteriocins with anti-MRSA activity produced by water and soil isolated bacteria. *Ann. Microbiol.*, 57: 9-14.
- Benitez LB, Velho RV, Medina LFC, Brandelli A (2010). Isolation and characterization of antifungal peptides produced by *Bacillus amyloliquefaciens* LBM 5006. *J. Microbiol.*, 48: 791-797.
- Brosius J, Palmer LM, Kennedy PJ, Noller H (1978). Complete nucleotide sequence of 16S ribosomal RNA gene from *Escherichia coli*. *Proc. Nat. Acad. Sci.*, 75: 4801-4805.
- Cherif A, Chehimi S, Limem F, Hansen BM, Hendriksen NB, Daffonchio D, Boudabous A (2003). Detection and characterization of the novel bacteriocin entomocin 9 and safety evaluation of its producer, *Bacillus thuringiensis* ssp. *entomocidus* HD9. *J. Appl. Microbiol.*, 95: 990-1000.
- Gray EJ, Lee KD, Souleimanove AM, Falco MR, Zhou X, Ly A, Charles TC, Driscoll BT, Smith DL (2006). A novel bacteriocin, thuricin 17, produced by plant growth promoting rhizobacteria strain *Bacillus thuringiensis* NEB17: isolation and classification. *J. Appl. Microbiol.*, 100: 545-554.
- Jansen EF, Hirschmann DJ (1994). Subtilin, an antibacterial product of *Bacillus subtilis*, culturing conditions and properties. *Arch. Biochem.*, 4: 297-309.
- Klaenhammer TR (1988). Bacteriocins of lactic acid bacteria. *Biochemie.*, 70: 337-349.
- Korenblum E, von der Weid I, Santos ALS, Rosado AS, Sebastian GV, Coutinho CMLM, Magalhaes FCM, de Paiva MM, Seldin L (2005). Production of antimicrobial substances by *Bacillus subtilis* LFE-1, *B. firmus* H2O-1 and *B. licheniformis* T6-5 isolated from an oil reservoir in Brazil. *J. Appl. Microbiol.*, 98: 667-675.
- Krishnan PU, Miles K, Sketty N (2002). Detection of methicillin and mupirocin resistance in *Staphylococcus aureus* isolates using conventional and molecular methods: a descriptive study from a burns unit with high prevalence of MRSA. *J. Clin. Pathol.*, 55: 745-748.
- Marx R, Stein T, Entian K.-D, and Glaser SJ. (2001). Structure of the *Bacillus subtilis* peptide antibiotic subtilisin A determined by 1H-NMR and matrix assisted laser desorption/ionization time-of-flight mass spectrometry. *J. Prot. Chem.*, 20: 501-506.
- Pattnaik P, Grover S, Batish VK. (2005). Effect of environmental factors on production of lichenin, a chromosomally encoded bacteriocin-like compound produced by *Bacillus licheniformis* 26L-10/3RA. *Microbiol. Res.*, 160: 213-218.
- Papagianni M (2003). Ribosomally synthesized peptides with antimicrobial properties: biosynthesis, structure, function, and applications. *Biotechnol. Adv.*, 21: 465-499.
- Schmitz FJ, Lindenlauf E, Hofman B, Fluit AC, Verhoef J, Heinz HP, Jones ME (1998). The prevalence of low- and high-level mupirocin resistance in *Staphylococci* from 19 European hospitals. *J. Antimicrob. Chemother.*, 42: 489-495
- Tagg JR, Mac-Given AR (1971). Assay system for bacteriocin. *Appl. Microbiol.*, 21: 943.