

3-1-2011

## Presence of Infectious Pancreatic Necrosis Virus on Rainbow Trout (*Oncorhynchus mykiss*) by Histopathology, ELISA and RT-PCR

Gabriel Aguirre-Guzmán

Ned Iván Iván de la Cruz-Hernández

Jesús Genaro Sánchez-Martínez

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

---

### Recommended Citation

Aguirre-Guzmán, Gabriel; Cruz-Hernández, Ned Iván Iván de la; and Sánchez-Martínez, Jesús Genaro (2011) "Presence of Infectious Pancreatic Necrosis Virus on Rainbow Trout (*Oncorhynchus mykiss*) by Histopathology, ELISA and RT-PCR," *The Thai Journal of Veterinary Medicine*: Vol. 41: Iss. 1, Article 19. Available at: <https://digital.car.chula.ac.th/tjvm/vol41/iss1/19>

This Short Communication is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact [ChulaDC@car.chula.ac.th](mailto:ChulaDC@car.chula.ac.th).

# **Presence of Infectious Pancreatic Necrosis Virus on Rainbow Trout (*Oncorhynchus mykiss*) by Histopathology, ELISA and RT-PCR**

**Gabriel Aguirre-Guzmán\* Ned Iván de la Cruz-Hernández  
Jesús Genaro Sánchez-Martínez**

## ***Abstract***

Trout (*Oncorhynchus mykiss*) culture in Mexico experiences sporadic and significant fish mortalities, where disease signs are associated with infectious pancreatic necrosis virus disease (IPNV). The purpose of the present work was to support the potential of ELISA, histopathology, and RT-PCR as routine techniques for IPNV detection. Fourteen trout farms were monitored after a disease outbreak. Positive results were confirmed in different collected organisms from 21% of tested farms by all 3 techniques. Virus detection by ELISA and RT-PCR was successfully performed in only two days after the initial signs of disease were observed in fish, since these methods are faster when compared to histopathology, which only detected signs of disease such as necrosis of the pancreatic acinus and intestine after 5 days. The results suggest that ELISA and RT-PCR offer an early, sensitive, faster and cheaper alternative for the routine detection of IPNV compared to cell culture and histopathology. These techniques can be performed for fish health monitoring and are reliable tools in the control, prevention and timely detection of IPNV.

---

**Keywords:** ELISA, infectious pancreatic necrosis, trout, reverse transcription polymerase chain reaction

---

<sup>1</sup>Facultad de Medicina Veterinaria y Zootecnia. Universidad Autónoma de Tamaulipas. Km. 5 Carr. Cd. Victoria - Mante,

Ciudad Victoria, Tamaulipas, México.

Corresponding author E-mail: gaguirre\_guzman@hotmail.com

## บทคัดย่อ

### การพบ Infectious Pancreatic Necrosis Virus ในปลาเรนโบว์เทราท์ (*Oncorhynchus mykiss*) โดยวิธีทางจุลพยาธิวิทยา ELISA และ RT-PCR

Gabriel Aguirre-Guzmán<sup>1\*</sup> Ned Iván de la Cruz-Hernández<sup>1</sup> Jesús Genaro Sánchez-Martínez<sup>1</sup>

อุตสาหกรรมการเพาะเลี้ยงปลาเรนโบว์เทราท์ (*Oncorhynchus mykiss*) ในประเทศเม็กซิโก เคยประสบปัญหาการตายของปลา ซึ่งพบว่าอาการดังกล่าวมีความสัมพันธ์กับโรคติดเชื้อไวรัส infectious pancreatic necrosis virus disease (IPNV) วัตถุประสงค์ของการศึกษานี้เพื่อศึกษาสภาพของวิธีการวินิจฉัยโรค IPNV โดยวิธี ELISA จุลพยาธิวิทยา และ RT-PCR. โดยทำการศึกษาในฟาร์มปลาเรนโบว์เทราท์ จำนวน 14 ฟาร์มที่เคยพบการระบาดของโรค พบว่าร้อยละ 21 ของตัวอย่างทั้งหมดที่เก็บจากฟาร์มให้ผลบวกทั้ง 3 วิธี การตรวจหาแอนติเจนของไวรัส โดยวิธี ELISA และ RT-PCR สามารถที่ให้ผลลบภายในวันที่ 2 ของวันที่ปลาเริ่มป่วย ซึ่งทำการยืนยันผลโดยวิธีทางจุลพยาธิวิทยาที่สามารถตรวจพบรอยโรคในวันที่ 5 จากผลการศึกษาพบว่าวิธี ELISA และ RT-PCR มีความไว รวดเร็ว และเป็นแนวทางเลือกที่ดีในการวินิจฉัยโรค IPNV เมื่อเปรียบเทียบกับเพาะแยกเชื้อไวรัส และการตรวจรอยโรคทางจุลพยาธิวิทยา ซึ่งสามารถนำมาใช้ดังกล่าวมาใช้ในการเฝ้าระวัง ควบคุมและป้องกันโรคในฟาร์มปลาได้เป็นอย่างดี

**คำสำคัญ:** ELISA, infectious pancreatic necrosis, ปลาเรนโบว์เทราท์, reverse transcription polymerase chain reaction

<sup>1</sup> Facultad de Medicina Veterinaria y Zootecnia. Universidad Autónoma de Tamaulipas. Km. 5 Carr. Cd. Victoria - Mante, Ciudad Victoria, Tamaulipas, México.

\*ผู้รับผิดชอบบทความ E-mail: gaguirre\_guzman@hotmail.com

## Introduction

Infectious pancreatic necrosis virus (IPNV) of the Aquabirnavirus genus and Birnaviridae family is a well-known pathogen of rainbow trout (*O. mykiss*). This virus is one of the major etiological agents found in farms and wild salmonid fish, and its genome has a double-stranded RNA packaged in a nonenveloped icosahedral shell, 60 nm in diameter (Cutrín et al., 2000; Zhang and Suzuki, 2004). This is a highly contagious disease initially detected in North America (Roberts and Pearson, 2005) with mortalities ranging from 70-100% in fingerlings and early juvenile fish, usually following culture stress (Evensen and Lorenzen, 1997). The susceptibility to this virus decreases with age, but asymptomatic organisms facilitate horizontal and vertical transmission (Brown and Bruno, 2002). Clinical signs include skin hyperpigmentation, distended abdomen with ascites, empty gut, presence of clear or milky mucus in stomach and anterior intestine, long, thin white fecal casts, spiral swimming, pale spleen, heart, liver, and petechiae observed in the viscera (Crane et al., 2000; Brown and Bruno, 2002).

Rainbow trout farming in Mexico is a growing and new industry whose production increased from 1612 to 11,792 metric tons in one decade (1998-2008); with most of the rainbow trout farms and hatcheries located in Chihuahua, Durango, Mexico, Michoacán, Morelos, and Puebla (CONAPESCA 2008). However, other states such as Hidalgo, Jalisco, Nuevo León, and San Luis Potosi have small scale extensive farms with potential to increase their production because of their association to eco-tourism and proximity to big cities. The first report of IPNV in Mexico occurred in 2002, from *O. mykiss* samples obtained from a farm located in Central Mexico (Ortega et al., 2002); however, IPNV distribution in farm and wild stock in Mexico has been little understood and poorly evaluated (Barrera-Mejia et al., 2002).

The standard protocol for the routine detection of IPNV is usually based on virus isolation (Milne et al., 2006), complemented by histopathology, immunofluorescence, serology, and molecular biology (Barrera-Mejia et al., 2002; Bowden et al., 2002; Jencić et al., 2002). Viral isolation is considered as the gold standard, but it is laborious, expensive and time consuming (Kerr and Cunningham, 2006; Milne et al.,


2006). The purpose of the present study was to evaluate the use of RT-PCR and ELISA for the routine diagnosis of IPNV in aquatic health labs.

### Materials and Methods

Fourteen rainbow trout farms in the Mexican state of Hidalgo (Fig. 1) were monitored for the presence of IPNV virus. Four hundred and ninety six fish (*O. mykiss*) showing clinical signs such as erratic swimming, loss of appetite, skin darkening, distended abdomen, exophthalmia, and long thin white fecal casts were collected. Fingerlings and early juveniles were collected without dissection and immediately submerged into 10% neutral buffered formalin (Evensen and Lorenzen 1997). Older juveniles were dissected in situ and their gills, brain, heart, spleen, kidney, liver, pancreas, and intestine samples were collected (Barlic-Maganja et al., 2002; Kerr and Cunningham, 2006). The ELISA samples were washed with sterile PBS and stored at -80°C until used. Samples for RT-PCR were washed with PBS and placed in RNAlater® (Ambion, USA) for storage (Phelan et al., 2005).

The fixed samples (whole organisms or tissues) were processed for routine histology; embedded in paraffin according to standard procedures, sectioned at 4-6 µm, stained with H&E. The samples were examined by light microscopy (Carl Zeiss/Axiostar) coupled to a digital camera (Canon, Powershot G6 PC1089) to look for IPNV-associated morphologic changes such as necrotic lesions and ulcers in the pancreatic acini, esophagus, stomach, intestine, and renal hematopoietic elements (Bowden et al., 2002; Roberts and Pearson, 2005).


Fish samples from the same pond were pooled in groups of three or four organs, and homogenized with a pellet mixer (VWR, USA) at 200 rpm for 1 min with sterile PBS, and independently analyzed. All samples were tested with an ELISA kit for IPNV (Test-line®, Czech Republic) according to




**Figure 1** Trout (*Oncorhynchus mykiss*) farm locations in Hidalgo state, Mexico, and yearly aquaculture yield.

the manufacturer's instructions (Bowden et al., 2002; Jenčić et al., 2002). The samples were analyzed with a Bio-Rad (USA) microplate reader at 415 nm, and were positive when absorbance was greater than 0.5. Positive controls used were supplied by Laboratorio de Biotecnología 1, Facultad de Ciencias Químicas, Universidad Autónoma de Chihuahua (LB1-FCQ-UACH), and the test-line kit.

Rainbow trout samples (whole organisms or tissues) were sent for RT-PCR analysis to the LB1-FCQ-UACH. All samples were pooled in groups of three or four organs, homogenized with a pellet mixer at 200 rpm for 1 min with sterile RNAlater® at 4°C. All samples were processed with a RNA isolation kit (Quantum Prep AquaPure Genomic) according to the manufacturer's instructions (Holmes et al., 2003). Reverse transcription reactions were performed using the protocol of SuperScript™ II ARNse reverse transcriptase (Invitrogen Inc., USA) (Cutrín et al., 2000). For IPNV detection, PNF primers [ATCTGCGGTGTAGACATCAAA (Forward) and TGCAGTTCCTCGTCCATCCC (Reverse)] (Taksdal et al., 2001; Barrera-Mejia et al., 2002) were used for the PCR reaction where the visualization of a 224 bp band was considered as a positive result (Fig. 2). PCR was carried out in a total volume of 25 µl of FSB buffer (10 mM dNTP's, 50 mM Tris-HCl, 75 mM KCl, 3 mM MgCl<sub>2</sub>, pH 8.3) containing 15 pmol of each primer, and 2.5 U of Taq polymerase (Promega, Inc.). The PCR reactions were carried out in a thermocycler (Corbett Research) under the following conditions: 96°C for 5 min followed by 94°C for 60 sec (denaturalization), 55°C for 30 sec (annealing) and 72°C for 60 sec (extension). After 35 cycles, the reactions were cooled down to 4°C, and PCR products were analyzed on 2% agarose gel in 1x TBE buffer (89 mM Tris HCl, 89 mM boric acid, 2 mM EDTA, pH 8) at 80 V for 60 min. The gels were stained with ethidium bromide, and analyzed with imaging system software from Kodak (Model Gel logic-200).



**Figure 2** IPNV amplification from trout samples in 2% agarose gel electrophoresis in 1x TBE at 80 volts by 60 min. MWM: molecular weight marker; Lane 1 and 2: negative control (trout tissue sample and distillate sterile water, respectively); Lane 3: positive control from LB1-FCQ-UACH; Lane 4-6: trout samples.



**Figure 3** Histological sections from *Oncorhynchus mykiss*. (A) Acute necrosis in pancreas (arrow) (H&E, 4x). (B) Generalized necrosis in intestine (arrow) and cells with nuclear pyknosis and karyorrhexis (arrow head) (H&E, 40x).

### Results and Discussion

The detection protocols for different pathogens (virus, bacteria, fungi, etc.) on aquatic organisms help to increase the knowledge about their presence, biogeographical distribution, effects, and control on aquaculture and fisheries production settings (Ghittino et al., 2003). In Mexico, rainbow trout production areas have shown a sudden and significant increase in mortality in fingerlings and early-juveniles during 2003-2006 (CONAPESCA 2008), which had a direct effect on fish farms. The international diagnosis protocol for IPNV is based on virus isolation by cell culture; which is a laborious, expensive and time-consuming activity. Histology, immunofluorescence, immunohistochemistry (Ellis et al. 2010), ELISA (Bowden et al., 2002; Jencič et al., 2002), and RT-PCR (Taksdal et al., 2001; Barrera-Mejia et al., 2002) are other methods that have been suggested as complementary for IPNV detection, because of their sensitivity, speed and economy.

In this study, the organisms collected (496) from rainbow trout farms (14) consisted of fingerlings (13%), early-juveniles (26%), juveniles (52%) and adults (9%). Significant histopathological lesions were observed in the different fish samples examined; however, only samples from farms 1, 4 and 8 showed lesions associated to IPNV, such as pancreatic necrosis, necrotizing enteritis; mucosal epithelial necrosis and catharrhal enteritis (Fig 3). Renal tissues displayed moderate degenerative tubules with necrotic areas (Brown and Bruno, 2002; Roberts and Pearson, 2005). However, the usefulness of histopathology in IPNV diagnosis is only effective when: i) a specific sign is associated to a specific pathogen; ii) this specific sign is expressed at a high enough level; and iii) the aquatic pathologist experience. The histopathological changes observed in the collected samples were not exclusive of IPNV diseases and the use of other diagnostic protocols were necessary for a confirmed diagnosis of the specific pathogen affecting the organism.

Table 1 shows the results of the ELISA and RT-PCR analyses for all trout samples. It can be seen that samples from three farms (1, 4, and 8; = 21% of

tested farms) were positive to IPNV (100%, pool of samples) by ELISA (1.12, 1.17, and 1.13 of S/P ratio mean, respectively) and RT-PCR, in a much shorter time frame compared to histopathology and exhibited a higher sensitivity to IPNV detection.

Prevention of pathogen exposure is the most effective way of controlling disease, with farms including routine diagnostic examinations for their aquatic health programs. These techniques may represent an advantage for the routine and timely diagnosis of IPNV in trout culture and wild stock studies, since early pathogen identification is crucial in the prevention of spread of infection, and in decision making for treatment and disease control (Barrera-Mejia et al., 2002), which represent possible effective containment and elimination of an emergency disease outbreak and epidemic.

**Table 1** ELISA and RT-PCR results from trout farm of Hidalgo state, Mexico.

Trout farm	Sample number	ELISA*				RT-PCR		
		K+	T+	W-	Sample	T+	W-	Sample
1	40	R	R	ND	R	+	-	+
2	8	R	R	ND	ND	+	-	-
3	5	R	R	ND	ND	+	-	-
4	14	R	R	ND	R	+	-	+
5	96	R	R	ND	ND	+	-	-
6	8	R	R	ND	ND	+	-	-
7	10	R	R	ND	ND	+	-	-
8	155	R	R	ND	R	+	-	+
9	14	R	R	ND	ND	+	-	-
10	16	R	R	ND	ND	+	-	-
11	16	R	R	ND	ND	+	-	-
12	16	R	R	ND	ND	+	-	-
13	60	R	R	ND	ND	+	-	-
14	38	R	R	ND	ND	+	-	-

\*Reactive (R) where the mean absorbance in wells with positive antigen is < 0.5. ND: not determined.

Control positive: T+ (sample from LB1-FCQ-UACH), K+ (control positive from test-line kit)

Control negative: W- (distilled water).

### Acknowledgements

This study was funded by grants from the Mexican Council of Science and Technology (CONACyT), project SAGARPA-CONACyT-2003-002-227). We thank Laboratorio de Biotecnología 1, Fac. de Ciencias Químicas, Universidad Autónoma de

Chihuahua (Dr. G. Erosa de la Vega) for supplying the positive control and the laboratory facilities for this work.

## References

- Barlic-Maganja, D., Strancar, M., Hostnik, P., Jencic, V. and Grom, J. 2002. Comparison of the efficiency and sensitivity of virus isolation and molecular methods for routine diagnosis of infectious haematopoietic necrosis virus and infectious pancreatic necrosis virus. *J Fish Dis.* 25: 73-80.
- Barrera-Mejia, M., Simon-Martinez, J., Salgado-Miranda, C., Vega, F., Ortega, C. and Aragon, A. 2002. Development and validation of a short-time cell culture and multiplex reverse transcriptase polymerase chain reaction assay for infectious pancreatic necrosis virus in mexican farm-sampled rainbow trout. *J Aquat Anim Health.* 21: 167-172.
- Bowden, T.J., Smail, D.A. and Ellis, A.E. 2002. Development of a reproducible infectious pancreatic necrosis virus challenge model for Atlantic salmon, *Salmo salar L.* *J Fish Dis.* 25: 555-563.
- Brown, L.L. and Bruno, D.W. 2002. Infectious diseases of coldwater fish in fresh water. In: *Diseases and Disorders of Finfish in Cage Culture*, P.T.K. Woo, D.W. Bruno and L.H.S. Lim (eds) CABI publishing, New York, 107-171.
- CONAPESCA (Comisión Nacional de Acuicultura y Pesca). 2008. *Anuario Estadístico de Pesca*.
- Crane, M.S.J., Hardy-Smith, P., Williams, L.W., Hyatt, A.D., Eaton, L.M., Gould, A., Handlinger, J., Kattenbelt, J. and Gudkovs, N. 2000. First isolation of an aquatic birnavirus from farmed and wild fish species in Australia. *Dis Aquat Org.* 43: 1-14
- Cutrín, J.M., Barja, J.L., Nicholson, B.L., Bandin, I., Blake, S. and Dopazo, C.P. 2000. Restriction fragment length polymorphisms and sequence analysis: an approach for genotyping infectious pancreatic necrosis virus reference strains and other aquabirnaviruses isolated from northwestern Spain. *Appl Environ Microbiol.* 70: 1059-1067.
- Evensen, O. and Lorenzen, E. 1997. Simultaneous demonstration of infectious pancreatic necrosis virus (IPNV) and *Flavobacterium psychrophilum* in paraffin embedded specimens of rainbow trout *Oncorhynchus mykiss* fry by use of paired immunohistochemistry. *Dis Aquat Org.* 29: 227-232.
- Ghittino, C., Latini, M., Agnetti, F., Panziera, C., Lauro, L., Ciappelloni, R. and Petracca, G. 2003. Emerging pathologies in aquaculture: Effects on production and food safety. *Vet Res Comm.* 27: 471-479.
- Ellis, A.E., Cavaco, A., Petrie, A., Lockhart, K., Snow, M. and Collet, B. 2010. Histology, immunocytochemistry and qRT-PCR analysis of Atlantic salmon, *Salmo salar L.*, post-smolts following infection with infectious pancreatic necrosis virus (IPNV). *J Fish Dis.* 33: 787-863.
- Holmes, S.P., Witbaard, R. and van der Meer, J. 2003. Phenotypic and genotypic population differentiation in the bivalve mollusc *Arctica islandica*: Results from RAPD analysis. *Mar Ecol Prog Ser.* 254: 163-176.
- Jenčič, V., Hostnik, P., Barlič Maganja, D. and Grom, J. 2002. The spread of salmonid viral diseases in Slovenia. *Slov Vet Res.* 39: 197-205.
- Kerr, C. and Cunningham, C. 2006. Moving molecular diagnostics from laboratory to clinical application: A case study using infectious pancreatic necrosis virus serotype A. *Lett Appl Microbiol.* 43: 98-104.
- Milne, S.A., Gallacher, S., Cash, P. and Porter, A.J.R. 2006. A reliable RT-PCR-ELISA method for the detection of infectious pancreatic necrosis virus (IPNV) in farmed rainbow trout. *J Virol Methods.* 132: 92-96.
- Ortega, C., Montes de Oca, R., Groman, D., Yason, C., Nicholson, B. and Blake, S. 2002. Viral infectious pancreatic necrosis in farmed rainbow trout from Mexico. *J Aquat Anim Health.* 14: 305-310.
- Phelan, P.E., Pressley, M.E., Witten, P.E., Mellon, M.T., Blake, S. and Kim, C.H. 2005. Characterization of snakehead rhabdovirus infection in zebrafish (*Danio rerio*). *J Virol.* 79: 1842-1852.
- Roberts, R.J. and Pearson, M.D. 2005. Infectious pancreatic necrosis in Atlantic salmon, *Salmo salar L.* *J Fish Dis.* 28: 383-390.
- Taksdal, T., Dannevig, B.H. and Rimstad, E. 2001. Detection of infectious pancreatic necrosis (IPN) virus in experimentally infected atlantic salmon parr by RT-PCR and cell culture isolation. *Fish Pathol.* 21: 214-218.
- Zhang, C.X. and Suzuki, S. 2004. Aquabirnaviruses isolated from marine organisms form a distinct genogroup from other aquabirnaviruses. *J Fish Dis.* 27: 633-643.

