

3-1-2014

Molecular Characterization and Expression Analysis of miR-29a in Porcine Cells and Porcine Reproductive and Respiratory Syndrome Virus Infected Peripheral Blood Mononuclear Cells

Win Surachetpong

Sunaree Nantakhruea

Porntippa Lekcharoensuk

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Surachetpong, Win; Nantakhruea, Sunaree; and Lekcharoensuk, Porntippa (2014) "Molecular Characterization and Expression Analysis of miR-29a in Porcine Cells and Porcine Reproductive and Respiratory Syndrome Virus Infected Peripheral Blood Mononuclear Cells," *The Thai Journal of Veterinary Medicine*: Vol. 44: Iss. 1, Article 4.

Available at: <https://digital.car.chula.ac.th/tjvm/vol44/iss1/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Molecular Characterization and Expression Analysis of miR-29a in Porcine Cells and Porcine Reproductive and Respiratory Syndrome Virus Infected Peripheral Blood Mononuclear Cells

Win Surachetpong* Sunaree Nantakhruea Porntippa Lekcharoensuk

Abstract

MicroRNAs are a group of small non-coding RNA that are involved in posttranscriptional regulation through target mRNA degradation or inhibition of protein synthesis. The objective of this present study was to characterize and determine expression level of porcine miR-29a in various tissues. Our results indicated that porcine miR-29a shared high conserved sequence with miR-29a from other organisms. Furthermore, *in silico* analysis suggested that the seed region of porcine miR-29a partially matched IFN- γ mRNA. RT-PCR analysis indicated that miR-29a expressed in various tissues including lymph node, spleen, peripheral blood mononuclear cells (PBMCs) and PK-15 cells. Expression profiling of miR-29a in PRRSV-infected PBMCs *in vitro* revealed an upregulation at 24 and 48 h. Moreover, expression analysis of miR-29a showed higher transcripts in both low pathogenic and highly pathogenic-PRRSV infected pigs. Together, all of these results revealed that porcine miR-29a was highly conserved with miR-29a from other organisms and miR-29a transcripts increased in PRRSV-infected PBMCs both *in vitro* and *in vivo*. Additional studies of the role of miR-29a will provide valuable information on virus: host interaction and could be applied for PRRS control in the future.

Keywords: immunity, miR-29a, porcine, PRRSV

Department of Veterinary Microbiology and Immunology, Faculty of Veterinary Medicine, Kasetsart University, Chatuchak, Bangkok 10900, Thailand

**Correspondence:* fvetswp@ku.ac.th

Introduction

Porcine reproductive and respiratory syndrome (PRRS) is an important viral disease affecting pigs worldwide. The causative agent is PRRS virus, which is a single stranded, positive sense RNA virus in the family of *Arteriviridae* (Frossard et al., 2013). Presently, two different genotypes of PRRSV, genotype I (European) and genotype II (North American), have been characterized since its initial outbreak. To avoid host immune response, PRRSV employs multiple strategies to inhibit host innate and adaptive immune response. For example, PRRSV non-structural proteins can interfere with type I IFN production (Sun et al., 2012^b). Moreover, the virus nucleocapsid protein can stimulate regulatory T cell differentiation and IL-10 production (Wongyanin et al., 2012). All of these mechanisms are responsible for poor adaptive immune response as suggested by delay neutralizing antibody production and less IFN- γ secreting cells in PRRSV-infected pigs (Labarque et al., 2000; Díaz et al., 2005). Such an inefficient immune response allows viral persistent in pigs up to 251 days (Wills et al., 2003).

MicroRNAs are a group of conserved small RNA with 22 nucleotide lengths that participate in the process of posttranscriptional regulation. MiRNAs have been identified across multiple organisms including plants, animals, human as well as viruses (Bartel, 2009). MiRNAs have been associated with a wide variety of biological processes such as oncogenesis, organ development, and host pathogen interaction (Krol et al., 2010). To regulate gene expression, specific sequence located within 5' untranslated region (UTR) of miRNA binds to target mRNA, leading to mRNA degradation or inhibition protein translation (Bartel, 2009). Recent studies clearly suggested that both innate and adaptive immunity were finely regulated by multiple miRNAs (Chen et al., 2013). Among these, members of the miR-29 family have been shown as important regulators of diverse immunological pathways (Liston et al., 2012; Brain et al., 2013). In particular, miR-29 targeted IFN- γ production by directly inhibiting IFN- γ mRNA (Ma et al., 2011; Savan et al., 2011) or indirectly regulating two transcription factors Eomes and T-bet (Steiner et al., 2011).

To date, there is no available information on miR-29a expression during viral infection in porcine cells. Based on the conserved sequence of miR-29a among different species and its critical role in immune system, it is interesting to characterize porcine miR-29a and determine expression level of miR-29a in normal porcine tissues and cell lines. In addition, analysis of miR-29a expression in PBMCs was investigated during PRRSV infection both *in vitro* and *in vivo*. Distinct expression pattern of miR-29a may provide valuable information to the role of this microRNA in pig.

Materials and Methods

Characterization of porcine miR-29a: To characterize porcine miR-29a, sequences of miR-29a were retrieved from miRbase (www.miRbase.org) and NCBI (www.ncbi.nlm.nih.gov). The *Sus scrofa* miR-29a (NR_038524) sequence was compared with miR-29a from other organisms including *Homo sapiens* (NR_029503), *Rattus norvegicus* (NR_031836), *Mus musculus* (NR_029744), *Bos taurus* (NR_031267), *Danio rerio* (NR_030039), and *Gallus gallus* (NR_031391). The secondary structure of precursor miRNA was predicted by RNA Folding Form, Mfold program (Zuker, 2003). Comparisons of miR-29a precursor and mature sequence among species were examined by multiple sequences alignment using ClustalW2 program, European Molecular Biology Laboratory-European Bioinformatics Institute (EMBL-EBI).

PBMC preparation and expression of miR-29 in normal cells and tissues: Tissues including spleen and mesenteric lymph nodes were collected from three healthy pigs. PBMC were isolated from heparinized whole blood samples using Ficoll-Hypaque Plus (GE healthcare, USA) density gradient centrifugation. Blood samples were initially diluted in sterile 1X phosphate buffer saline (PBS) (PBS : whole blood = 1 : 1). The diluted blood was overlaid on Ficoll-Hypaque Plus (diluted blood : Ficoll = 10 : 3) in 15 ml conical tube and centrifuged at 2500 rpm for 20 min at 20°C. After centrifugation, PBMCs layer was carefully collected into a new conical tube. To remove contaminated red blood cells, samples were incubated with 5 ml of sterile Red Blood cell (RBC) lysis solution

Table 1 All Primers used in this study

Target	Primer name	Primer sequence (5' - 3')	Tm (°C)	Size (bp)	GenBank Accession number
miR-29a	miR29A_F	CCCCTTAGAGGATGACTGAT	57.3	86	NR_038524.1
	miR29A_R	CCCCAATCATTATAACCGAT	53.2		
GAPDH	GAPDH_F	AGGTCATCCATGACAACCTCGGCA	72.5	155	AF017079
	GAPDH_R	AGCACCAGTAGAAGCAGGGATGAT	67.9		
β -actin	ACTB_F	GGAATTCGAGCAGGAGATGG	66.2	233	AY550069.1
	ACTB_R	GCACCGTGTGGCGTAGAGG	70.1		

Table 2 Secondary structure of precursor miR-29a from various organisms

Species	Secondary structure of miR-29a					
<i>Sus scrofa</i>	UUAGAGG	UUU	C	UCAAU		
	CCCC	AUGACUGAUUUC	UGGUGUU	AGAG	\	
	GGGG	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A	
	UUAGUAA	UCU	-	UUAUU		
<i>Homo sapiens</i>	UUU	C	UCAAU			
	AUGACUGAUUUC	UGGUGUU	AGAG	\		
	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A		
	UCU	-	UUAUU			
<i>Rattus norvegicus</i>	A	UUAGAGG	UUU	C	UCAAU	
	CCCC	AUGACUGAUUUC	UGGUGUU	AGAG	\	
	GGGG	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A	
	A	UUAGUAA	UCU	-	UUAAG	
<i>Mus musculus</i>	A	UUAGAGG	UUU	C	UCAAU	
	CCCC	AUGACUGAUUUC	UGGUGUU	AGAG	\	
	GGGG	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A	
	A	UUAGUAA	UCU	-	UUAAG	
<i>Bos taurus</i>	UUU	C	UCAAU			
	AUGACUGAUUUC	UGGUGUU	AGAG	\		
	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A		
	UCU	-	UUAUU			
<i>Danio rerio</i>	U	CCUCAUCUCUCUCUCUCUC	CCAAACG	CUU	U	GUCCCA
	GAAGA		UCCCCA	AUGACUGAUUUC	UGGUGCU	AGA
	CUUCU		AGGGGU	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCU
	A	UAACU	-----	CAGUAA-	UUU	-
						ACUGUC
<i>Gallus gallus</i>	A	UUUAGAGG	UUU	C	UCAAU	
	CCCC	AUGACUGAUUUC	UGGUGUU	AGAG	\	
	GGGG	UAUUGGCUAAAAG	<u>ACCACGA</u>	UCUU	A	
	A	UUAGUGA	-	UUU	-	UUAUA




Figure 2 Expression of miR-29a in normal porcine tissues including PBMCs, mesenteric lymph nodes and spleen. β -actin was used as internal control gene. M: 10 bp marker, N: no template control, P: positive control (pGEM-miR29a), PBMC: peripheral blood mononuclear cells, LN: mesenteric lymph nodes




Figure 3 RT-PCR analysis of miR-29a expression in PK-15 cell line. M: 10 bp marker, N: no template control and P: positive control (pGEM-miR29a)

from mesenteric lymph nodes, spleen and PBMCs. RT-PCR analysis revealed a single specific product of miR-29a at 86 bp. Interestingly, miR-29a was highly expressed in mesenteric lymph nodes and spleen, whereas the expression of miR-29a was low in PBMCs (Fig 2). In addition to the porcine tissues, we examined miR-29a expression in porcine kidney cell line (PK-15). As shown in Figure 3, miR-29a transcript could be detected in normal PK-15 cells. These findings indicated that miR-29a was expressed in both porcine immune and non-immune cells.

Expression of miR-29a in PRRSV-infected PBMCs: To determine the expression pattern of miR-29a during PRRSV infection, PBMCs were isolated from PRRSV-free pigs and plated in 96 well plate. The cells were then inoculated with LP-PRRSV at MOI 0.1 for 24 and 48 h. Expression of miR-29a was investigated by qRT-PCR method and normalized against GAPDH control gene. After PRRSV infection, we found that the expression of miR-29a increased 5.2 and 4.1 folds at 24 and 48 h, respectively, with higher expression at 24 h post infection (Fig 4). However, there was no statistical significant between control and PRRSV-infected groups ($p < 0.05$).

Expression of miR-29a in PRRSV-infected pigs: Based on the expression of miR-29a in PRRSV-

infected PBMCs *in vitro*, it is interesting to investigate level of miR-29a in PBMCs from PRRSV-infected animals. Although there is no statistical significance among the control, LP-PRRSV and HP-PRRSV infected animals ($p < 0.05$), we observed an increasing trend of miR-29a in both LP-PRRSV and HP-PRRSV infected pigs (Fig 5a). Specifically, the pigs infected with LP-PRRSV and HP-PRRSV has miR-29a transcript increase at 2.2 and 1.8 folds, respectively (Fig 5b).

Discussion

MicroRNAs are evolutionary conserved, non-coding RNAs that negatively regulates target gene expression at posttranscriptional process. To date, hundreds of microRNAs have been characterized in pigs during normal biological process and pathogen response (Xie et al., 2011; Podolska et al., 2012). In this study, we characterized porcine miR-29a and determined its expression level during normal condition and viral infection. Our results showed that porcine miR-29a shared high conserved sequence with miR-29a from other organisms (Xie et al., 2011; Chen et al., 2012). Furthermore, the sequence of porcine miR-29a seed region shared 100% sequence similarity to miR-29a seed region from human, mouse, rat, cow, chicken and fish. In human and mice, the seed region of miR-29a directly targeted 3'UTR of IFN- γ mRNA, leading to low IFN- γ production (Ma et al., 2011; Savan et al., 2011). Moreover, miR-29a regulated IFN- γ production in helper T cells through indirectly targeting two transcription factors Eomes and T-bet (Steiner et al., 2011).

Following the characterization of porcine miR-29a, the expression pattern of miR-29a was investigated in various porcine tissues (mesenteric lymph nodes, spleen and PBMCs) and PK-15 cell line. Our findings suggested that miRNA-29a expressed in all normal porcine tissues and cell lines with higher expression in spleen and lymph nodes than PBMCs. Similar to this finding, different expression patterns of miRNA in mice and human were reported in various tissues and correlated with function and disease pathogenesis (Lagos-Quintana et al., 2002; Liang et al., 2007).

One of the important viral diseases affecting pig production is PRRS (Neumann et al., 2005). To establish infection, PRRSV uses multiple strategies to suppress pig immune response including the delayed and weak IFN- γ production (Dotti et al., 2013). Based on the sequence similarity with other organisms and expression pattern of porcine miR-29a in stimulated PBMC (Nantakhruea et al., 2013), it is interesting to examine miR-29a transcripts in PBMC inoculated with PRRSV. At 24 and 48 h post infection, we found an upregulation of miR-29a in PRRSV-infected PBMC relative to uninfected control. Similar to *in vitro* observation, expression analysis of miR-29a showed higher miR-29a transcript in both LP and HP-PRRSV infected animals. Although both *in vitro* and *in vivo* data did not provide any statistical significance, we still observed similar trend of upregulation of miR-29a in both studies. In addition, our results revealed higher expression of miR-29a in pigs infected with LP-

animals. At 10 days post challenged, PBMCs were isolated from pigs. Although there is no PRRSV than pigs infected with HP-PRRSV. Therefore, it is intriguing to note that different patterns of




Figure 4 MiR-29a expression in PRRSV-infected PBMCs. PBMCs were incubated with PRRSV for 24 and 48 h. Expression of miR-29a was determined by qRT-PCR and normalized to GAPDH gene. The relative fold change of miR-29a in PRRSV-infected cells was shown as the relative to control




Figure 5 Expression of miR-29a in PBMCs isolated from PRRSV-infected pigs (Fig 5a). Relative fold change of miR-29a was normalized to β -actin gene (Fig 5b). LP-PRRS : low pathogenic PRRS virus infection, HP-PRRS : highly pathogenic PRRS virus infection, control : non-infected PBMC. The β -actin gene was used as an internal control. M : 10 bp marker, N : no template control, P : positive control (pGEM-miR29a)

miR-29a might lead to different immunological outcome between diverse strains of PRRSV infection (Han et al., 2013; Zhang et al., 2013). In conclusion, our results revealed that porcine miR-29a was highly conserved with miR-29a from other organisms and miR-29a transcripts were expressed in normal porcine cells and tissues. Moreover, expression analysis of miR-29a suggested that miR-29a increased in PRRSV-infected PBMCs both *in vitro* and *in vivo*. A more detailed analysis of target gene expression should be examined to better understand the role of miR-29 during PRRSV infection. These data provide valuable information on PRRS virus: host interaction and could be applied for PRRSV control in the future.

Acknowledgements

This project was partially funded by the Kasetsart University Research and Development Institute, the Thailand Research Fund (Grant no. MRG5580180), the National Research Council of Thailand, and Graduate Study Research Scholarship for International Publication, the Graduate School of Kasetsart University. The authors would like to thank Professor Roongroje Thanawongnuwech (Faculty of Veterinary Science, Chulalongkorn University), Dr. Susichol Sitthinan and Betagro hybrid farm, Nakorn Prathom for providing pig tissues and blood samples.

References

- Bartel DP 2009. MicroRNAs: Target recognition and regulatory functions. *Cell*. 136(2): 215-233.
- Brain O, Owens Benjamin MJ, Pichulik T, Allan P, Khatamzas E, Leslie A, Steevens T, Sharma S, Mayer A, Catuneanu Ana M, Morton V, Sun MY, Jewell D, Coccia M, Harrison O, Maloy K, Schönfeldt S, Bornschein S, Liston A and Simmons A 2013. The intracellular sensor NOD2 induces microRNA-29 expression in human dendritic cells to limit IL-23 release. *Immunity*. 39(3): 521-536.
- Chen CZ, Schaffert S, Fragoso R and Loh C 2013. Regulation of immune responses and tolerance: the microRNA perspective. *Immunol Rev*. 253(1): 112-128.
- Chen C, Deng B, Qiao M, Zheng R, Chai J, Ding Y, Peng J and Jiang S 2012. Solexa sequencing identification of conserved and novel microRNAs in backfat of large white and Chinese Meishan pigs. *PLoS ONE*. 7(2): e31426.
- Díaz I, Darwich L, Pappaterra G, Pujols J and Mateu E 2005. Immune responses of pigs after experimental infection with a European strain of porcine reproductive and respiratory syndrome virus. *J Gen Virol*. 86(7): 1943-1951.
- Dotti S, Guadagnini G, Salvini F, Razzuoli E, Ferrari M, Alborali GL and Amadori M 2013. Time-course of antibody and cell-mediated immune responses to Porcine Reproductive and Respiratory Syndrome virus under field conditions. *Res Vet Sci*. 94(3): 510-517.
- Frossard JP, Hughes GJ, Westcott DG, Naidu B, Williamson S, Woodger NGA, Steinbach F and Drew TW 2013. Porcine reproductive and respiratory syndrome virus: Genetic diversity of recent British isolates. *Vet Microbiol*. 162(2-4): 507-518.
- Han Z, Liu Y, Wang G, He Y, Hu S, Li Y, Shi W, Wu J, Wang S, Liu H and Cai X 2013. Comparative analysis of immune responses in pigs to high and low pathogenic porcine reproductive and respiratory syndrome viruses isolated in China. *Transbound Emerg Dis*. 6(10): 12190.
- Krol J, Loedige I and Filipowicz W 2010. The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet*. 11(9): 597-610.
- Labarque GG, Nauwynck HJ, Van Reeth K and Pensaert MB 2000. Effect of cellular changes and onset of humoral immunity on the replication of porcine reproductive and respiratory syndrome virus in the lungs of pigs. *J Gen Virol*. 81(5): 1327-1334.
- Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W and Tuschl T 2002. Identification of tissue-specific microRNAs from mouse. *Curr Biol*. 12(9): 735-739.
- Liang Y, Ridzon D, Wong L and Chen C 2007. Characterization of microRNA expression profiles in normal human tissues. *BMC Genomics*. 8(1): 166.
- Liston A, Papadopoulou A, Danso-Abeam D and Dooley J 2012. MicroRNA-29 in the adaptive immune system: setting the threshold. *Cell Mol Life Sci*. 69(21): 3533-3541.
- Livak KJ and Schmittgen TD 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods*. 25(4): 402-408.
- Ma F, Xu S, Liu X, Zhang Q, Xu X, Liu M, Hua M, Li N, Yao H and Cao X 2011. The microRNA miR-29 controls innate and adaptive immune responses to intracellular bacterial infection by targeting interferon- γ . *Nat Immunol*. 12(9): 861-869.
- Nantakhruea S, Lekcharoensuk P and Surachetpong W 2013. Analysis of miRNA-29a expression in porcine peripheral blood mononuclear cells using quantitative reverse transcription polymerase chain reaction. *Thai J Vet Med*. 43(4): 595-600.
- Neumann E, Kliebenstein J, Johnson C, Mabry J, Bush E, Seitzinger A, Green A and Zimmerman J 2005. Assessment of the economic impact of porcine reproductive and respiratory syndrome on swine production in the United States. *J Am Vet Med Assoc*. 227(3): 385-392.
- Podolska A, Anthon C, Bak M, Tommerup N, Skovgaard K, Heegaard P, Gorodkin J, Cirera S and Fredholm M 2012. Profiling microRNAs in lung tissue from pigs infected with *Actinobacillus pleuropneumoniae*. *BMC Genomics*. 13(1): 459.
- Savan R, Legiewicz M, McFarland A, Schwerk J, Bindewald E, Orr S, Ramakrishnan K, Yalamanchili R, Kronfli A, McVicar D, Carrington M, Anderson S, Shapiro B, LeGrice S and Young H 2011. MicroRNA-29 stabilizes interferon-gamma mRNA by antagonizing AU-rich element-mediated decay. *J Immunol*. 186(57): 53.
- Steiner DF, Thomas MF, Hu JK, Yang Z, Babiarz JE, Allen CDC, Matloubian M, Billech R and Ansel KM 2011. MicroRNA-29 regulates T-Box transcription factors and interferon- γ production in helper T cells. *Immunity*. 35(2): 169-181.

- Sun N, Liu D, Chen H, Liu X, Meng F, Zhang X, Chen H, Xie S, Li X and Wu Z 2012^a. Localization, expression change in PRRSV infection and association analysis of the porcine TAP1 gene. *Int J Biol Sci.* 8(1): 10.
- Sun Y, Han M, Kim C, Calvert JG and Yoo D 2012^b. Interplay between interferon-mediated innate immunity and porcine reproductive and respiratory syndrome virus. *Viruses.* 4(4): 424-446.
- Wills RW, Doster AR, Galeota JA, Sur JH and Osorio FA 2003. Duration of infection and proportion of pigs persistently infected with porcine reproductive and respiratory syndrome virus. *J Clin Microbiol.* 41(1): 58-62.
- Wongyanin P, Buranapraditkul S, Yoo D, Thanawongnuwech R, Roth JA and Suradhat S 2012. Role of porcine reproductive and respiratory syndrome virus nucleocapsid protein in induction of interleukin-10 and regulatory T-lymphocytes (Treg). *J Gen Virol.* 93(Pt 6): 1236-1246.
- Xie SS, Li XY, Liu T, Cao JH, Zhong Q and Zhao SH 2011. Discovery of porcine microRNAs in multiple tissues by a Solexa deep sequencing approach. *PLoS ONE.* 6(1): e16235.
- Zhang H, Guo X, Nelson E, Christopher-Hennings J and Wang X 2012. Porcine reproductive and respiratory syndrome virus activates the transcription of interferon alpha/beta (IFN- α/β) in monocyte-derived dendritic cells (Mo-DC). *Vet Microbiol.* 159(3-4): 494-498.
- Zhang L, Liu J, Bai J, Wang X, Li Y and Jiang P 2013. Comparative expression of Toll-like receptors and inflammatory cytokines in pigs infected with different virulent porcine reproductive and respiratory syndrome virus isolates. *Virol J.* 10(1): 135.
- Zuker M 2003. Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 31(13): 3406-3415.

บทคัดย่อ

ลักษณะทางโมเลกุลและระดับการแสดงออกของ miR-29a ในเซลล์สุกร และเซลล์เม็ดเลือดขาวที่ติดเชื้อไวรัสพาร์อาร์เอส

วิน สุรเชษฐพงษ์* สุนารี นันตะเครือ พรทิพภา เล็กเจริญสุข

ไมโครอาร์เอ็นเอเป็นอาร์เอ็นเอขนาดเล็กที่ควบคุมการแสดงออกของยีนผ่านการทำลายเอ็มอาร์เอ็นเอหรือยับยั้งการสร้างโปรตีน วัตถุประสงค์ของงานวิจัยนี้เพื่อศึกษาลักษณะทางโมเลกุลและระดับการแสดงออกของ miR-29a ในเนื้อเยื่อสุกร การศึกษาพบว่าลำดับเบสของ miR-29a สุกรมีความเหมือนในระดับที่สูงกับ miR-29a ในสิ่งมีชีวิตอื่น นอกจากนี้การวิเคราะห์ด้วยโปรแกรมคอมพิวเตอร์พบว่าลำดับเบสบริเวณ seed region ของ miR-29a สุกรสามารถจับได้บางส่วนกับอินเตอร์เฟอรอนแกมมาเอ็มอาร์เอ็นเอ การศึกษาด้วยเทคนิคอาร์ทีพีซีอาร์พบว่า miR-29a มีการแสดงออกในเนื้อเยื่อ ต่อม้ำเหลือง ม้าม เม็ดเลือดขาว และเซลล์ PK-15 การแสดงออกของ miR-29a ในเซลล์เม็ดเลือดขาวที่ติดเชื้อไวรัสพาร์อาร์เอสมีปริมาณเพิ่มขึ้นที่ 24 และ 48 ชั่วโมง การแสดงออกยังเพิ่มขึ้นในสุกรที่ติดเชื้อไวรัสพาร์อาร์เอสทั้งสายพันธุ์ไม่รุนแรงและสายพันธุ์รุนแรง จากผลการทดลองทั้งหมดแสดงให้เห็นว่า miR-29a สุกรมีความคล้ายคลึงกับ miR-29a ของสิ่งมีชีวิตชนิดอื่น และระดับการแสดงออกของ miR-29a มีปริมาณเพิ่มขึ้นในเซลล์เม็ดเลือดขาวที่ติดเชื้อพาร์อาร์เอสในหลอดทดลองและในสุกรมีชีวิต การศึกษาเพิ่มเติมเกี่ยวกับหน้าที่ของ miR-29a จะช่วยอธิบายปฏิสัมพันธ์ระหว่างไวรัสและโฮสต์ ซึ่งอาจนำไปสู่การพัฒนาวิธีป้องกันโรคพาร์อาร์เอสในอนาคต

คำสำคัญ: ภูมิภาคัมกัน miR-29a สุกร ไวรัสพาร์อาร์เอส

ภาควิชาจุลชีววิทยาและวิทยาภูมิคุ้มกัน คณะสัตวแพทยศาสตร์ มหาวิทยาลัยเกษตรศาสตร์ จตุจักร กรุงเทพฯ 10900

* ผู้รับผิดชอบบทความ E-mail: fvetwsp@ku.ac.th