

6-1-2015

Effects of Vitamin D3 and Calcium on Productive Performance, Egg quality and Vitamin D3 Content in Egg of Second Production Cycle Hens

Hatairat Plaimast

Suwanna Kijparkorn

Punyaphat Ittitanawong

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Plaimast, Hatairat; Kijparkorn, Suwanna; and Ittitanawong, Punyaphat (2015) "Effects of Vitamin D3 and Calcium on Productive Performance, Egg quality and Vitamin D3 Content in Egg of Second Production Cycle Hens," *The Thai Journal of Veterinary Medicine*: Vol. 45: Iss. 2, Article 4.

Available at: <https://digital.car.chula.ac.th/tjvm/vol45/iss2/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Effects of Vitamin D₃ and Calcium on Productive Performance, Egg quality and Vitamin D₃ Content in Egg of Second Production Cycle Hens

Hatairat Plaimast* Suwanna Kijparkorn Punyaphat Ittitanawong

Abstract

This experiment was conducted to investigate the effect of supplementary vitamin D₃ on productive performance, internal egg quality, eggshell quality, apparent calcium digestibility and vitamin D₃ content in egg of second production cycle hens fed on different levels of calcium. A total of 140 second cycle brown-egg laying hens, aged 98 weeks, were randomly divided into 5 treatments with 4 replicates. The treatments consisted of diets containing 3.5%Ca and 2,000 IU of vitamin D₃/kg of feed (control), 3.5%Ca and 6,000 IU of vitamin D₃/kg of feed (treatment 2), 3.75%Ca and 2,000 IU of vitamin D₃/kg of feed (treatment 3), 3.75%Ca and 6,000 IU of vitamin D₃/kg of feed (treatment 4), and 4.0%Ca and 2,000 IU of vitamin D₃/kg of feed (treatment 5). All diets were fed for 8 weeks. Thereafter, apparent calcium digestibility was studied for 5 days. There was no significant difference in productive performance and internal egg quality among the dietary treatments ($p>0.05$). Thickness of eggshell increased with the addition of 6,000 IU vitamin D₃ in the diet containing 3.5%Ca or with the supplementation of calcium higher than 3.5% in the diet ($p<0.05$). The hens fed on 4.0%Ca in the diet produced eggs with greater specific gravity than the 3.5%Ca groups ($p<0.05$). Apparent calcium digestibility was not affected by the dietary treatments ($p>0.05$). Vitamin D₃ content in egg yolks increased with the content of vitamin D₃ in the diet ($p<0.01$). In conclusion, second production cycle brown-egg laying hens require dietary calcium higher than 3.5% or the addition of 6,000 IU vitamin D₃ in diet containing 3.5%Ca for eggshell quality improvement. Vitamin D₃ content in egg yolks can be increased at least 2.5 times by adding 6000 IU of vitamin D₃/kg feed to hens feed.

Keywords: calcium, eggshell quality, hens, performance, vitamin D₃

Department of Animal Husbandry, Faculty of Veterinary Science, Chulalongkorn University, Bangkok 10330, Thailand

**Correspondence: hatairatplmt@yahoo.com*

Introduction

Many nutritional factors are involved in eggshell quality of laying hens. Among those factors, calcium and vitamin D₃ appear to be the most relevant and important. Calcium plays an important role in eggshell formation and maintenance of eggshell quality. The shell of egg is composed primarily of calcium carbonate; Ca makes up about 38% of the eggshell (Leeson and Summers, 2001; McDowell, 2003). Functions of vitamin D₃ are associated with calcium. Vitamin D₃ is essential for intestinal uptake and subsequent calcium metabolism. The active form of vitamin D₃ promotes the synthesis of Ca-binding protein which is involved in the calcium transportation in the gut and probably in the uterus. The requirement for vitamin D₃ is increased if the supply of calcium is not at an optimum level (Bar et al., 1992; Leeson and Summers, 2001; Sreenivasaiiah, 2006; Bar et al., 2008). In aged hens, eggshell quality is reduced due to disorders related to calcium and vitamin D metabolism. Eggshell deposition is not enough to prevent a decline in eggshell quality when egg size is larger. A large number of eggs with poor shell quality are produced from older hens (Roland, 1980; Abe et al., 1982; Albatshan et al., 1994; Bar et al., 2002). This problem can be ameliorated by forced molting. During the molt the reproductive system is allowed a complete rest from laying and the bird builds up its body reserves of nutrients. There are evidences that layers improve their calcium and Vitamin D metabolism before starting a second laying cycle (Albatshan et al., 1994; Bar et al., 2001; Sreenivasaiiah, 2006; Nascimento et al., 2014). However, the beneficial effect of calcium and Vitamin D on eggshell quality does not remain after a long period. In a study, Arpasova et al. (2010) indicated that some eggshell qualitative parameters and dietary calcium utilization for shell formation decreased while the number of cycles increased. Therefore, supplementing calcium and vitamin D₃ to a diet might be a good strategy to maintain eggshell quality of second cycle laying hens. Limited research information is available on calcium and vitamin D₃ requirements of repeat cycle brown-egg laying hens. Zapata and Gernat (1995) observed that increasing the level of Ca from 3.0 to 3.5% significantly increased specific gravity of egg produced by molted white leghorn hens from 1.073 to 1.075. Cufadar et al. (2011) suggested that molted brown laying hens should be fed 3.6%Ca in the diet to maintain eggshell quality. Recently, Nascimento et al. (2014) have recommended that the Ca level for second cycle white egg layers should be approximately 4%. In older brown egg layers, Safaa et al. (2008) reported that more than 3.5%Ca in the diet was required to maintain egg production and eggshell quality. Vitamin D for poultry must be in the form of vitamin D₃ (NRC, 1994). Some studies reported that shell quality increased when vitamin D₃ or its metabolites were supplemented in laying hen diets (Frost and Roland, 1990; Tsang and Grunder, 1993). However, excessive vitamin D₃ can produce adverse effects. The maximum tolerance of poultry to vitamin D₃ is 4 to 10 times of the minimum requirement (NRC, 1994). Therefore, the vitamin D₃ content used in this study was 3 times of the CP brown layer nutrient recommendation and this level was not

toxic for hens (Mattila et al., 2004). In addition, many researches showed that vitamin D content in egg could be increased by supplementing laying hen diets with vitamin D (Mattila et al., 1999; Mattila et al., 2003; Yao et al., 2013; Browning and Cowieson, 2014). In the present study, the objective was to evaluate the effect of the additional vitamin D₃ on productive performance, internal egg quality, eggshell quality, apparent calcium digestibility and vitamin D₃ content in egg of second cycle laying hens fed various levels of calcium.

Materials and Methods

Animals and experimental diets: All experimental procedures used in this experiment were approved by the Animal Ethics Committee of Chulalongkorn University. One hundred and forty 98-week-old commercial brown-egg laying hens (CP Brown) in their second production cycle were divided into five treatments. Twenty-eight birds were assigned to each treatment consisting of four replicates. The treatments consisted of diets containing 3.5%Ca and 2,000 IU of vitamin D₃/kg of feed (control), 3.5%Ca and 6,000 IU of vitamin D₃/kg of feed (treatment 2), 3.75%Ca and 2,000 IU of vitamin D₃/kg of feed (treatment 3), 3.75%Ca and 6,000 IU of vitamin D₃/kg of feed (treatment 4), and 4.0%Ca and 2,000 IU of vitamin D₃/kg of feed (treatment 5). A control diet was formulated to meet the calcium and vitamin D₃ requirements of CP brown layer. Vitamin D₃ was supplemented in two lower calcium level treatments (treatments 2 and 4). The basal or control diet was based on corn and soybean meal. The ingredients and calculated nutrient composition of the basal diet are shown in Table 1. All experimental diets were fed for 8 weeks. At the end of the 8th week, chromic oxide (0.25%) was added to the diets to conduct a calcium digestibility study. These diets were fed for 5 days. All experimental diets were prepared and presented in a mash form. The diet and water were provided ad libitum throughout the experimental period. The experimental diets were fed twice a day at 08.00 h and 13.00 h. All hens were reared in an evaporative cooling system with a lighting program, 16 h per day. Temperature in the hen house was recorded daily during the experimental period.

Data collection and chemical analyses: The number of total eggs were recorded and weighed daily by replicate. Feed intake was measured weekly on a replicate. Egg production, average egg weight and average daily feed intake were calculated from these data. At the end of feeding period, sixteen eggs from each treatment were randomly collected to measure internal egg quality, eggshell quality and vitamin D₃ content. The internal egg quality was determined by Haugh unit and percentage of albumen and yolk. The Haugh unit measurement was performed by using a standard tripod micrometer which measured the albumen height in millimeters. This device was adjusted manually for egg weight and displayed HU directly. The albumen and yolk were separated and weighed. Then, the percentage of albumen and yolk were calculated. The eggshell quality was determined

by shell thickness and specific gravity of egg. The shell thickness was measured at three different parts (aircell, equator and sharp end) by micrometer. Egg specific gravity was determined by using NaCl solutions made of incremental concentration of 0.004 in the range from 1.070 to 1.110. The vitamin D₃ content was determined in freeze-dried egg yolk using the method of the Association of German Agricultural Analytic and

Research Institutes (VDLUFA, 2007). For the calcium digestibility study, fecal content of each replicate was collected in approximately the same amount at the same time of the day, during the last 3 days. Chromic oxide of diets and feces were measured by the method described by Fenton and Fenton (1979). Dry matter and calcium content of diets and feces were determined by the methods of AOAC (2005).

Table 1 Ingredient composition and calculated nutrient composition of basal diet

Ingredient	Amount (kg/100kg)
Corn	54.66
Soybean meal	22.55
Rice bran	12.00
Limestone	8.28
Monocalciumphosphate	1.33
Lard	0.33
Sodiumbicarbonate	0.24
Salt	0.18
Choline Chloride	0.11
DL-Methionine	0.11
Vitamin Premix*	0.10
Mineral Premix**	0.10
Antioxidant	0.01
Calculated nutrient content	
Crude Protein (%)	16.30
ME (Kcal/kg)	2750
Crude Fat	4.35
Crude Fiber	3.70
Calcium	3.50
Available Phosphorus	0.40
Vitamin D ₃ (IU/kg)	2000

*Vitamin premix per kilogram of diet provided 10,000 IU vitamin A; 2,000 IU vitamin D₃; 20 IU vitamin E; 2 mg vitamin K₃; 2 mg vitamin B₁; 5 mg vitamin B₂; 5 mg vitamin B₆; 10 mg vitamin B₁₂; 30 mg niacin; 10 mg panthothenic acid; and 0.74 mg folic acid.

**Mineral premix per kilogram of diet provided 25 mg zinc, 10 mg copper, 120 mg iron, 60 mg manganese, and 0.2 mg selenium.

Calculation and statistical analysis: The apparent calcium digestibility was calculated using the following formula:

$$\text{Apparent Ca digestibility} = [1 - (\% \text{ Chromic oxide in diet} \times \% \text{ calcium in feces}) / (\% \text{ Chromic oxide in feces} \times \% \text{ calcium in diet})] \times 100$$

All dependent variables were performed as a completely randomized design to determine the effect of treatment groups. The dependent variables, production performance egg quality and calcium digestibility were analyzed by using a one-way analysis of variance. Difference between treatments was determined by Duncan's New Multiple Range test at the level of $p < 0.05$.

Results

The effects of supplementary vitamin D₃ on the productive performance and internal egg quality of second production cycle laying hens fed different levels of calcium are shown in Table 2. The dietary treatments had no effect ($p > 0.05$) on the amount of feed

consumed, egg weight and egg production. The average of feed intake, egg weight and egg production of hens from all treatments ranged from 110.9 to 111.8 g/d, 63.4 to 65.9 g and 72.8 to 74.6%, respectively. The Haugh units and percentage of albumen and yolk were not influenced by the dietary treatments ($p > 0.05$).

The eggshell quality and apparent calcium digestibility are shown in Table 3. The eggshell quality parameters were affected by the dietary calcium levels or vitamin D₃ supplementation. The eggshell thickness was lowest in eggs from hens fed the control diet ($p < 0.05$). The thickness of eggshell increased with the addition of 6,000 IU vitamin D₃ containing 3.5%Ca or with the increase in calcium higher than 3.5% in the diet. The hens fed 4.0%Ca in the diet produced eggs with greater specific gravity than the 3.5%Ca groups ($p < 0.05$). There was no significant difference in the apparent calcium digestibility among the treatment groups ($p > 0.05$).

As depicted in figure 1, the vitamin D₃ content in egg yolks increased with the content of vitamin D₃ in the diet ($p < 0.01$). There was no significant difference in the vitamin D₃ content of egg yolk among the treatments with the same level of vitamin D₃ in the diet.

The hens fed on the higher vitamin D₃ level (6,000 IU/kg feed) produced higher vitamin D₃ content in egg yolks (3360-3527 IU/kg), while the hens fed on the lower vitamin D₃ level (2,000 IU/kg feed) produced lower vitamin D₃ content in egg yolks (1324-1495 IU/kg).

Discussion

Increasing the dietary calcium levels and vitamin D₃ supplementation did not affect the feed intake, egg weight and egg production of the laying hens in the current study. Several published researches also reported that no additional improvement in productive performance or internal egg quality were obtained when dietary calcium was increased to more than 3.5% (Keshavarz et al., 1993; Roland and Bryant, 1994; Castillo et al., 2004). In addition, Bar et al. (2002) found that egg production, egg weight and feed intake were not affected when the calcium content of the diet was increased from 3.6 to very high level of 4.9% in Lohmann Brown hens aged 66 to 78 weeks. However, Safaa et al. (2008) indicated that an increase in calcium level from 3.5 to 4% of the diet improved some performance parameters, including egg production,

egg mass, and feed conversion ratio in Lohmann Brown laying hens aged 58 to 73 weeks. The reason for the disparity among researchers might be due to the differences in age, number of cycle, rate of egg production and nutrient content of the diet used. For vitamin D₃ supplementation, it did not affect the productive performance and internal egg quality of the hens receiving 3.5 and 3.75% of calcium in the diet. This finding agrees with the report of Mattila et al. (2004), who found that vitamin D₃ supplements (6,000 IU/kg feed) had no effects on production traits compared with the control diet (2,500 IU/kg feed) for groups of laying hens (20 to 67 weeks of age) receiving 3.69% of calcium in the diet. Browning and Cowieson (2014) also reported no difference in layer performance or egg quality with higher levels of vitamin D₃ (up to 10,000 IU/kg feed). From this study, it is revealed that increasing calcium levels and supplementing vitamin D₃ cannot improve animal productive performance and internal egg quality. This is probably because 3.5% of calcium and 2,000 IU/kg feed of vitamin D₃ in the diet are adequate for maintaining productive performance and internal egg quality of second production cycle laying hens.

Table 2 Productive performance and internal egg quality as influenced by dietary Ca and vitamin D₃ in second production cycle hens

Item	Ca 3.5%	Ca 3.5% + Vit D ₃	Ca 3.75%	Ca 3.75% + Vit D ₃	Ca 4%	SEM	p-value
<u>Performance</u>							
Egg production (%)	73.21	73.66	74.11	72.77	74.55	2.316	0.825
Egg weight (g)	64.40	64.21	64.95	65.89	63.43	2.884	0.805
Feed intake (g/d)	110.9	111.8	111.1	111.4	110.9	1.779	0.951
<u>Internal egg quality</u>							
Haugh unit	79.84	81.13	80.44	82.85	83.42	3.382	0.525
Percentage of albumen (%)	64.42	62.08	61.19	61.66	60.06	2.928	0.349
Percentage of yolk (%)	25.53	23.75	24.95	24.56	24.07	1.3542	0.397

Table 3 Eggshell quality and apparent calcium digestibility as influenced by dietary Ca and vitamin D₃ in second production cycle hens

Item	Ca 3.5%	Ca 3.5% + Vit D ₃	Ca 3.75%	Ca 3.75% + Vit D ₃	Ca 4%	SEM	p-value
<u>Eggshell quality</u>							
Eggshell thickness (mm)	0.335 ^a	0.370 ^b	0.361 ^b	0.373 ^b	0.379 ^b	0.031	0.011
Egg specific gravity	1.092 ^a	1.093 ^a	1.095 ^{ab}	1.097 ^{ab}	1.099 ^b	0.006	0.047
Apparent calcium digestibility (%)	51.25	40.69	51.28	43.46	48.44	8.667	0.348

^{a,b} means in a row with different superscripts are significantly different ($p < 0.05$).

The improvement in eggshell quality was observed with the increase in level of calcium or supplementation of vitamin D₃ content in the diet. The eggshell thickness was improved by adding 6,000 IU vitamin D₃ in the diet containing 3.5%Ca or supplementing calcium higher than 3.5% in the diet. This finding indicates that the supplementation of vitamin D₃ into diet has the ability to compensate for the adverse effect of low calcium level (3.5% in feed) on eggshell quality. More vitamin D₃ content might

improve the eggshell quality by increasing the active form of vitamin D₃ (1,25-dihydroxycholecalciferol; 1,25(OH)₂D₃) production in the kidney. 1,25(OH)₂D₃ stimulated the synthesis of calcium-binding protein which is essential for transportation of calcium across the intestinal membrane and may be essential for transportation of calcium for eggshell formation in the shell gland (Bar et al., 1992; Yoshimura et al., 1997; Bar et al., 2008). However, the thickness of eggshell was not changed by supplementing vitamin D₃ content in the

group of hen fed on the higher level of calcium (3.75%Ca). Some authors also observed that egg specific gravity and shell thickness were not significantly different between diets containing 1,500-2,500 and 6,000 IU vitamin D₃/kg feed when laying hens were fed on 3.69-3.87%Ca (Yannakopoulos and Morris, 1979; Mattila et al., 2004). However, the group receiving 4.0%Ca in the diet had a significantly greater specific gravity than those receiving 3.5%Ca. This finding is supported by the report of Safaa et al. (2008), who observed improvement in shell thickness (from 0.342 to 0.351 mm) and shell density (82.0 to 83.8 mg/cm²) of eggs in Lohmann Brown laying hens aged 58 to 73 weeks when the calcium level of the diet was increased from 3.5 to 4%. The results also agree with

the report of Lichovnikova (2007), who suggested 4.1% of calcium in the diet for maintaining eggshell quality in the last period of the production cycle. The result of calcium digestibility study may support this finding. The second production cycle laying hens fed on different diets had similar apparent calcium digestibility while the amount of feed consumed did not differ among the treatment groups. Therefore, the total amount of calcium absorbed from higher calcium diets should be greater than the lower calcium diets with similar feed intake. The better eggshell quality could be obtained from the higher amount of absorbable calcium. These might be the reason for the higher egg specific gravity when the calcium content of the diet was increased to 4%.




Figure 1 Vitamin D₃ content (IU/kg) in egg yolk as influenced by dietary Ca and vitamin D₃ in second production cycle hens

Vitamin D is found naturally in only a few food sources of human, and eggs are considered to be one of the most important sources of vitamin D. In present study, the vitamin D₃ content in egg yolks could be increased by supplementing hens feed with vitamin D₃. The addition of 6000 IU of vitamin D₃/kg feed to laying hen diets resulted in vitamin D₃-enriched eggs which contained more than 2.5 times of the amount of vitamin D₃ found in those produced from the control group. The capacity of vitamin D deposition in egg yolk was similar to previous studies. Mattila et al. (2004) reported that eggs produced from hens receiving 6000 IU of vitamin D₃/kg feed contained 2.7 to 3.6 times more vitamin D than eggs from hens receiving 2500 IU of vitamin D₃/kg feed. A similar result was also reported by Browning and Cowieson (2014), who observed that the vitamin D content of egg yolk increased (1.6 times) with the addition of vitamin D₃ (5000 IU/kg feed) compared with the lower concentration of vitamin D₃ (2500 IU/kg feed) in the layer feed. It is evident that eggs with higher vitamin D content can be produced by feeding hens with increasing level of vitamin D₃.

In conclusion, during the second production cycle commercial brown-egg strain of laying hens require dietary calcium higher than 3.5% or the addition of 6,000 IU vitamin D₃ in diet containing

3.5%Ca for eggshell quality improvement. Vitamin D₃ content in egg can be increased (>2.5 times) by adding 6000 IU of vitamin D₃/kg feed to hen feed.

Acknowledgements

This research was supported by the Chulalongkorn University-Veterinary Science Research Fund. The authors are grateful to BASF (Thai) limited for Vitamin D₃ analysis in yolk.

References

- Abe E, Horikawa H, Masumura T, Sugahara M, Kubota M and Suda T 1982. Disorders of cholecalciferol metabolism in old egg-laying hens. *J Nutr.* 112: 436-446.
- Albatshan HA, Scheideler SE, Black BL, Garlich JD and Anderson KE 1994. Duodenal calcium uptake, femur ash and eggshell quality decline with age and increase following molt. *Poult Sci.* 73: 1590-1596.
- AOAC International. 2005. Official Methods of Analysis of AOAC International. 18th ed. Gaithersburg: AOAC International.
- Arpasova H, Halaj M and Halaj P 2010. Eggshell quality and calcium utilization in feed of hens in

- repeated laying cycles. *Czech J Anim Sci.* 55(2): 66-74.
- Bar A, Vax E and Striem S 1992. Relationships between calbindin (Mr 28,000) and calcium transport by the eggshell gland. *Comp Biochem Physiol B.* 101: 845-848.
- Bar A, Razaphkovsky V, Wax E and Malkat Y 2001. Effect of age at molting on post molting performance. *Poult Sci.* 80: 874-878.
- Bar A, Razaphkovsky V and Vax E 2002. Re-evaluation of calcium and phosphorus requirements in aged laying hens. *Br Poult Sci.* 43:261-169.
- Bar A 2008. Calcium homeostasis and vitamin D metabolism and expression in strongly calcifying laying birds *Comp Biochem Physiol A.* 151(4): 477-490.
- Bolin DW, King RP and Klosterman EW 1952. A simplified method for the determination of chromic oxide (Cr₂O₃) when used as an index substance. *Sci.* 116: 634-635.
- Browning LC and Cowieson AJ 2014. Vitamin D fortification of eggs for human health. *J Sci Food Agric.* 94:1389-1396.
- Castillo C, Cuca M, Pro A, Gonzalez M and Morales E 2004. Biological and economic optimum level of calcium in White Leghorn laying hens. *Poult Sci.* 83(6): 868-72
- Cufadar Dr Y, Olguna O and Yildiza AO 2011. The effect of dietary calcium concentration and particle size on performance, eggshell quality, bone mechanical properties and tibia mineral contents in moulted laying hens. *Br Poult Sci.* 52(6): 761-768.
- Fenton TW and Fenton M 1979. Determination of chromic oxide in feed and feces. *Canadian J. Anim. Sci.*, 58: 631-634.
- Frost TJ and Roland DA Sr 1990. Influence of vitamin D₃, 1 α -hydroxyvitamin D₃, and 1-25-dihydroxyvitamin D₃ on eggshell quality, tibia strength, and various production parameters in commercial laying hens. *Poult Sci.* 69: 2008-2016.
- Keshavarz K and Nakajima S 1993. Re-evaluation of calcium and phosphorus requirements of laying hens for optimum performance and eggshell quality. *Poult Sci.* 72: 144-153.
- Leeson S and Summers JD 1997. *Commercial poultry nutrition.* 2nd ed. Guelph: University books 350pp.
- Leeson S and Summers JD 2001. *Nutrition of the Chicken.* 4th ed. Guelph: University books 591pp.
- Lichovnikova M 2007. The effect of dietary calcium source, concentration, and particle size on calcium retention, eggshell quality, and overall calcium requirements in laying hens. *Br Poult Sci.* 48: 71-75.
- Mattila P, Lehikoinen K, Kiiskinen T and Piironen V 1999. Cholecalciferol and 25-hydroxycholecalciferol content of chicken egg yolk as affected by the cholecalciferol content of feed. *J Agric Food Chem.* 47:4089-4092.
- Mattila P, Rokka T, Konko K, Valaja J, Rossow L and Ryhanen EL 2003. Effect of cholecalciferol-enriched hen feed on egg quality. *J Agric Food Chem.* 51(1): 283-287.
- Mattila, P, J Valaja, L Rossow, E Vanalainen and T Tupasela 2004. Effect of vitamin D₂ and D₃ enriched diets on egg vitamin D content, production, and bird condition during an entire production period. *Poult Sci.* 83: 433-440.
- McDowell LR 2003. *Minerals in animal and human nutrition.* 2nd ed. Amsterdam: Elsevier science 644pp.
- Nascimento GR do, Murakami AE, Guerra AFQM, Ospinas-Rojas IC, Ferreira MFZ and Fanhani JC 2014. Effect of Different Vitamin D Sources and Calcium Levels in the Diet of Layers in the Second Laying Cycle. *Braz J Poult Sci.* 16(2): 37-42.
- National Research Council 1994. *Nutrient Requirements of Poultry.* 9th rev. ed. Washington DC: National Academy Press 155pp.
- Roland DA Sr 1980. The ability of young and old hens to change shell deposition with sudden neutral drastic changes in egg size. *Poult Sci.* 59:924-926.
- Roland DA Sr and Bryant MM 1994. Influence of calcium on energy consumption and egg weight of commercial leghorns. *J Appl Poult Res.* 3: 184-189.
- Safaa HM, Serrano MP, Valencia DG, Frikha M, Jimenez-Moreno E and Mateos GG 2008. Productive performance and egg quality of brown egg-laying hens in the late phase of production as influenced by the level and source of calcium in the diet. *Poult Sci.* 87: 2043-2051.
- Sreenivasiah PV 2006. *Scientific poultry production.* A Unique Encyclopaedia. 3rd ed. Lucknow : International Book Distributing Co. 1487pp.
- Tsang CPW and Grunder AA 1993. Effect of dietary contents of cholecalciferol, 1 α ,25-dihydroxycholecalciferol and 24,25-dihydroxycholecalciferol on blood concentrations of 25-hydroxycholecalciferol, 1 α ,25-dihydroxycholecalciferol, total calcium and eggshell quality. *Br Poult Sci.* 34: 1021-1027.
- Verband Deutscher Landwirtschaftlicher Untersuchungs und Forschungsanstalten (VDLUFA) (Hrsg.) 2007. *Methode 13.8.1 Determination of Vitamin D₃ by HPLC.* In: *Handbuch der Landwirtschaftlichen Versuchs- und Untersuchungsmethodik (VDLUFA-Methodenbuch), Bd. III Die chemische Untersuchung von Futtermitteln* 3. Aufl. VDLUFA-Verlag: Darmstadt.
- Yannakopoulos A and Morris T 1979. Effect of light, vitamin D and dietary phosphorus on egg-shell quality late in the pullet laying year. *Br Poult Sci.* 20:337-342.
- Yao L, Wang T, Persia M, Horst RL and Higgins M 2013. Effects of vitamin D₃-enriched diet on egg yolk vitamin D₃ content and yolk quality. *J Food Sci.* 78:178-83.
- Yoshimura Y, Ohira H and Tamura T 1997. Immunocytochemical localization of vitamin D receptors in the shell gland of immature, laying, and molting hens. *Gen Comp Endocr.* 108: 282-289.
- Zapata LF and Gernat AG 1995. The effect of four levels of ascorbic acid and two levels of calcium on eggshell quality. *Poult Sci.* 74: 1049-1052.

บทคัดย่อ

ผลของวิตามินดีสามและแคลเซียมต่อสมรรถภาพการผลิต คุณภาพไข่และ ปริมาณวิตามินดีสามในไข่ของไก่ไข่ที่ให้ผลผลิตรอบที่สอง

หทัยรัตน์ พลายมาศ* สุวรรณมา กิจภากรณ์ ปัญพัสตร์ อธิธิธนาวงษ์

การทดลองนี้ศึกษาผลของการเสริมวิตามินดีสามต่อสมรรถภาพการผลิต คุณภาพภายในของไข่ คุณภาพเปลือกไข่ การย่อยได้แบบ ปรากฏของแคลเซียมและปริมาณวิตามินดีสามในไข่ของไก่ไข่ที่ให้ผลผลิตรอบที่สองที่ได้รับปริมาณแคลเซียมในระดับต่างๆ แบ่งไก่ไข่ให้ เปลือกไข่สีน้ำตาลที่ให้ผลผลิตรอบสองจำนวนทั้งหมด 140 ตัว อายุ 98 สัปดาห์ ออกเป็น 5 กลุ่มๆละ 4 ซ้ำ อาหารทดลองประกอบด้วยอาหาร ควบคุมที่มีระดับแคลเซียม 3.5% และวิตามินดีสาม 2,000 IU ต่อ กิโลกรัมของอาหาร อาหารที่มีระดับแคลเซียม 3.5% และวิตามินดีสาม 6,000 IU ต่อ กิโลกรัมของอาหาร อาหารที่มีระดับแคลเซียม 3.75% และวิตามินดีสาม 2,000 IU ต่อ กิโลกรัมของอาหาร อาหารที่มีระดับ แคลเซียม 3.75% และวิตามินดีสาม 6,000 IU ต่อ กิโลกรัมของอาหาร และอาหารที่มีระดับแคลเซียม 4.0% และวิตามินดีสาม 2,000 IU ต่อ กิโลกรัมของอาหาร การทดลองพบว่าสมรรถภาพการผลิตและคุณภาพภายในของไข่แต่ละกลุ่มทดลองไม่แตกต่างกัน ($p>0.05$) การเสริม วิตามินดีสามในอาหารที่มีระดับของแคลเซียม 3.5% หรือการให้แคลเซียมในระดับที่สูงกว่า 3.5% มีผลทำให้เปลือกไข่มีความหนาเพิ่มขึ้น ($p<0.05$) ไก่ไข่ที่ได้รับอาหารที่มีระดับของแคลเซียม 4.0% ให้ผลผลิตไข่ที่มีความถ่วงจำเพาะสูงกว่ากลุ่มที่ได้รับแคลเซียม 3.5% ($p<0.05$) อาหารทดลองที่แตกต่างกันไม่มีผลต่อการย่อยได้แบบปรากฏของแคลเซียม ($p>0.05$) ปริมาณวิตามินดีสามในไข่แดงเพิ่มขึ้นตามปริมาณ วิตามินดีสามที่เพิ่มขึ้นในอาหาร ($p>0.01$) จากผลการทดลองสรุปได้ว่าไก่ไข่ให้เปลือกไข่สีน้ำตาลที่ให้ผลผลิตรอบสองต้องการแคลเซียมใน อาหารสูงกว่า 3.5% หรือวิตามินดีสามเสริมในปริมาณ 6,000 IU ต่อ กิโลกรัมของอาหารที่มีระดับของแคลเซียม 3.5% เพื่อปรับปรุงคุณภาพ ของเปลือกไข่ ปริมาณวิตามินดีสามในไข่แดงเพิ่มขึ้นอย่างน้อย 2.5 เท่าเมื่อเสริมวิตามินดีสามในอาหารไก่ไข่ 6,000 IU ต่อ กิโลกรัมของอาหาร

คำสำคัญ: แคลเซียม คุณภาพเปลือกไข่ ไก่ไข่ สมรรถภาพการผลิต วิตามินดีสาม

ภาควิชาสัตวบาล คณะสัตวแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ปทุมวัน กรุงเทพฯ 10330

*ผู้รับผิดชอบบทความ E-mail: hatairatplmt@yahoo.com