

4-1-1990

Acquired secondary syphilis in 10-year-old boy

Pomarong Chotiwan

Pirom Kamolratanakul

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Chotiwan, Pomarong and Kamolratanakul, Pirom (1990) "Acquired secondary syphilis in 10-year-old boy," *Chulalongkorn Medical Journal*: Vol. 34: Iss. 4, Article 8.

Available at: <https://digital.car.chula.ac.th/clmjournal/vol34/iss4/8>

This Case Report is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Acquired secondary syphilis in 10-year-old boy

Pornarong Chotiwan*
Pirom Kamolratanakul*

Chotiwan P, Kamolratanakul P. Acquired secondary syphilis in 10-year-old boy. Chula Med J 1990 Apr; 34(4) : 307-310

A 10-year-old Thai boy attended out-patient clinic because of patchy alopecia. Acquired secondary syphilis was diagnosed after complete investigation and responded to benzathine penicillin injections. This is the first proved acquired secondary syphilis in a child reported in Thailand.

Reprint request : Chotiwan P, Department of Preventive and Social Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. April 19, 1990.

พรณรงค์ โชติวรรณ, ภิรมย์ กมลรัตนกุล. ซิฟิลิสในเด็ก 10 ขวบ. จุฬาลงกรณ์เวชสาร 2533 เมษายน ; 34(4) : 307-310

รายงานผู้ป่วยเด็กชายไทยอายุ 10 ขวบ 1 ราย มาโรงพยาบาลด้วยอาการผื่นร่วงเป็นหย่อม ๆ ตรวจสอบพบว่ามีอาการติ่งเนื้อซิฟิลิส มีหลักฐานแสดงว่าน่าจะเป็นการติดเชื้อภายหลังคลอด แต่ไม่ทราบวิธีการติดเชื้อ ผู้ป่วยหายดีหลังจากให้ยาเพนนิซิลินชนิดฉีด ยังไม่มีรายงานผู้ป่วยวัยนี้มาก่อนในประเทศไทย

Acquired syphilis in children is relatively rare when compared with congenital syphilis. There has been no report of acquired syphilis in children in Thailand. Most of the reports were about congenital syphilis.⁽¹⁻⁸⁾ There are few reported cases of acquired syphilis in prepubertal children. Most cases are caused by sexual abuse. Ginsbury 1983⁽⁹⁾ reported 3 cases in 9 to 11 years old children. White et al 1983⁽¹⁰⁾ reported 6 cases of syphilis from 409 under 12-year-old cases of suspected sexual abuse. Goldenring 1989⁽¹¹⁾ reported a 10-year-old child with secondary syphilis presenting with condyloma lata. Van Niekerk et al 1985⁽¹²⁾ reported that 2% of 1575 primary school children in South Africa got positive test for syphilis. We report a case of proved acquired secondary syphilis in 10-year-old boy presenting with patchy alopecia.

Case Report

A 10-year-old Thai boy came to out-patient

department because of patchy alopecia for more than a month. He was delivered by a traditional midwife together with his twin brother. His mother did not get any antenatal care. The boy was small for date at birth but there was no evidence of congenital syphilis. The parents separated when the child was 3 years old. The boy is in primary school and stays with his mother. Four months before, he ran away from home and came back a month later on. There was no definite history of child sexual abuse. Patchy hair falling began three months after he returned. The boy has 5 elder sisters. There was no stillbirth or abortion in his family.

The boy weighed 18 kg. and was 121.5 cm. high which is much below Thai standard.⁽¹³⁾ The chest circumference was 48.5 cm., and the body temperature was 37.2 degree celsius. He had poor mouth hygiene and no Hutchinson's teeth. The hair showed typical patchy alopecia (Fig. 1). Liver and spleen were not palpable. There was no joint swelling. Normal eyes and eye-ground was noted by an ophthalmologist.





Figure 1. Shows typical patchy alopecia, presenting sign of secondary syphilis in this case.

VDRL gave reactive result but the serum was not enough for titre. TPHA and FTA - ABS was reactive but FTA - ABS (IgM) was non-reactive with normal x-ray survey of long bones and normal findings from lumbar puncture. The serum from the boy's mother and his twin brother were non-reactive for VDRL, TPHA, FTA - ABS and FTA - ABS (IgM).

Treatment with Benzathine penicillin 1.2 million units was given weekly for 3 weeks. The hair returned to normal appearance after 3 months of treatment. VDRL remain reactive at low titre of 1:2 and TPHA also reactive at 2,4, and 6 months follow up. The patient loss follow up after that.

Discussion

Signs and symptoms of secondary syphilis in a child may be the result of either congenital or acquired syphilis. Congenital syphilis is much more common especially in a child under 2 years old. Typical evidence of congenital syphilis such as saddle nose, Hutchinson's teeth, saber-shin, tibial osteitis, Clutton's joints can be used together with history and serum examination to differentiate acquired from congenital syphilis in children.

In this case, the evidence pointed to acquired secondary syphilis because there was no evidence of infection in his mother and twin brother, and no sign of congenital syphilis. Normal eye and lumbar puncture confirmed that there was no late congenital syphilis. Physicians should be alert to the possibility that patchy alopecia in children may be a manifestation of secondary syphilis and consider doing VDRL for secondary syphilis work up.

References

1. Jutamas Sudasna. Congenital syphilis. เชียงใหม่เวชสาร 1984 มิถุนายน 23(2) ; 123-31
2. เสมอ มิ่งวานิช. Syphilis in the newborns and their mothers. วารสารโรงพยาบาลสระบุรี 1983 มิถุนายน-กันยายน ; 8(3) : 1-15
3. เสมอ มิ่งวานิช. Congenital syphilis : report of a case. วารสารโรงพยาบาลสระบุรี 1983 ตุลาคม-ธันวาคม ; 8(4) : 15-23
4. Yupin Thanasophon. Diagnosis of neonatal congenital syphilis. วารสารสมาคมกุมารแพทย์แห่งประเทศไทย 1981 มีนาคม ; 20 : 37-45
5. Yupin Thanasophon. Congenital syphilis. วารสารกรมการแพทย์ 1977 ตุลาคม-ธันวาคม ; 2(4) : 188-99
6. Chorfa Kaewjinda, Dusdee Prabhasawat. Congenital syphilis. สารศิริราช 1973 พฤศจิกายน ; 25(11) : 1988-91
7. Vitoon Osathanondh, Tongdee Ruangpiroj, Oonjai Veuwsorn. A study on current perinatal deaths due to congenital syphilis. J Med Assoc Thai 1966 Nov; 49(11) : 899-926
8. Peythai Mansuwan. Congenital syphilis. วารสารสมาคมกุมารแพทย์แห่งประเทศไทย 1963 มีนาคม ; 2(1) : 44-54
9. Ginsburgh CM. Acquired syphilis in prepubertal children. Pediatr Infect Dis 1983; 2 : 232-4
10. White ST, Loda FA, Ingram DL, Pearson A. Sexually transmitted diseases in sexually abused children. Pediatrics 1983 Jul; 72(1) : 16-21
11. Goldenring JM. Secondary syphilis in a prepubertal child. Differentiating condylomata lata from condylomata acuminata. N Y State J Med 1989 Mar; 89(3) : 180-1
12. van Niekerk CH, van Niekerk LC, Van den Ende J. Positive serological tests for syphilis in black primary school children in Bloemfontein: a pilot study. S Afr Med J 1985 Jan 19; 67(3) : 90-1
13. Chavalittamrong B, Vathakanon R. Standards of height and weight of Bangkok Children. J Med Assoc Thai 1978 Feb; 61 Suppl 2 : 1-28