

3-1-2017

Effects of the higher parity and litter size on piglet birth weight and survival rate of later born piglets

Supatee Kitkha

Alongkot Boonsoongnern

Nattavut Ratanavanichrojn

Pichai Jirawattanapong

Anuchai Pinyopummin

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Kitkha, Supatee; Boonsoongnern, Alongkot; Ratanavanichrojn, Nattavut; Jirawattanapong, Pichai; and Pinyopummin, Anuchai (2017) "Effects of the higher parity and litter size on piglet birth weight and survival rate of later born piglets," *The Thai Journal of Veterinary Medicine*: Vol. 47: Iss. 1, Article 18. Available at: <https://digital.car.chula.ac.th/tjvm/vol47/iss1/18>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Effects of the higher parity and litter size on piglet birth weight and survival rate of later born piglets

Supatee Kitkha^{1,2} Alongkot Boonsoongnern³ Nattavut Ratanavanichrojn³

Pichai Jirawattanapong³ Anuchai Pinyopummin^{3*}

Abstract

The objectives of this study were to evaluate the effects of higher parity and litter size (LS) on piglet birth weight (BW) and to compare the BW and survival rate of piglets born in different birth orders (BO). The study was conducted at a commercial pig farm in Thailand. Data from 230 sows and 3,500 piglets were analyzed including parity, litter size (LS), average birth weight (AvBW), coefficient of variation of piglets' BW (%CV), standard deviation of piglets' BW (SD), percentage of small piglets (%Small), individual piglet's BW, piglet birth order (BO; 1-10, 11-16, 17-25), sex and characteristics at birth (mummified fetuses (MM), stillborn piglets (SB) and born alive piglets (BAL)). Results revealed that the reduction in AvBW was prominent when LS increased ($p < 0.001$) and the expansion of BW variability (%CV, SD, %Small) was significant in the higher parity sows with large LS ($p < 0.001$). A difference in piglet BW was found among different BO; the later born piglets in BO 17-25 were lighter than the piglets born in BO 1-10 and 11-16 ($p < 0.001$). The percentage of BAL steadily decreased concomitantly with the rising of BO ($p < 0.001$), whereas the born dead piglets including MM and SB increased significantly ($p < 0.001$). This study demonstrated that higher parity and LS had negative effects on piglet BW including the reduction in mean BW with a greater variability. Moreover, the later born piglets in large litters had a lower survival rate.

Keywords: birth order, birth weight, litter size, parity, piglet, survival rate

¹Center for Agricultural Biotechnology, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

²Center of Excellence on Agricultural Biotechnology: (AG-BIO/PERDO-CHE), Bangkok 10900, Thailand

³Faculty of Veterinary Medicine, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

*Correspondence: foetacp@ku.ac.th

Introduction

Over the last decade the improvement in sow prolificacy and the ability to increase piglets per litter or litter size (LS) of middle-aged or old sows (Milligan et al., 2002) have assisted the swine industry in confronting piglet losses during peri- and post-natal periods, due to lower mean birth weight (BW) with greater variability (Quiniou et al., 2002) and prolongation of the expulsive stage (van Dijk et al., 2005).

Increased LS causes a greater proportion of small piglets with inadequate body energy, resulting in higher sensitivity to cold. Besides, they have lower ability to compete with their littermates to obtain sufficient colostrum and milk (Le Dividich, 1999). Therefore, most of them have a higher risk of post-natal mortality during the first week after birth (Quiniou et al., 2002).

Moreover, the prolonged expulsive stage in large litters raises the risk of death in later born piglets with lighter weight peri-partum, due to reduced stimuli production, which is necessary for maintenance of uterine contraction (Fraser, 1984), resulting in asphyxia (Herpin et al., 1996).

However, the severity of piglet losses varies among countries and farms (Furtado et al., 2012; Kilbride et al., 2012). In Thailand, the information on piglet losses is scarce. Therefore, the objectives of this study were to evaluate the effects of parity and LS on piglet BW and compare piglet BW and survival rate among different birth orders (BO), of a commercial pig farm in Thailand.

Materials and Methods

Animals: Animal manipulations were approved by the Animal Usage and Ethics Committee, Kasetsart University (ACKU 03555). The study was conducted from June 2012 to August 2013 in Chiang Mai, Thailand (18°47'25" N, 98°59'04" E) at a commercial sow breeding farm housing 5,000 sows with an evaporative cooling system which maintained the temperature at 25°C and relative humidity below 80%. The external temperature and relative humidity varied from 11.6°C to 39.7°C and 48% to 88% (Meteorological Department, 2015), respectively.

All the sows received a gestation and lactation diet that contained 15.5% crude protein (CP), 3,048 Kcal/kg metabolizable energy (ME) and 0.9% total lysine (Lys) during weeks 1-11 of gestation; 16.4% CP, 3,048 Kcal/kg ME and 0.9% Lys during weeks 12-15 of gestation; and 18.1% CP, 3,314 Kcal/kg ME and 1.2% Lys. During lactation and prior to conception, the sows were fed ad libitum four times a day. After mating, they were fed twice a day and the feed level was controlled depending on the sow's body condition score (BCS) (BCS < 3, 2.0 kg/day; BCS = 3, 1.5 kg/day; BCS > 3, 1.0 kg/day). Water was supplied through a nipple drinker and was available at all times.

Estrus monitoring after weaning was carried out at 08.00 and 15.00 by trained farm technicians using fence-line boar contact. Estrus signs were detected when the sows exhibited a standing heat reflex during a back pressure test in the presence of a boar.

Traditional artificial insemination (AI) with pooled semen was done at 12 and 24 h after estrus expression.

Gestating sows were moved to a farrowing barn one week before the expected day of parturition. The farrowing pen was composed of a farrowing crate on a slatted floor under the sow and a metal slatted floor under piglets. A heat lamp was placed inside a metal box which was beside the sow to provide heat for the piglets. At one day of age, all piglets had their teeth clipped and tail docked and were injected with iron.

Data collection: Cross-bred (Large white x Landrace) sows (n = 230) were recorded for parity (1, 2, 3, 4, 5, 6, ≥ 7) and LS. After birth, piglets (n = 3,500) were individually weighed. Their individual BW, BO, sex and characteristics (mummified fetuses (MM), stillborn piglets (SB) and born alive piglets (BAL)) were also recorded. Then, the piglets' data were calculated for average birth weight (AvBW), coefficient of variation of BW (%CV), standard deviation of BW (SD) and percentage of small piglets (weighing less than one kilogram within the litter) (%Small piglets). The data collection from both sows and their litters was conducted by questionnaires.

Statistical analysis: Factors (LS and parity) that affected AvBW and BW variation (%CV, SD and %Small piglets) were identified by linear regression (Winter, 2013). The model after this procedure was $Y_{ijk} = L_i + P_j + E_{ijk}$, where Y_{ijk} = response variable (AvBW, %CV, SD and %Small piglets), L_i = litter size, P_j = parity, and E_{ijk} = residual error.

Comparison of piglet BW among different BO (1-10, 11-16, 17-25) with correction of the effect of sex (male and female) (Čechová, 2006) was determined by the two-way analysis of covariance (ANCOVA), in which LS was accounted a covariate (Quiniou et al., 2002). Significance of the ANCOVA results was further analyzed by post hoc analyses using Bonferroni (SPSS version 18, SPSS Inc., Chicago, IL, USA).

Proportions of piglets' characteristic in different BO (1-10, 11-16, 17-25) were tested by the chi-square test and Tukey-type multiple comparison (Zar, 1996).

A significant level was considered when the *p*-value was < 0.05. Data were presented as mean ± SEM.

Results

Effects of increased parity and LS on piglet BW: The increase in LS decreased the mean BW as described by the following equation: $AvBW = 2.015 - 0.033 LS$, $R^2 = 0.251$, $p < 0.001$. For every one piglet added to a litter, the AvBW would be reduced by 0.033 kg (Fig. 1). There was no parity effect on AvBW ($p > 0.05$) (Fig. 2). However, the variability of piglet BW depended on both parity and LS (Figs. 1-4). The higher parity and larger LS triggered the rising of SD, %CV and %Small piglet within a litter as described by the three equations as follows: $SD = 0.141 + 0.012 Parity + 0.009 LS$, $R^2 = 0.151$, $p < 0.001$; $\%CV = 2.492 + 0.958 Parity + 1.067 LS$, $R^2 = 0.245$, $p < 0.001$; and $\%Small\ piglet = -10.778 + 0.873 Parity + 1.037 LS$, $R^2 = 0.194$, $p < 0.001$.

Comparison of piglet BW among different BO: After adjusting for LS ($p < 0.001$) and correcting the effect of sex ($p = 0.003$), the main effect of BO on individual piglet BW was significant ($p < 0.001$). There was no interaction (BO \times sex) effect ($p = 0.683$). The piglets born in BO 17-25 (1.40 ± 0.01 kg) were lighter than the piglets born in BO 1-10 (1.51 ± 0.01 kg) and BO 11-16 (1.46 ± 0.01 kg) ($p < 0.05$). However, there was no

difference in piglet BW between BO 1-10 and BO 11-16 ($p \geq 0.05$) (Fig. 5).

Different survival rates among different BO: The percentage of BAL decreased with the increase in BO (94.34%, 84.70%, 71.09%, respectively) ($p < 0.05$), while the percentage of SB (3.06%, 10.88%, 18.58%) and MM (2.61%, 4.42%, 10.32%) grew consecutively ($p < 0.05$) (Fig. 6).




Figure 1 Piglets average birth weight (AvBW) and standard deviation of birth weight (SD) categorized by litter size (vertical error bars = \pm SEM)




Figure 2 Piglets average birth weight (AvBW) and standard deviation of birth weight (SD) categorized by parity (vertical error bars = \pm SEM)




Figure 3 Coefficient of variation of piglets' birth weight (%CV) and percentage of small piglets (%Small piglet) categorized by litter size (vertical error bars = \pm SEM)




Figure 4 Coefficient of variation of piglets' birth weight (%CV) and percentage of small piglets (%Small piglet) categorized by parity (vertical error bars = \pm SEM)




Figure 5 Piglets birth weight in different groups of birth order
 a, b = Significant differences at $p < 0.05$




Figure 6 Percentage of mummified fetuses (MM), stillborn piglets (SB) and born alive piglets (BAL) in different birth orders
 a, b, c = Significant differences at $p < 0.05$

Discussion

Negative effects of higher parity and LS on piglet BW:

The reduction in mean BW was mainly affected by the increase in LS in the present study, whereas parity and LS contributed to the higher piglet BW variability. The results of this study agreed with those of Quiniou et al. (2002), who reported that the increase in LS was inappropriate for the rising of litter BW. The increase

in LS from 9 to 17 piglets was an 88% increase, whereas the total litter BW increased only by 50% (21 vs 14 kg), or the increase in LS from ≤ 11 to ≥ 16 decreased mean BW from 1.59 to 1.26 kg. Besides, SD of BW rose from 0.26 to 0.30 kg and the percentage of low BW piglets (weighing less than one kilogram) rose to 11%. Increased LS is likely related to a higher ovulation rate and embryo survival of middle-aged or older sows (Wrathall, 1971; Milligan et al., 2002). Therefore,

uterine crowding is a consequent problem for these sows, especially after day 35 of pregnancy, when the uterine capacity is limited for a larger number of porcine fetal development (Knight et al., 1977). Reducing uterine blood flow per fetus (Père and Etienne, 2000) and insufficient nutrient supply might be the cause of low BW in large litters (Caugant and Guéblez, 1993). Moreover, the variation of fetal development might depend on the position of implantation (Perry and Rowell, 1969) because the rates of blood flow and nutritional supply vary greatly along the length of the uterus, due to difference in vasculature structure and density (Ford et al., 2002; Wu et al., 2010), resulting in a variation of piglet weight at birth.

Observation of lighter BW in later born piglets: The piglets born in the last group of BO were lighter than the piglets born earlier. This finding is normally observed in large litters (Fraser, 1984). Homogenized embryos are distributed within the uterine horn before day 35 of pregnancy (Anderson and Parker, 1976). Thereafter, porcine fetal development depends on the position and number of fetuses in the uterus (Perry and Rowell, 1969). There is no difference in advantage between positions when there are only five or fewer fetuses in the horn (Perry and Rowell, 1969). However, in large litters, the fetal weight declines from the ovarian end of the uterus to the middle, related to the slight increase in weight of the fetuses near the cervical end (Perry and Rowell, 1969). During the farrowing process, which does not completely occur at random (Friend and Cunningham, 1966), piglets rarely overtake neighboring piglets in the same horn (Taverne et al., 1977), as a result later born piglets tend to be those positioned at the ovary end of the uterus (Friend and Cunningham, 1966) which are slightly lighter than the piglets positioned near the cervical end.

Reduction in survival rate of later born piglets caused by higher BO: The percentage of BAL significantly decreased concomitantly with the higher BO, whereas the percentage of MM and SB steadily increased. The number of piglets born positively correlates with the percentage of piglets born dead (Milligan et al., 2002; Motsi et al., 2006), due to the prolonged duration of the expulsive stage, which affects the degree of asphyxia during delivery (Herpin et al., 1996). Significantly, later born piglets located high up at the ovary end of the uterus tend to suffer from occlusion or rupture of the umbilical cord or detachment of the placenta, leading to inadequate blood flow and oxygen supply to unborn piglets (Fraser, 1984; van der Lende and van Rens, 2003). Therefore, 70% of SB are normally noticed during the last third of the farrowing period (Friend and Cunningham, 1966). In addition, the lighter BW of the last born piglets (Friend and Cunningham, 1966) might lead to a greater risk of death during the farrowing period due to the lower production of stimuli to trigger or maintain uterine contraction during fatigue (Fraser, 1984).

For the already dead piglets or MM, they could not move to reach the right position and enter the pelvic canal to be delivered. This was the reason

why the higher percentage of MM was noticeable in the higher BO. Besides, BO of MM might depend on their size. Large MM are generally expelled at the same time as their littermates, while the majority of small MM are expelled with the placenta (van der Lende and van Rens, 2003).

Conclusion

The increase in parity and LS decreased the mean piglet BW, raised the variability of BW within a litter and suppressed the opportunity of survival of the later and lighter born piglets. Therefore, intensive management should be employed for those sows and piglets. For example, better supervision during delivery could reduce the number of SB and the mortality of live-born piglets (Holyoake et al., 1995), colostrum supplementation at least 200 g per piglet during the first 24 h after birth enhances piglet survival rate (Quesnel et al., 2012) and cross-fostering of small piglets to small litters (Deen and Bilkei, 2004) with similar size littermates (English, 1998) encourages the survival and weight gain until weaning (Robert and Martineau, 2001; Bierhals et al., 2012).

Acknowledgements

The authors gratefully thank Center for Agricultural Biotechnology, Kasetsart University, Center of Excellence on Agricultural Biotechnology: (AG-BIO/PERDO-CHE), Thailand for their financial support; V.P.F. group and its employees for their assistance in this study; and Faculty of Veterinary Medicine, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom, Thailand for the facilities.

References

- Anderson, L.L., Parker, R.O., 1976. Distribution and development of embryos in the pig. *J Reprod Fertil* 46(2): 363-368.
- Bierhals, T., Magnabosco, D., Ribeiro, R.R., Perin, J., da Cruz, R.A., Bernardi, M.L., Wentz, I., Bortolozzo, F.P., 2012. Influence of pig weight classification at cross-fostering on the performance of the primiparous sow and the adopted litter. *Livest Sci* 146(2-3): 115-122.
- Caugant, A., Guéblez, R., 1993. Influence du poids du porcelet à la naissance sur les performances ultérieures. *J Rech Porcine France* 25: 123-128.
- Čechová, M., 2006. Analysis of some factors influencing the birth weight of piglets. *Slovak J Anim Sci* 39(3): 139-144.
- Deen, M.G.H., Bilkei, G., 2004. Cross fostering of low-birthweight piglets. *Livest Prod Sci* 90(2-3): 279-284.
- English, P.R., 1998. Ten basic principles of fostering piglets. *Pig Prog* 14, 39-41.
- Ford, S.P., Vonnahme, K.A., Wilson, M.E., 2002. Uterine capacity in the pig reflects a combination of uterine environment and conceptus genotype effects. *J Anim Sci* 80(E.Suppl.1): E66-E73.
- Fraser, D., 1984. The role of behavior in swine production: a review of research. *Appl Anim Behav* 11(4): 317-339.

- Friend, D.W., Cunningham, H.M., 1966. Piglet birthweights and the order of farrowing. *Can J Comp Med Vet Sci* 30(7): 179-182.
- Furtado, C.S.D., G., M.A.P., Cypriano, C.R., Gaggini, T.S., Bernardi, M.L., Wentz, I., Bortolozzo, F.P., 2012. Influence of birth weight and of oral, umbilical or limb lesions on performance of suckling piglets. *Acta Sci Vet* 40(4): 1077.
- Herpin, P., Le Dividich, J., Hulin, J.-C., Fillaut, M., DeMarco, F., Bertin, R., 1996. Effect of level of asphyxia during delivery on viability at birth and early post-natal viability of new born pig. *J Anim Sci* 74(9): 2062-2076.
- Holyoake, P.K., Dial, G.D., Trigg, T., King, V., 1995. Reducing pig mortality through supervision during the perinatal period. *J Anim Sci* 73(12): 3543-3551.
- Kilbride, A.J., Mendl, M., Statham, P., Held, S., Harris, M., Cooper, S., Green, L.E., 2012. A cohort study of preweaning piglets mortality and farrowing accommodation on 112 commercial pig farm in England. *Prev Vet Med* 104(3-4): 281-291.
- Knight, J.W., Bazer, F.W., Thatcher, W.W., Frank, D.E., Wallace, H.D., 1977. Conceptus development in intact and unilaterally hysterectomized-ovariectomized gilts: Interrelationships among hormonal status, placental development, fetal fluids, and fetal growth. *J Anim Sci* 44, 620-637.
- Le Dividich, J., 1999. Review: management to reduce variation in pre- and post-weaning pigs. In: Cranwell, P.D. (Ed.), *Manipulating Pig Production VII*. Australian Pig Science Association, pp. 135-155.
- Meteorological Department, 2015. "Statistics of rainfall at meteorology station, Chiang mai province: 2003 - 2015." [Online]. Available: <http://service.nso.go.th/nso/web/statseries/statseries27.html>.
- Milligan, B.N., Fraser, D., Kramer, D.L., 2002. Within-litter birth weight variation in the domestic pig and its relation to pre-weaning survival, weight gain, and variation in weaning weights. *Livest Prod Sci* 76(1-2): 181-191.
- Motsi, P., Sakuhuni, C., Halimani, T.E., Bhebhe, E., Ndiweni, P.N.B., Chimonyo, M., 2006. Influence of parity, birth order, litter size and birth weight on duration of farrowing and birth intervals in commercial exotic sows in Zimbabwe. *Anim Sci* 82(4): 569-574.
- Père, M.C., Etienne, M., 2000. Uterine blood flow in sows: effects of pregnancy stage and litter size. *Reprod Nutr Dev* 40(4): 369-382.
- Perry, J.S., Rowell, J.G., 1969. Variations in foetal weight and vascular supply along the uterine horn of the pig. *J Reprod Fertil* 19(3): 527-534.
- Quesnel, H., Farmer, C., Devillers, N., 2012. Colostrum intake: Influence on piglet performance and factors of variation. *Livest Sci* 146(2-3): 105-114.
- Quinioua, N., Dagorna, J., Gaudre, D., 2002. Variation of piglets' birth weight and consequences on subsequent performance. *Livest Prod Sci* 78(1): 63-70.
- Robert, S., Martineau, G.P., 2001. Effects of repeated cross-fostering on preweaning behavior and growth performance of piglets and on maternal behavior of sows. *J Anim Sci* 79(1): 88-93.
- Taverne, M.A.M., Weyden, v.d., Fontijne, P., Ellendorff, F., Naakteboren, C., Smidt, D., 1977. Uterine position and presentation of minipig-fetuses and their order and presentation at birth. *Am J Vet Res* 38(11): 1761-1764.
- van der Lende, T., van Rens, B.T.T.M., 2003. Critical periods for foetal mortality in gilts identified by analysing the length distribution of mummified foetuses and frequency of non-fresh stillborn piglets. *Anim Reprod Sci* 75(1-2): 141-150.
- van Dijk, A.J., van Rens, B.T.T.M., van der Lende, T., Taverne, M.A.M., 2005. Factors affecting duration of the expulsive stage of parturition and piglet birth intervals in sows with uncomplicated, spontaneous farrowings. *Theriogenology* 64(7): 1573-1590.
- Winter, B., 2013. "Linear models and linear mixed effects models in R with linguistic applications." [Online]. Available: <https://arxiv.org/ftp/arxiv/papers/1308/1308.5499.pdf>.
- Wrathall, A.E., 1971. Prenatal survival in pigs. Part I. Ovulation rate and its influence on prenatal survival and litter size in pigs. Commonwealth Agriculture Bureau, Slough, UK.
- Wu, G., Bazer, F.W., Burghardt, R.C., Johnson, G.A., Kim, S.W., Li, X.L., Satterfield, M.C., Spencer, T.E., 2010. Impacts of amino acid nutrition on pregnancy outcome in pigs: Mechanisms and implications for swine production. *J Anim Sci* 88(113 Suppl.): E195-E204.
- Zar, J.H., 1996. *Biostatistical analysis*. Prentice-Hall, Inc. Silmon & Schuster/A Viacom Company, New Jersey, USA.

บทคัดย่อ

ผลของลำดับท้องและขนาดครอกที่เพิ่มขึ้นต่อน้ำหนักแรกเกิดของลูกสุกร และอัตราการรอดชีวิตของลูกสุกรที่คลอดที่หลัง

สุพาทิ กิจคำ^{1,2} อลงกต บุญสูงเนิน³ ณัฐวุฒิ รัตนวณิชโรจน์³ พิชัย จิรวัดนาพงศ์³ อนุชัย ภิญโญภูมิมินทร์^{3*}

วัตถุประสงค์ของการศึกษาเพื่อศึกษาผลของลำดับท้องและขนาดครอกที่เพิ่มขึ้นต่อน้ำหนักแรกเกิดของลูกสุกร และเปรียบเทียบ น้ำหนักแรกเกิดและอัตราการรอดชีวิตของลูกสุกรที่คลอดในลำดับการคลอดที่ต่างกัน ในฟาร์มสุกรเอกชนแห่งหนึ่งในประเทศไทย ทำการ เก็บข้อมูลจากแม่สุกรจำนวน 230 ตัวและลูกสุกรจำนวน 3,500 ตัว ดังนี้ ลำดับท้อง ขนาดครอก น้ำหนักแรกเกิดเฉลี่ยภายในครอก ค่าสัมประสิทธิ์ความผันแปรของน้ำหนักแรกเกิด ค่าส่วนเบี่ยงเบนมาตรฐานของน้ำหนักแรกเกิด ร้อยละของลูกสุกรตัวเล็ก น้ำหนักแรกเกิดของ ลูกสุกรรายตัว กลุ่มลำดับการคลอด (ตัวที่ 1-10, 11-16 และ 17-25) เพศ และลักษณะของลูกสุกร (ลูกกรอก ตายคลอด และคลอดมีชีวิต) ผล การศึกษาแสดงให้เห็นว่า น้ำหนักแรกเกิดน้อยลง ($p < 0.001$) เมื่อขนาดครอกใหญ่ขึ้น และความผันแปรของน้ำหนักแรกเกิดเพิ่มขึ้น ($p < 0.001$) เมื่อแม่สุกรมีลำดับท้องมากขึ้นร่วมกับการมีครอกขนาดใหญ่ น้ำหนักแรกเกิดของลูกสุกรมีความแตกต่างกันระหว่างกลุ่มของลำดับการ คลอด โดยลูกสุกรที่คลอดในกลุ่ม 17-25 มีน้ำหนักแรกเกิดน้อยกว่า ($p < 0.001$) ลูกสุกรที่คลอดในกลุ่ม 1-10 และ 11-16 ร้อยละของลูกสุกร คลอดมีชีวิตรอดลง ($p < 0.001$) เมื่อลำดับการคลอดเพิ่มขึ้น โดยมีการเพิ่มขึ้น ($p < 0.001$) ของลูกสุกรที่เป็นลูกกรอกและตายคลอด การศึกษา นี้แสดงให้เห็นว่าลำดับท้องที่มากขึ้นและขนาดครอกใหญ่มีผลต่อน้ำหนักแรกเกิดของลูกสุกร ทั้งในลักษณะของการมีน้ำหนักแรกเกิดลดลง และการมีความผันแปรของน้ำหนักแรกเกิดเพิ่มขึ้น นอกจากนี้ ร้อยละของลูกสุกรที่คลอดมีชีวิตในกลุ่มลำดับการคลอดที่หลังยังมีจำนวน ลดลง

คำสำคัญ: ขนาดครอก น้ำหนักแรกเกิด ลูกสุกร ลำดับการคลอด ลำดับท้อง อัตราการรอดชีวิต

¹ศูนย์เทคโนโลยีชีวภาพเกษตร มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140

²ศูนย์ความเป็นเลิศด้านเทคโนโลยีชีวภาพเกษตร สำนักพัฒนาบัณฑิตศึกษาและวิจัยด้านวิทยาศาสตร์และเทคโนโลยี สำนักงานคณะกรรมการ การอุดมศึกษา กรุงเทพฯ 10900

³คณะสัตวแพทยศาสตร์ มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140

*ผู้รับผิดชอบบทความ E-mail: fvetacp@ku.ac.th