

3-1-2018

Crossover feeding test of a new milk mixture for giant panda (*Ailuropoda melanoleuca*) cubs

Zhihe Zhang

Rong Hou

Jingchao Lan

Hairui Wang

Tatsuko Nakao

See next page for additional authors

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Zhang, Zhihe; Hou, Rong; Lan, Jingchao; Wang, Hairui; Nakao, Tatsuko; Takatsu, Zenta; Kobayashi, Toyokazu; Koie, Hiroshi; Kamata, Hiroshi; Kanayama, Kiichi; and Watanabe, Toshi (2018) "Crossover feeding test of a new milk mixture for giant panda (*Ailuropoda melanoleuca*) cubs," *The Thai Journal of Veterinary Medicine*: Vol. 48: Iss. 1, Article 2.

Available at: <https://digital.car.chula.ac.th/tjvm/vol48/iss1/2>

This Short Communication is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Crossover feeding test of a new milk mixture for giant panda (*Ailuropoda melanoleuca*) cubs

Authors

Zhihe Zhang, Rong Hou, Jingchao Lan, Hairui Wang, Tatsuko Nakao, Zenta Takatsu, Toyokazu Kobayashi, Hiroshi Koie, Hiroshi Kamata, Kiichi Kanayama, and Toshi Watanabe

Crossover feeding test of a new milk mixture for giant panda (*Ailuropoda melanoleuca*) cubs

Zhihe Zhang¹ Rong Hou¹ Jingchao Lan¹ Hairui Wang¹ Tatsuko Nakao² Zenta Takatsu^{3*}

Toyokazu Kobayashi⁴ Hiroshi Koie⁵ Hiroshi Kamata⁵ Kiichi Kanayama⁵ Toshi Watanabe¹

Abstract

A milk substitute developed in 1988 (OM) has been used to feed giant panda cubs, but its contents do not match those of panda breast milk. OM contains 10% more protein than the breast milk, and so a novel milk substitute (NM) was developed based on breast milk data. NM has the following nutrition: protein, 38%; fat, 40%; carbohydrates, 12% (containing lactose, 7%); ash, 6%; and moisture, 3%; and it also contains lactoferrin, nucleotides, oligosaccharides and docosahexaenoic acid. In order to ensure that NM is safe before it is fed to premature newborns, a plan was made to subject twin cubs (Kaihin and Youhin, around one year old) to the first feeding test of NM. Here, a crossover feeding test of OM mixture (OMM) and NM-containing mixture (NMM) is described. Kaihin and Youhin were fed 800-1300 ml/d NMM and OMM, respectively, combined with 200 ml/d breast milk for 280-380 days after birth. Then, they were fed 1370-1570 ml/d OMM and NMM, respectively, without breast milk for the next 100 days. Both cubs exhibited constant growth of 90 g/day throughout the test, which is comparable with the growth of previous panda cubs (including twins). Moreover, no NMM-associated safety concerns were observed. Feces were excreted every 3-5 d during milk feeding, whereas they were excreted several times a day when bamboo was consumed. It could be concluded that NM can be fed safely to newborn panda cubs.

Keywords: feeding, giant panda, cub, milk

¹Sichuan Key Laboratory of Conservation Biology for Endangered Wildlife, Chengdu Research Base of Giant Panda Breeding, 1375 Panda Road, Northern Suburb, Chengdu, Sichuan 610081, P. R. China

²Adventure World, Aws Co., Ltd, 2399, Katada, Shirahama, Nishimuro-gun, Wakayama 649-2201, Japan

³Morinyu Sunworld Co. Ltd., Research & Information Center, Morinaga Milk, 5-1-83, Higashihara, Zama, Kanagawa 252-8583, Japan

⁴Department of Animal Sciences, Faculty of Life & Environmental Sciences, Teikyo University of Science, 2-2-1, Senju-Sakuragi, Adachi-ku, Tokyo 120-0045, Japan

⁵Department of Veterinary Medicine, College of Bioresource Sciences, Nihon University, 1866, Kameino, Fujisawa, Kanagawa 252-8510, Japan

*Correspondence: z_takatu@morinagamilk.co.jp

Introduction

Milk substitutes are used to feed giant panda cubs in captivity. OM, a milk substitute for neonate panda cubs, was released in 1988 (Tanabe, 1995); however, it has a slightly higher protein content than panda breast milk (protein, 48%; fat, 41%; carbohydrates, 4%; ash, 4%; moisture, 3%; and lactose, nil) (Table 1). Other formulas have also been used to feed panda cubs, but their contents differ from those of panda breast milk (Edwards et al., 2006). While these substitutes have been successfully used for nursing, their effects on growth have not been reported in

detail. Zhang (2013) reported that the survival rates of giant panda newborns in captivity around the world at one year old were 71% (20/28) in 2012 and 86% (42/49) in 2013.

At Adventure World, OM has been administered to newborn panda cubs, a mixture of OM, dog milk powder, and human infant formula (OMM), during mid-term feeding for 10 years. Nine cubs grew well with the OMM; and Nakao et al. (2013) reported that the survival rate of panda cubs fed the OMM between 2001 and 2013 was 92% (11/12). Fig. 1 shows growth curves for another set of twins that were born in 2008, Meihin (a female) and Eihin (a male).

Table 1 Composition of the milk substitutes, milk mixtures, and breast milk (DM basis)

The composition of the four milk substitutes used as raw materials, and the mixtures fed to the cubs are shown.

OLD MILK, TD, and NL were mixed at a ratio of 4:5:4 and administered as the control mixture (OMM).

NEW MILK, TD, and NL were mixed at a ratio of 4:5:4 and administered as the test mixture (NMM).

The nutritional contents of the breast milk of Dashuang, YaYa, and Qi Zhen were reported previously (Zhang et al., 2016).

Milk	Protein %	Fat %	Carbohydrates %	Ash %	Moisture %	Energy kcal/100 g	Lactose %
OLD MILK	48	41	4	4	3	572	0
NEW MILK	38	40	12	6	3	561	7
TD	47	28	17	5	3	510	8
NL	13	20	62	2	3	479	0
OLD MILK+TD+NL (4:5:4)	37	30	27	4	3	519	3
NEW MILK+TD+NL (4:5:4)	34	29	29	5	3	516	5
Dashuang (25 days after delivery)	38.7	40.8	14.7	5.8	0.0	580	5.3
YaYa (160 days)	29.0	35.9	32.0	3.2	0.0	567	3.6
QiZhen (154 days)	25.9	50.1	21.5	3.2	0.0	640	1.6




Figure 1 Growth curves of another set of twins reared at Adventure World

The growth curves of another set of twins, Meihin, a female, and Eihin, a male, who were born in 2008 and fed OMM during the same period (FH, d280-380), are shown.

Zhang et al. (2016) proposed NM as a milk substitute for nursing panda cubs that exhibited a closer nutritional profile to panda breast milk than previous milk substitutes. NM powder has the following nutritional profile: protein, 38%; fat, 40%; carbohydrates, 12%; ash, 6%; moisture, 3%; and lactose, 7%. It also contains docosahexaenoic acid/eicosapentaenoic acid, nucleotides, arachidonic acid, taurine, lactoferrin, and L-carnitine. There has been little opportunity to test the feasibility of feeding newborn panda cubs with NM because panda cubs are of global importance. There is similar difficulty in

performing experiments on neonates of other captive animals. A mixture of fish fillets and a milk substitute was fed to *Tursiops* neonates (Sweeney et al., 2010), but the comparison with breast milk was insufficient. In human infants, new ingredients should be fully studied in children and young adults before they are tested on neonates (Aggett et al., 2001; CEAI, 2004). For example, while probiotics are widely used in adults and young people, their use in infants aged from 0-6 months is still being studied (Bertelsen et al., 2016; FAO, 2001).

The other foods made smaller contributions to the pandas' nutritional intake in FH than during LH. The pandas ate 60 g/day (24-168 g) apples from d279 to 380 (Kaihin and Youhin ate 5708 g and 6480 g, respectively). The twins took small amounts of biscuits and carrots from their mother's supplies. In LH, the pandas consumed 120-250 g/day apples after d404, 10-60 g/day biscuits after d433, and 10 g/day carrots after d488. The pandas consumed 3-20 g of bamboo shoots on d304 and d309 in FH, consumed some bamboo with their mother from d404 to 425 in LH, and took 1600~12500 g/day of young bamboo after d426. The maximum total amount of food, including bamboo, consumed was estimated during d495-504 in LH (Table 2) from published composition data (Zhang et al., 2016; CST, 2015; Dierenfeld et al., 1982; Schaller et al., 1985).

Both twins were very healthy and demonstrated reasonable growth throughout the study; Kaihin and Youhin grew by 93 and 108 g/d in FH, and by 93 and 92 g/d in LH, respectively (Fig. 4).

Kaihin was more active, and thus, gained slightly less weight than Youhin. The cubs' growth rates were compared using the t-test. First, the 82 and 79 pieces of growth rate data obtained for Kaihin and Youhin in FH were compared, respectively, and the 44 and 43 pieces of growth rate data collected for the pandas in LH, respectively. Then, the 82 and 44 pieces of growth rate data collected for Kaihin in FH and LH were compared, respectively, and the 79 and 43 pieces of growth rate data obtained for Youhin in FH and LH, respectively. The four sets of growth data did not differ significantly ($t=0.01\sim0.44$, t -critical one tail= $1.65\sim1.66$ with $\alpha 0.05$). The twins' BW values at 12 months of age were comparable with those of healthy cubs, including twins, which had previously been raised on the OMM at Adventure World in the past 10 years. Both milk mixtures allowed the twins to grow appropriately, and no NMM-associated safety concerns arose during the test.




Figure 2 Volume of breast milk fed to the cubs (g/day)

Only data obtained during FH are shown, as data during the exhibition or free nursing periods could not be obtained. The pandas were fed breast milk one to three times a day in FH and LH until d425. M: the male cub, F: the female cub.




Figure 3 Milk mixture consumption

Both pandas were fed similar volumes of milk mixture in FH and identical amounts of milk mixture in LH. M: the male cub, F: the female cub.

The consumption of apples increased the amount of fecal matter produced by the cubs, and some pieces of apple were observed in the pandas' feces. Most of the young bamboo was excreted, but this did not affect the condition of the pandas' digestive tracts. The cubs' bamboo consumption was recorded, but as the degree of moisture evaporation from the bamboo changed daily, the accurate amount taken could not be determined, so the pandas' feces were monitored instead. The total amounts of collected feces are shown in Fig. 5. Kaihin and Youhin produced 1731 g and 3245 g of fecal matter during FH, respectively. In addition, they excreted feces every 5.1 days and 2.6 days, and the mean amount of fecal matter excreted per day was 87 g and 83 g, respectively. The amount of feces produced

by the pandas during FH was very small compared with the amount of milk solids they consumed, i.e. the cubs consumed 167 g/day milk solids, which would have produced 560 g/day of feces if none of the milk had been digested (assuming a fecal moisture content of 70%) (Schaller et al., 1985). Therefore, it was speculated that more than 95% of the milk was absorbed, as described previously (NAFR, 2009). The differences in fecal production in FH between the twins might have been due to variations in apple consumption. In LH, the feces excreted by the two pandas could not be collected separately after d432 because the amount of feces produced increased greatly after the twins started to consume bamboo regularly (Dierenfeld et al., 1982).




Figure 4 Growth curves for the twins

Separate growth curves for FH and LH are shown. The crossover feeding of the two mixtures did not result in any differences in BW gain. M: the male cub, F: the female cub.




Figure 5 Weight of the collected feces

Most feces could be collected during FH, but the feces produced by each of the pandas were difficult to be gathered separately after d433 because the amount of feces produced by the pandas increased; therefore, only the total amount is shown in LH. M: the male cub, F: the female cub.

Kaihin suffered a fever from d321 to 322, and exhibited mouth breathing and coughing after consuming milk or sliced apples from d373 to 377. In addition, he consumed less milk on d374. Youhin excreted soft feces containing milk particles from d316 to 320. However, none of these issues were considered to be related to the milk mixtures. Both cubs grew well during LH, and no health-related issues or abnormal feces were observed in this period.

Growth is an essential endpoint in all safety assessment of new foodstuffs for human infants, and our findings suggest that the NMM meets this criterion. The normal growth demonstrated by the twins indicates that it is safe to conduct a nutritional study of formulas containing NM involving newborn panda cubs. Although it is impossible to carry out large clinical trials, small amounts of data should be accumulated, such as that reported in this study, in

order to build up a clearer picture of how wild panda cubs should be raised in captivity.

Acknowledgements

We thank the staff at Adventure World, especially Teruaki Hayashi, Koji Imazu, and Norikatsu Yasuda, who fed the twins and kept comprehensive records. We are also grateful to Nihon University for facilitating this study and providing financial support. Finally, we thank the giant panda staff at the Chengdu Research Base of Giant Panda Breeding for their guidance and cooperation. This study received grants from the International Science and Technology Cooperation Program of China (2011DFA33530) and China Key Projects in the National Science & Technology Pillar Program (2012BAC01B06).

References

- Aggett PJ, Agostini C, Goulet O, Hernell O, Koletzko B, Lafeber HL, Michaelsen KF, Rigo J and Weaver LT 2001. The nutritional and safety assessment of breast milk substitutes and other dietary products for infants: A commentary by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol.* 32 (3): 256-258.
- Bertelsen RJ, Jensen ET and Ringel-Kulka T 2016. Use of probiotics and prebiotics in infant feeding. *Best Pract Res Clin Gastroenterol.* 30 (1): 39-48.
- CEAI (Committee on the evaluation of the addition of ingredients new to infant formula, food and nutrition board) 2004. Going beyond current clinical studies. In: *Infant Formula Evaluating the Safety of New Ingredients*. Washington DC: The National Academies Press p. 98-159.
- CST (The Council for Science and Technology, Ministry of Education, Culture, Sports, Science and Technology, Japan) 2015. *Standard Tables of Food Composition in Japan*. Tokyo, Japan: Official Gazette Co-operation Japan 589 pp.
- Dierenfeld ES, Hintz HF, Robertson JB, Van SPJ and Oftedal OT 1982. Utilization of bamboo by the giant panda. *J Nutr.* 112 (4): 636-641.
- Edwards MS, Wei R, Hawes J, Sutherland-Smith M, Tang C, Li D, Hu D and Zhang G 2006. The neonatal giant panda: hand-rearing and medical management. In: *Giant Pandas Biology, Veterinary Medicine and Management*. Wildt DE, Zhang A, Zhang H, Janssen DL and Ellis S eds, Cambridge: Cambridge Univ Press p. 315-333.
- FAO (Food and Agriculture Organization of the United Nations) 2001. Report of joint FAO/WHO expert consultation on evaluation of health and nutritional properties of probiotics in food including powder milk with live lactic acid bacteria. p. 1-34.
- NAFR (National Agriculture and Food Research Organization) 2009. *Standard Tables of Feed Composition in Japan*. Tokyo, Japan: Japan Livestock Industry Association p. 287.
- Nakao T, Yasuda N, Imazu K and Hayashi T 2013. The breeding records of giant panda past 19 years (1994-2013) and overview of the new facility in Adventure World. In: *Thesis Collection, International Symposium on Giant Panda Conservation, Chengdu, China: The Chengdu Research Base of Giant Panda Breeding* p. 89-93.
- Schaller GB, Jinchu H, Wenshi P and Jing Z 1985. *The Giant Pandas of Wolong, Chicago, U.S.A.*: University of Chicago Press p. 298.
- Sweeney JC, Stone R, Campbell M, McBrain J, St. Leger J, Xitco M, Jensen E and Ridgway S 2010. Comparative survivability of *Tursiops* neonates from three U.S. institutions for the decades 1990-1999 and 2000-2009.
- Tanabe K 1995. Development of panda milk. In: *Documents of Giant Panda Keeping in Ueno Zoological Garden, 20 Year's Report (in Japanese)*. Tokyo, Japan: Ueno Zoological Garden p. 116-120.
- Xie Z 2013. The 2013 International Studbook for Giant Panda: Chinese Association of Zoological Gardens Beijing, China p. 245.
- Zhang Z 2013. Retrospect and prospects on the giant pandas ex-situ conservation. In: *Thesis Collection, International Symposium on Giant Panda Conservation, Chengdu, China: The Chengdu Research Base of Giant Panda Breeding* p. 1-14.
- Zhang Z, Hou R, Lan J, Wang H, Kurokawa H, Takatsu Z, Kobayashi T, Koie H, Kamata H, Kanayama K and Watanabe T 2016. Analysis of the breast milk of giant pandas (*Ailuropoda melanoleuca*) and the preparation of substitutes. *J Vet Med Sci.* 78(5): 747-754.

บทคัดย่อ

การทดลองครอสโอเวอร์โดยการให้อาหารผสมนมชนิดใหม่แก่ลูกแพนด้ายักษ์ (Ailuropoda melanoleuca)

จางจื้อเหอ¹ โทหวร¹ หลันจิงเซา¹ หวังไห่รู่ย¹ ทัดลีโกะ นากาโอะ² เซ็นตะ ทาคัตสึ^{3*}
โทโยคัตสึ โคบายาชิ⁴ อิโรชิ โคอิเอะ⁵ อิโรชิ คามาตะ⁵ คิอิชิ คานายามะ⁵ โทชิ วาตานาเบะ¹

อาหารทดแทนนม (OM) ที่ได้รับการพัฒนาขึ้นในปี 1988 ถูกนำมาใช้สำหรับเลี้ยงลูกแพนด้ายักษ์ แต่ส่วนประกอบของอาหารดังกล่าวยังไม่สามารถเทียบเท่ากับน้ำนมของแม่แพนด้าได้ OM มีปริมาณโปรตีนมากกว่านมแพนด้า 10% จึงมีการพัฒนาอาหารทดแทนนมตัวใหม่ (NM) ขึ้นมาโดยอ้างอิงจากข้อมูลของนมแพนด้า ซึ่ง NM มีส่วนประกอบของสารอาหาร ได้แก่ โปรตีน 38%, ไขมัน 40%, คาร์โบไฮเดรต 12% (ประกอบด้วยแลคโตส 7%), เกล็ด 6% และ ความชุ่มชื้น 3% นอกจากนี้ ยังมีแลคโตเฟอรัลิน นิวคลีโอไทด์ โอลิโกแซ็กคาไรด์ และกรดโดโคซาเฮกซาอีโนอิกด้วย เพื่อตรวจสอบความปลอดภัยก่อนให้อาหาร NM แก่แพนด้าแรกเกิด จึงได้ทำการทดลองป้อน NM ให้กับลูกแพนด้าฝาแฝด (โคฮินและโยฮิน อายุประมาณ 1 ปี) ก่อน โดยทดลองป้อนอาหารแบบสลับกันระหว่างอาหาร OMM และอาหารที่มีส่วนประกอบของ NM (NMM) ทั้งโคฮินและโยฮินได้รับ NMM และ OMM 800-1,300 มล./วัน ตามลำดับ โดยผสมกับนมแม่ 200 มล./วัน เป็นเวลา 280-380 วันหลังจากที่ลูกแพนด้าทั้งสองเกิด หลังจากนั้นลูกแพนด้าทั้งสองได้รับ OMM และ NMM 1,370-1,570 มล./วัน ตามลำดับ โดยไม่ผสมกับนมแม่ต่ออีกเป็นเวลา 100 วัน การทดลองพบว่า ลูกแพนด้าทั้งสองตัวมีอัตราการเติบโตอยู่ที่ 90 กรัม/วันตลอดระยะเวลาการวิจัย ซึ่งเทียบเท่ากับอัตราการเจริญเติบโตของลูกแพนด้าตัวอื่น ๆ (รวมถึงตัวที่เป็นฝาแฝดด้วย) และไม่พบว่ามีข้อกังวลด้านความปลอดภัยเกี่ยวกับ NMM นอกจากนั้น ยังพบว่าในระหว่างที่กินอาหารทดแทนนมลูกแพนด้ามีการขับถ่ายอุจจาระทุก ๆ 3-5 วัน ในขณะที่การกินต้นไผ่ทำให้ลูกแพนด้าขับถ่ายอุจจาระวันละหลายครั้ง ดังนั้น จึงสรุปได้ว่า NM มีความปลอดภัยสำหรับใช้ป้อนให้กับลูกแพนด้าแรกเกิด

คำสำคัญ: การให้อาหาร แพนด้ายักษ์ ลูกแพนด้า นม

¹ศูนย์วิจัยหลักด้านชีววิทยาและการอนุรักษ์สัตว์ป่าเสฉวน, ฐานวิจัยการขยายพันธุ์แพนด้ายักษ์เฉิงตู, เมืองเฉิงตู, มณฑลเสฉวน, ประเทศสาธารณรัฐประชาชนจีน

²แอดเวนเจอร์ เวิลด์, บริษัท AWS จำกัด, อำเภอชิราฮามะ, จังหวัดวากายามะ, ประเทศญี่ปุ่น

³บริษัท โมรินิว ซันเวิลด์ จำกัด, เขตมินาโตะ, โตเกียว, ประเทศญี่ปุ่น

⁴ภาควิชาสัตวศาสตร์ คณะชีวิตและสิ่งแวดล้อม มหาวิทยาลัยวิทยาศาสตร์เทเคียว, เขตอาดาจิ, โตเกียว, ประเทศญี่ปุ่น

⁵ภาควิชาสัตวแพทยศาสตร์ คณะวิทยาศาสตร์และทรัพยากรชีวภาพ มหาวิทยาลัยนิงงง, อำเภอฟูจิซาวะ, จังหวัดคานากาวะ, ประเทศญี่ปุ่น

*ผู้รับผิดชอบบทความ E-mail: z_takatu@moringamilk.co.jp