

3-1-2018

Mutations in QRDRs of DNA gyrase and topoisomerase IV genes in nalidixic acid and ciprofloxacin-resistant *Salmonella enterica* isolated from chicken meat, pork and humans

Nuananong Sinwat

Jiratchaya Pongseree

Sunpetch Angkittitrakul

Rungtip Chuanchuen

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Sinwat, Nuananong; Pongseree, Jiratchaya; Angkittitrakul, Sunpetch; and Chuanchuen, Rungtip (2018) "Mutations in QRDRs of DNA gyrase and topoisomerase IV genes in nalidixic acid and ciprofloxacin-resistant *Salmonella enterica* isolated from chicken meat, pork and humans," *The Thai Journal of Veterinary Medicine*: Vol. 48: Iss. 1, Article 16.

Available at: <https://digital.car.chula.ac.th/tjvm/vol48/iss1/16>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

**Mutations in QRDRs of DNA gyrase and topoisomerase IV
genes in nalidixic acid and ciprofloxacin-resistant
Salmonella enterica isolated from chicken meat,
pork and humans**

Nuananong Sinwat¹ Jiratchaya Pongseree¹

Sunpetch Angkittitrakul² Rungtip Chuanchuen^{1*}

Abstract

Twenty-eight nalidixic acid-resistant *Salmonella* isolates originated from chicken meat, pork and humans were examined for mutation in the quinolone resistance determining region (QRDR) of *gyrA*, *gyrB*, *parC* and *parE* genes. Four single-point mutations in *gyrA* (i.e. C248A, C248T, G259T and G259A) leading to amino acid substitutions Ser83Tyr, Ser83Phe, Asp87Tyr and Asp87Asn in GyrA, respectively, were identified in 11 quinolone-resistant strains. Amino acid change at position Ser83 was most frequently identified. Resistance to nalidixic acid was not always associated with resistance to ciprofloxacin. The presence of mutations in GyrA did not well correlate with ciprofloxacin resistance phenotype. No mutations were observed in *gyrB* and *parE*. A serovar Typhimurium resistant to nalidixic acid did not carry any mutations in QRDR of all genes tested. The results highlight the high frequency of mutation in *gyrA* in the quinolone-resistant isolates and the existence of alternative-quinolone resistance mechanisms.

Keywords: DNA gyrase, fluoroquinolone resistance, nalidixic acid resistance, *Salmonella enterica*, topoisomerase IV

¹Research Unit in Microbial Food Safety and Antimicrobial Resistance, Department of Veterinary Public Health, Faculty of Veterinary Science, Chulalongkorn University, Bangkok 10330, Thailand

²Faculty of Veterinary Medicine, KhonKaen University, KhonKaen 40000, Thailand

*Correspondence: rchuanchuen@yahoo.com

Introduction

Non typhoidal *Salmonella enterica* infections are a major threat to global public health and is commonly associated with consumption of contaminated food of animal origin (e.g. chicken meat, pork). Salmonellosis is usually self-limited and does not require antibiotic therapy. However, antibiotic treatment may be required for young children or the elderly. Fluoroquinolones have been recommended by WHO as empirical treatment for multidrug resistant enteric fever (WHO, 2007) and are often used to treat invasive *Salmonella* infection in humans (Crement et al., 2011). Particular concern is that the increasing use of fluoroquinolones may lead to the emergence and spread of fluoroquinolone-resistant *Salmonella*. Recently, fluoroquinolone-resistant *Salmonella* has been increasingly reported (Lertworapreecha et al., 2013; Wang et al., 2015).

Quinolones are broad-spectrum antibiotics with four generations, of which nalidixic acid is the first-generation drug exhibiting a narrow-spectrum activity against Gram-negative bacteria. Ciprofloxacin is the second generation quinolone with a broad-spectrum activity against Gram-negative and Gram-positive bacteria (Fabrega et al., 2009). These antibiotics selectively inhibit topoisomerase II (DNA gyrase) and topoisomerase IV and prevent bacterial DNA replication during growth and reproduction. DNA gyrase has two A subunits and two B subunits that are encoded by *gyrA* and *gyrB* genes, respectively. Topoisomerase IV is composed of two C and two E subunits encoded by *parC* and *parE* genes, respectively. During DNA replication, DNA gyrase catalyses negative DNA supercoiling and topoisomerase IV is responsible for segregation daughter chromosome (Hopkins et al., 2005).

Three major mechanisms of quinolone resistance have been recognized including mutations in the quinolone resistance determining region (QRDR) of DNA gyrase and topoisomerase IV genes; reduced drug accumulation in bacterial cell by overexpression of efflux pump; impermeability of outer membrane and the presence of plasmid-mediated quinolone resistance genes (Jacoby, 2005). Among these, mutations in QRDR of DNA gyrase and topoisomerase IV genes are frequently present in quinolone-resistant bacteria (Wasyl et al., 2014; Garcia-Fernandez et al., 2015). QRDR is a small region located between amino acid 67-107 in GyrA and amino acid 426-447 in GyrB (Yoshida et al., 1991) and may be quinolone-binding site (Madurga et al., 2008). The amino acid substitutions in QRDR could result in reduced fluoroquinolone binding affinity, leading to decrease in fluoroquinolone susceptibility in bacteria (Vashist et al., 2009).

Quinolone resistant *Salmonella* has been previously reported in Thailand. However, most studies focused on the resistance phenotype (Minami et al., 2010; Sanpong et al., 2010; Lertworapreecha et al., 2013) and little published information on the quinolone resistance genotype is available. The present study provides an update on mutations in the QRDR of *gyrA*, *gyrB*, *parC* and *parE* genes in nalidixic acid and/or

ciprofloxacin-resistant *Salmonella* from pork, chicken meat and humans in Thailand.

A total of 28 nalidixic-resistant *Salmonella* isolates that were originated from faecal samples of humans (n=7), pigs (n=16) and chicken (n=5) in 2010-2011 (Table 1) were examined. The strains were isolated using ISO6579:2002 (E) (ISO, 2002) and serotyping at Center of Antimicrobial Resistance in Foodborne Pathogens (in cooperation with the World Health Organization), Faculty of Veterinary Science, Chulalongkorn University. These *Salmonella* isolates were previously tested against nalidixic acid (30 µg) using the CLSI guidelines for antimicrobial disc susceptibility testing (CLSI, 2008). All the isolates were stored at -80°C in 20% glycerol as our strain collection.

Materials and Methods

All the isolates were determined for minimum inhibitory concentration (MIC) of nalidixic acid and ciprofloxacin by two-fold agar dilution method as described by CLSI (CLSI, 2008). Clinical breakpoints for nalidixic acid and ciprofloxacin were ≥ 32 and ≥ 4 µg/ml, respectively. *Escherichia coli* ATCC® 25922, *Pseudomonas aeruginosa* ATCC® 27853, and *Staphylococcus aureus* ATCC® 29213 were included as quality control strains.

The QRDRs of *gyrA*, *gyrB*, *parC* and *parE* genes were PCR amplified by using specific primer pairs as follows: *gyrA*, *gyrA*-F (5'-GCTGAAGAGCTCCATCTGG-3')/*gyrA*-R (5'-GGTCGGCATGACGTCCGG-3'); *gyrB*, *gyrB*-F (5'-GCGCGCTCGATTTAGCCG-3')/*gyrB*-R (5'-TGATAGCGCAGCTTGTCCG-3'); *parC*, *parC*-F (5'-GTACGTGATCATGGATCGTG-3')/*parC*-R (5'-TTCCTGCATGGTGCCGTCG-3'); and *parE*, *parE*-F (5'-GCCATCGCGAATATCAGGCG-3')/*parE*-R (5'-CAGTTGTTCCAGTACGCC 3') (Chuanchuen and Padungtod, 2009). The PCR amplicons were gel purified using Nucleospin® Gel and PCR clean up (Düren, Germany) and submitted for DNA sequencing (First Base Laboratories, Selangor Darul Ehsan, Malaysia). All sequences obtained were compared with those in GenBank database using the Blast algorithm available at www.ncbi.nlm.nih.gov (Genbank accession numbers AE008801, AE008878 and AE008846 for *gyrA*, *gyrB*, *parC* and *parE*, respectively). Two *Salmonella* strains susceptible to nalidixic acid and ciprofloxacin were included as control strains.

Results

All the nalidixic acid-resistant *Salmonella* isolates had MIC value ranging from 32 to 256 µg/ml. Six isolates (4 human isolates and 2 chicken isolates) were additionally resistant to ciprofloxacin with MIC 4 µg/ml (Table 1).

Nucleotide sequencing analysis revealed that 11 isolates carried a single point mutation in *gyrA* including C248A, C248T, G259T and G259A, leading to amino acid substitutions Ser83Tyr, Ser83Phe, Asp87Tyr, Asp87Asn in GyrA, respectively (Table 1). Amino acid change at codon 83 (i.e. Ser83Phe and Ser83Tyr) was most commonly identified (n=8). Four isolates resistant to both nalidixic acid and ciprofloxacin harbored Ser83Phe (i.e. A50, A53 and

E21) or Asp87Tyr (i.e. E21). Seven *Salmonella* isolates resistant to nalidixic acid but susceptible to ciprofloxacin carried amino acid change Ser83Tyr (i.e. A43, C64, C92 and D10), Ser83Phe (i.e. E37), Asp87Tyr (i.e. D63) or Asp87Asn (i.e. A7). Twenty-two isolates

carried a single point mutation G283C in *parC*, leading to Val95Leu substitution in ParC, which was also found in the two susceptible-control strains. None of the isolates harbored mutations either in *gyrB* or *parE*.

Table 1 MIC values of nalidixic acid and ciprofloxacin in *Salmonella* isolates from humans, chicken meat and pork (n=28)

Source	Strain ID	Serotype	Sample type	Location	MIC ($\mu\text{g/ml}$) ^{a)}		Mutation ^{b)}
					Nalidixic acid	Ciprofloxacin	
Humans (n=7)	A43	Anatum	Rectal swab	Hospital	256	2	C248A(Ser83Tyr)
	A45	Weltevreden	Rectal swab	Hospital	256	0.125	- ^{c)}
	A50	Enteritidis	Rectal swab	Hospital	256	4	C248T(Ser83Phe)
	A53	Enteritidis	Rectal swab	Hospital	256	4	C248T(Ser83Phe)
	A54	Corvallis	Rectal swab	Hospital	32	4	-
	A60	Rissen	Rectal swab	Hospital	256	4	C248T(Ser83Phe)
	C48	Typhimurium	Rectal swab	Hospital	128	1	-
Chicken (n=5)	B36	Anatum	Carcass swab	Slaughterhouse	32	4	-
	C64	Amsterdam	Carcass swab	Slaughterhouse	128	1	C248A(Ser83Tyr)
	D63	Virchow	Chicken meat	Market	128	0.5	G259T(Asp87Tyr)
	D70	Agona	Chicken meat	Market	128	0.125	-
	E21	Virchow	Chicken meat	Market	128	4	G259T(Asp87Tyr)
Pig (n=16)	A4	Anatum	Carcass swab	Slaughterhouse	32	0.125	-
	A5	Anatum	Carcass swab	Slaughterhouse	32	0.25	-
	A7	Albany	Carcass swab	Slaughterhouse	128	0.5	G259A(Asp87Asn)
	C77	Weltevreden	Carcass swab	Slaughterhouse	32	0.125	-
	C78	Stanley	Carcass swab	Slaughterhouse	128	0.125	-
	C90	Anatum	Retail pork	Market	32	0.125	-
	C91	Corvallis	Retail pork	Market	32	2	-
	C92	Anatum	Retail pork	Market	>128	1	C248A(Ser83Tyr)
	D1	Corvallis	Retail pork	Market	32	2	-
	D5	Rissen	Retail pork	Market	32	0.125	-
	D10	Anatum	Retail pork	Market	128	1	C248A(Ser83Tyr)
	D11	Rissen	Retail pork	Market	128	1	-
	D54	Stanley	Retail pork	Market	128	1	-
	E16	Anatum	Retail pork	Market	128	1	-
	E37	Anatum	Retail pork	Market	128	1	C248T(Ser83Phe)
	E82	Worthington	Retail pork	Market	32	1	-

^{a)} The clinical breakpoints for nalidixic acid and ciprofloxacin were $\geq 32 \mu\text{g/ml}$ and $\geq 4 \mu\text{g/ml}$, respectively.

^{b)} nucleotide change in *gyrA* (amino acid substitution in GyrA)

^{c)} -, no mutation found

Discussion

Based on the CLSI ciprofloxacin breakpoint, only six nalidixic acid-resistant isolates in this collection were additionally resistant to ciprofloxacin. It indicates that resistance to nalidixic acid is not always associated with resistance to ciprofloxacin, consistent with previous studies (Hakanen et al., 1999; Kim et al., 2011). However, it has been suggested that CLSI interpretative criteria for ciprofloxacin (i.e. $\geq 4 \mu\text{g/ml}$) should be reevaluated (Ryan et al., 2011). Several studies showed that the nontyphoidal and typhoidal *Salmonella* isolates had the ciprofloxacin MIC much lower than 4 mg/liter (Kim et al., 2011; Wasyl et al., 2014) and a breakpoint of $\geq 0.125 \mu\text{g/ml}$ was recommended (Aarestrup et al., 2003). Based on the nucleotide sequencing analysis, similar mutations in GyrA (e.g. Asp87Asn, Ser83Phe, Asp87Thr) were observed in the ciprofloxacin-susceptible and -resistant isolates. Seven isolates resistant to nalidixic acid but susceptible to ciprofloxacin carried amino acid mutation at positions Ser83 and Asp87 in GyrA. It is possible that mutations may reduce the ciprofloxacin susceptibility at certain extent but not enough to bring

their MIC value above 4 $\mu\text{g/ml}$. These may be another support for the requirement of CLSI-ciprofloxacin breakpoint revision.

In addition, the isolates with high nalidixic acid MIC (128-256 $\mu\text{g/ml}$) exhibited varied ciprofloxacin MIC (0.125 to 4 $\mu\text{g/ml}$). Therefore, there was no correlation between nalidixic acid MIC and the raised ciprofloxacin MIC, in agreement with a previous study (Kim et al., 2011).

Mutations within the QRDR of *gyrA* are considered one of the main mechanisms of quinolone resistance. Amino acid substitutions at positions Ser83 and Asp87 in GyrA have been previously reported (Wasyl et al., 2014; Wang et al., 2015). The replacement of the hydrophilic hydroxyl group of serine by the hydrophobic residue of phenylalanine or tyrosine results in the loss of OH group of serine, leading to the loss of hydrogen-bonding interaction between amino acid residue and quinolone molecule and, eventually, reduced drug binding affinity with target site (Piddock, 2002). Similar phenomenon is applied for the amino acid change at Asp87. It was previously shown that a mutation at Ser83 conferred a higher quinolone resistance level, in comparison to that at position 87

(Ogbolu et al., 2012). However, it was not the case in this study. Further studies, e.g. site-directed mutagenesis, is required to better understand the contribution of mutations in these target genes.

All four *Salmonella* isolates resistant to both nalidixic acid and ciprofloxacin containing a single mutation at positions Ser 83 and Asp87 had nalidixic acid MIC value of ≥ 128 $\mu\text{g/ml}$ and ciprofloxacin MIC value of 4 $\mu\text{g/ml}$. This finding is consistent with a previous study showing that *S. Typhi* and *S. Paratyphi A* isolates carrying Ser83Phe had a high nalidixic acid resistance level (MIC > 256 $\mu\text{g/ml}$) with a ciprofloxacin MIC value of 4 $\mu\text{g/ml}$ (Hirose et al., 2002).

Previous studies demonstrated that high resistance level to fluoroquinolones was associated with double mutations in *gyrA* at positions 83 and 87 coupled with *parC* mutation (Cui et al., 2008; Garcia-Fernandez et al., 2015). However, double mutations in *gyrA* gene were not observed in our study. Most quinolone-resistant strains in the present study (n=22) carried amino acid substitution Val95Leu in ParC, which was also found in the quinolone-susceptible strains. Therefore, this amino acid change may be a result of biological variation among the *Salmonella* strains.

The lack of mutation in *gyrB* and *parE* in this study supports that the quinolone-resistant strain of *Salmonella* with mutations in *gyrB* or *parE* is rare (Yang et al., 2012; Lertworapreecha et al., 2013). In fact, the contribution of amino acid change in ParE to quinolone resistance remains unclear. In addition, some quinolone-resistant *Salmonella* isolates (n=17) lacked QRDR mutation of the tested genes, suggesting the existence of alternative resistance mechanisms that were not characterized in this study, e.g. efflux pump systems and plasmid-mediated quinolone resistance. This warranted further studies to elucidate the quinolone resistance mechanisms in these *Salmonella* isolates. Furthermore, the findings of the present study support that it is important to keep monitoring quinolone resistance-associated DNA gyrase mutations, particularly in *gyrA*, in order to predict the emergence of new mutations in quinolone-resistant *Salmonella* strains.

In conclusion, the results of this study highlight the complex pictures of quinolone resistance mechanisms in *Salmonella* isolate. Continuous monitoring of fluoroquinolone usage and resistance in *Salmonella* and other bacteria is indispensable.

Acknowledgements

This work was financially supported by the Research Grant for Mid-Career University Faculty RSA 5680051 cofunded by Thailand Research Fund (TRF), the Faculty of Veterinary Science, and Chulalongkorn University. It was also partially supported by the 90th Anniversary of Chulalongkorn University fund. NS is a recipient of the Royal Golden Jubilee PhD program PHD/006/2553, cofunded by TRF and Chulalongkorn University.

References

- Aarestrup FM, Wiuff C, Molbak K and Threlfall EJ 2003. Is it time to change fluoroquinolone breakpoints for *Salmonella* spp.? *Antimicrob Agents Chemother.* 47(2): 827-829.
- Chuanchien R and Padungtod P 2009. Antibiotic Resistance Genes in *Salmonella enterica* Isolates from Poultry and Swine. *J. Vet. Med. Sci.* 70: 1349-1355.
- CLSI 2008. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; Approved standard-Third edition M31-A3: Wayne, PA, USA. pp.
- Crement L, Caroff N, Dauvergne S, Reynaud A, Lepelletier D and Corvec S 2011. Prevalence of plasmid-mediated quinolone resistance determinants in ESBL *Enterobacteriaceae* clinical isolates over a 1-year period in a French hospital. *Pathol Biol (Paris).* 59(3): 151-156.
- Cui S, Li J, Sun Z, Hu C, Jin S, Guo Y, Ran L and Ma Y 2008. Ciprofloxacin-resistant *Salmonella enterica* serotype Typhimurium, China. *Emerg Infect Dis.* 14(3): 493-495.
- Fabrega A, Madurga S, Giralt E and Vila J 2009. Mechanism of action of and resistance to quinolones. *Microb Biotechnol.* 2(1): 40-61.
- Garcia-Fernandez A, Gallina S, Owczarek S, Dionisi AM, Benedetti I, Decastelli L and Luzzi I 2015. Emergence of Ciprofloxacin-Resistant *Salmonella enterica* Serovar Typhi in Italy. *PLoS One.* 10(6): e0132065.
- Hakanen A, Kotilainen P, Jalava J, Siitonen A and Huovinen P 1999. Detection of decreased fluoroquinolone susceptibility in *Salmonellas* and validation of nalidixic acid screening test. *J Clin Microbiol.* 37(11): 3572-3577.
- Hirose K, Hashimoto A, Tamura K, Kawamura Y, Ezaki T and Sagara H., & Watanabe, H 2002. DNA Sequence Analysis of DNA Gyrase and DNA Topoisomerase IV Quinolone Resistance-Determining Regions of *Salmonella enterica* Serovar Typhi and Serovar Paratyphi A. *Antimicrob Agents and Chemother.* 46(10): 3249-3252.
- Hopkins KL, Davies RH and Threlfall EJ 2005. Mechanisms of quinolone resistance in *Escherichia coli* and *Salmonella*: recent developments. *Int J Antimicrob Agents.* 25(5): 358-373.
- ISO 2002. Microbiology of food and animal feeding stuffs-Horizontal method for the detection of *Salmonella* spp: ISO6579-Fourth Edition. 1-27 pp.
- Jacoby GA 2005. Mechanisms of resistance to quinolones. *Clin Infect Dis.* 41 Suppl 2: S120-126.
- Kim KY, Park JH, Kwak HS and Woo GJ 2011. Characterization of the quinolone resistance mechanism in foodborne *Salmonella* isolates with high nalidixic acid resistance. *Int J Food Microbiol.* 146(1): 52-56.
- Lertworapreecha M, Sutthimusi S and Tontikapong K 2013. Antimicrobial resistance in *salmonella enterica* isolated from pork, chicken, and vegetables in southern Thailand. *Jundishapur J Microbiol* 6(6): 36-41.

- Madurga S, Sanchez-Cespedes J, Belda I, Vila J and Giralt E 2008. Mechanism of binding of fluoroquinolones to the quinolone resistance-determining region of DNA gyrase: towards an understanding of the molecular basis of quinolone resistance. *Chembiochem*. 9(13): 2081-2086.
- Minami A, Chaicupa W, Chongsa-Nguan M, Samosornsuk S, Monden S, Takeshi K, Makino S and Kawamoto K 2010. Prevalence of foodborne pathogens in open markets and supermarkets in Thailand. *Food Control*. 21(3): 221-226.
- Ogbolu DO, Daini OA, Ogunledun A, Alli AT, Olusoga-Ogbolu FF and Webber MA 2012. Effects of gyrA and parC Mutations in Quinolones Resistant Clinical Gram Negative Bacteria from Nigeria. *Afr J Biomed Res*. 15(2): 97-104.
- Piddock LJ 2002. Fluoroquinolone resistance in *Salmonella* serovars isolated from humans and food animals. *FEMS Microbiol Rev*. 26(1): 3-16.
- Ryan MP, Dillon C and Adley CC 2011. Nalidixic acid resistant strains of *Salmonella* showing decreased susceptibility to fluoroquinolones in the midwestern region of the Republic of Ireland due to mutations in the gyrA gene. *J. Clin. Microbiol*. 49(5): 2077-2079.
- Sanpong P, Theeragool G, Wajjwalku W and Amavisit P 2010. Characterization of multiple-antimicrobial resistant *Salmonella* isolated from pig farms in Thailand. *Kasetsart J*. 44(4): 643-651.
- Vashist J, Vishvanath, Kapoor R, Kapil A, Yennamalli R, Subbarao N and Rajeswari MR 2009. Interaction of nalidixic acid and ciprofloxacin with wild type and mutated quinolone-resistance-determining region of DNA gyrase A. *Indian J Biochem Biophys*. 46(2): 147-153.
- Wang Y, Yang B, Wu Y, Zhang Z, Meng X, Xi M, Wang X, Xia X, Shi X, Wang D and Meng J 2015. Molecular characterization of *Salmonella enterica* serovar Enteritidis on retail raw poultry in six provinces and two National cities in China. *Food Microbiol*. 46: 74-80.
- Wasył D, Hoszowski A and Zajac M 2014. Prevalence and characterisation of quinolone resistance mechanisms in *Salmonella* spp. *Vet Microbiol*. 171(3-4): 307-314.
- WHO 2007. Critically important antimicrobials for human medicine. Geneva, CH: Department of Food safety, zoonoses and foodborne diseases.
- Yang B, Xi M, Cui S, Zhang X, Shen J, Sheng M, Qu D, Wang X and Meng J 2012. Mutations in gyrase and topoisomerase genes associated with fluoroquinolone resistance in *Salmonella* serovars from retail meats. *Food Research International*. 45(2): 935-939.
- Yoshida H, Bogaki M, Nakamura M, Yamanaka LM and Nakamura S 1991. Quinolone resistance-determining region in the DNA gyrase gyrB gene of *Escherichia coli*. *Antimicrob Agents Chemother*. 35(8): 1647-1650.

บทคัดย่อ

การศึกษาการกลายพันธุ์ที่ตำแหน่ง QRDRs ของยีน DNA gyrase และ topoisomerase IV ของเชื้อแซลโมเนลลา เอนเทอริกาที่ดื้อต่อยากลุ่มควิโนโลนที่แยกได้จากเนื้อไก่ เนื้อสุกร และคน

นวลอนงค์ สินวัต¹ จิรัชญา พวงเสรี¹ สรรเพชญ์ อังกิตตระกูล² รุ่งทิพย์ ขวนชื่น^{1*}

เชื้อแซลโมเนลลา เอนเทอริกาที่ดื้อต่อยาในกลุ่มควิโนโลนจำนวน 28 isolates แยกมาจากเนื้อไก่ เนื้อสุกร และคน ตรวจหาการกลายพันธุ์ในตำแหน่ง quinolone resistant determining regions (QRDRs) ของยีน *gyrA*, *gyrB*, *parC* และ *parE* พบการเปลี่ยนแปลงของกรดอะมิโนแบบ single point mutation จำนวน 4 จุด (C248A, C248T, G259T และ G259A) ซึ่งทำให้เกิดการเปลี่ยนแปลงของกรดอะมิโนที่ตำแหน่ง Ser83Tyr, Ser83Phe, Asp87Tyr และ Asp87Asn บนโปรตีน GyrA ของเชื้อแซลโมเนลลาที่ดื้อต่อยากลุ่มควิโนโลนจำนวน 11 isolates โดยพบการเปลี่ยนแปลงของกรดอะมิโนที่ตำแหน่ง Ser 83 มากที่สุด นอกจากนี้ ไม่พบการกลายพันธุ์ในตำแหน่ง QRDRs ของยีน *gyrB* และ *parE* ทั้งนี้ ยังพบเชื้อแซลโมเนลลา ไทพิมุเรียมจำนวน 1 isolate ที่ดื้อต่อยานาลิดีซิก แอซิด ไม่พบการเปลี่ยนแปลงของกรดอะมิโนบนทุกยีนเป้าหมาย ซึ่งอาจเกิดมาจากกลไกอื่น ๆ ที่มีผลต่อการดื้อต่อยาในกลุ่มนี้

คำสำคัญ: ยีน DNA gyrase การดื้อต่อยาฟลูโอโรควิโนโลน การดื้อต่อยานาลิดีซิก แอซิด แซลโมเนลลา เอนเทอริกา ยีน topoisomerase IV

¹ภาควิชาสัตวแพทยสาธารณสุข หน่วยปฏิบัติการวิจัยความปลอดภัยอาหารทางจุลชีววิทยาและการดื้อยา คณะสัตวแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ปทุมวัน กรุงเทพฯ 10330

²ภาควิชาสัตวแพทยสาธารณสุข คณะสัตวแพทยศาสตร์ มหาวิทยาลัยขอนแก่น ขอนแก่น 40000

*ผู้รับผิดชอบบทความ E-mail: rchuanchuen@yahoo.com