

6-1-1988

Urinary enzymes and renal function changes followingVIPERA RUSSELLI SIAMENSIS snake venom injection in rats

Bungorn Chomdej

Waiter Pfaller

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Chomdej, Bungorn and Pfaller, Waiter (1988) "Urinary enzymes and renal function changes followingVIPERA RUSSELLI SIAMENSIS snake venom injection in rats," *Chulalongkorn Medical Journal*: Vol. 32: Iss. 6, Article 9.

Available at: <https://digital.car.chula.ac.th/clmjjournal/vol32/iss6/9>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Urinary enzymes and renal function changes following VIPERA RUSSELLI SIAMENSIS snake venom injection in rats.*

Bungorn Chomdej**
Walter Pfaller***

Chomdej B, Pfaller W. Urinary enzymes and renal function changes following VIPERA RUSSELLI SIAMENSIS snake venom injection in rats. Chula Med J 1988 Jun; 32(6): 573-578

The effects of Russell's viper venom (RVV) on urinary excretion of 4 enzymes and renal function changes were studied in 2 groups of 24 Wistar strain rats. Twelve control and 12 experimental rats were injected intraperitoneally with normal saline 1 ml/kg and RVV 1 mg/ml-kg body weight (bw) respectively. Urine collection was done in the period of 12 hours following injection. At the end of 12 hours, cardiac blood samples were taken after the rats were anesthetized with inactin at 120 mg/kg bw.

In the period of 12 hours following injection, the glomerular filtration rate (GFR) of experimental rats decreased significantly, whereas the urinary excretion of total protein increased ($p < .001$) when compared with control rats. The fractional excretion of sodium and potassium also increased significantly ($p < .001$). The urinary excretion of proximal tubular cytosolic enzyme fructose-1,6-biphosphatase (FBP), lysosomal N-acetyl- β -glucosaminidase (NAG) and glutathione-S-transferase (GST) rose drastically, while the pyruvate kinase (PK) fell significantly ($p < .001$).

From this study, both glomerular and tubular functions were reduced following venom injection. The enhanced urinary excretion of total protein and the reduction of GFR may have been due to the toxic damage of glomerular membrane leading to protein leakage into the urine. The increased proximal tubular enzyme excretion along with the fractional excretion of sodium and potassium might have reflected some destruction of proximal tubular cells. We concluded that RVV does exert a primary nephrotoxic action on both glomerular and tubular parts of the nephron. However, the secondary effect from its general circulation may also play a role.

Reprint requests: Chomdej B, Department of Physiology, Faculty of Medicine,
Chulalongkorn University, Bangkok 10500, Thailand.

Received for publication. October 5, 1987.

* Supported by the Austrian Fonds Zur Förderung der Wissenschaftlichen Forschung (project number P 5513.)

** Department of Physiology, Faculty of Medicine, Chulalongkorn University.

*** Institute of Physiology, University of Innsbruck, Austria.

บังอร ชมเดช, Pfaller W. ผลของพิษงูแมวเซาไทยต่อเอ็นซัยม์ในปัสสาวะและหน้าที่ของไตในหนูแรท.
จุฬาลงกรณ์เวชสาร 2531 มิถุนายน ; 32(6) : 573-578

ศึกษาผลของพิษงูแมวเซาไทยในหนูแรทพันธุ์วีสตาร์ 24 ตัว โดยแบ่งหนูออกเป็น 2 กลุ่มเท่า ๆ กัน กลุ่มควบคุมฉีคน้ำเกลือปกติ กลุ่มทดลองฉีคพิษงูขนาด 1 มิลลิกรัมต่อน้ำหนักตัว 1 กิโลกรัม โดยละลายในน้ำเกลือปกติเข้าทางช่องท้อง เก็บปัสสาวะในหนูทั้งสองกลุ่มตลอดระยะเวลา 12 ชั่วโมงภายหลังฉีค เมื่อครบ 12 ชั่วโมง นำหนูมาทำให้สลบโดยฉีคอินแอกซิน 120 มิลลิกรัมต่อน้ำหนักตัว 1 กิโลกรัม เจาะเลือดจากหัวใจนำไปปั่นแยกเอาพลาสมาออกทันทีทันใด

ผลการทดลองพบว่าการทำงานของไตในหนูกลุ่มที่ฉีคพิษงูลดลงไปอย่างมีนัยสำคัญทางสถิติ เมื่อเทียบกับกลุ่มควบคุม ($p < .001$) อัตราการกรองของไตลดลงพร้อมกับมีโปรตีนในปัสสาวะเพิ่มขึ้นแสดงถึงการทำลายผนังโกลเมอรูลอ อัตราการขับถ่ายต่ออัตราการกรองของโซเดียมและ โปแตสเซียมเพิ่มขึ้น พร้อมกับมีการขับถ่ายเอ็นซัยม์ของหลอดไตส่วนต้นในปัสสาวะมากขึ้นด้วย จึงพอจะสรุปได้ว่าพิษงูแมวเซามีผลทั้งโกลเมอรูลอและหลอดไตส่วนต้นโดยตรง ผลเสริมจากการเปลี่ยนแปลงของระบบไหลเวียนเลือดก็น่าจะมีส่วนเกี่ยวข้องด้วย

Clinical observations have shown that renal involvement is common following Russell's viper bites.^(1,2) The pathological changes of the kidney including acute tubular necrosis and cortical necrosis,^(3,4) severe glomerulonephritis⁽⁴⁾ and acute interstitial nephritis⁽⁵⁾ have been reported:

Many investigators believed that acute renal failure is secondary to intravascular hemolysis and intravascular coagulation.^(3,5) The hypotension following envenomation can cause a sudden decrease in renal blood flow leading to acute ischemic renal failure.^(7,9) However, acute renal failures have developed in some patients without hypotension⁽⁶⁾ and no correlation has been found between the severity of renal failure and coagulopathy.⁽³⁾

Enzymes along the nephron have been recorded in some conditions.⁽⁹⁻¹¹⁾ In patients with tubulointerstitial disease, enzymuria was frequent even in the absence of proteinuria.^(12,13) Fructose-1,6-biphosphatase (FBP), glutathione-S-transferase (GST), N-acetyl- β -glucosaminidase (NAG) are enzymes associated with the proximal tubular cells, while the pyruvate kinase (PK) is associated with the distal tubular cell functions.^(11,14)

The increased urinary excretion of renal enzymes in response to nephrotoxins may be indicative of various subcellular effects.⁽¹⁵⁾ NAG enzymuria is a sensitive indicator of the activity of renal disease and may prove to be a suitable screening test for significant renal diseases or injury,^(13,15,16) since it has been shown to be present in the renal tubules and to have a sufficiently high molecular weight not to ordinarily be filtered by the glomerulus.⁽¹³⁾ The high urinary excretion could indicate the conditions that produce primary glomerular as well as tubular damage.⁽¹³⁾

Renal function changes following injection of RVV in an animals have been reported.^(7,17,18) The direct or indirect toxicity or both have been attempted to be elucidated. This study investigated the role of venom on urinary excretion of four enzymes which may point out the mechanism of nephrotoxicity of the venom.

Materials and Methods

Four urinary enzymes and the renal functions were performed in the 2 groups of 24 Wistar strain rats, weighing 180-240 gm. Rats were fed with normal pellets in metabolic cage 2 days before the experiment. On the day of experiment, 12 control rats were injected with normal saline solution 1 ml/kg bw and 12 experimental rats were injected with lyophilized crude RVV, dissolved in normal saline 1 mg/ml-kg bw intraperi-

toneally. Urine samples were collected over a 12 hour period following injection. At the end of 12 hours, rats were anesthetized intraperitoneally with inactin 120 mg/kg bw. The whole cardiac blood was collected, centrifuged and separated for plasma sample at once.

Routine measurements of creatinine, sodium and potassium concentrations in plasma and urine were done. Total protein concentration in urine was also examined. Urinary enzyme activities were detected as described by Kotanko et al.⁽¹⁹⁾ One unit of enzyme activity was defined as the amount of enzyme catalyzing the formation of 1 μ mol reaction product per minute. Enzyme activities were related to urinary creatinine concentration in order to correct the differences in urinary volume output.^(13, 19)

The glomerular injury was indicated by the urinary excretion of protein, and the glomerular filtration rate was expressed as the clearance of creatinine. The tubular damage was exhibited by the fractional excretion of sodium, potassium and also 4 urinary enzyme activities.

The results were demonstrated as mean \pm S.D. The differences were analyzed by Student's unpaired t - test.

Results




Figure 1 The Effect of Russell's Viper Venom on Glomerular Filtration Rate (GFR)

As illustrated in figure 1, in the 12 hour period following the injection of normal saline and Russell's viper venom in control and experimental groups of rats, the glomerular filtration rate of the experimental rats fell significantly ($p < .001$) when compared with the control rats.




Figure 2 The Effect of Russell's Viper Venom on Urinary Total Protein.

Figure 2. showed the significant enhancement of urinary total protein during the 12 hours in the experimental rats compared to the control rats during the same period.




Figure 4 The Effect of Russell's Viper Venom on Fractional Excretion of Potassium (FEK)

Figures 3 and 4 demonstrated the fractional excretion of sodium and potassium in response to Russell's viper venom. The fractional excretion of both electrolytes of Russell's viper venom injected rats increased significantly when compared with the saline injected rats. The percent changes of fractional excretion of sodium were more than those of potassium.

As shown in the table, the proximal tubular enzymes : fructose-1, 6-biphosphatase, glutathione-S-transferase and N-acetyl-β-glucosaminidase increased drastically in the urine, whereas pyruvate kinase, the distal tubular enzyme, diminished in response to Russell's viper venom injection.




Figure 3 The Effect of Russell's Viper Venom on Fractional Excretion of Sodium (FENa)

Discussion

The RVV consists of a mixture of toxic proteins and enzymes which have hematotoxic and necrotizing properties.^(20,21) Acute renal failure is one of the clinical manifestations that may result from the toxicity.⁽²²⁾ The renal effects of the venom could be direct, or indirect as a consequence of shock, hemorrhage and vasculopathy.^(3, 23, 34)

Table The Effect of Russell's Viper Venom on Urinary Enzymes

Urinary Enzymes	mU/mg. Creatinine		
	Control group	Experimental group	p-value
F - 1,6 - Biphosphatase (FBP)	2.9 ± 0.3	17.4 ± 1.4	< .001
G - S - Transferase (GST)	2.9 ± 2.5	14,200 ± 6,900	< .001
Pyruvate Kinase (PK)	10.1 ± 2.6	1.7 ± 0.9	< .001
N - acetyl - β - Glucosaminidase (NAG)	34.1 ± 2.8	175.0 ± 13.1	< .001

In the present study, as shown in figures 1 and 2, RVV reduced GFR significantly along with producing proteinuria. It was possible that RVV interfered the filtering function by changing the glomerular permeability or surface area.⁽⁸⁾ The injurious effect of the venom could be mediated through complement activation,^(1, 20) causing glomerular membrane damage, hence resulting in an increase in urinary excretion of proteins.

Metabolism of renal cortex is characterized by fatty acid oxidation and gluconeogenesis. Proximal tubule is the main site of renal gluconeogenesis^(9, 11) and is the region in which gluconeogenesis is metabolically adapted.⁽⁹⁾ The specific enzymes from different parts of the nephron could be used as a marker of the damage.^(10, 11)

NAG possibly indicates the direct leakage with the damaged tubular lysosomes. It is also known that such lysosomes occur in renal tubular epithelial cells.^(25, 26) High levels appearing in the urine are always associated with active renal parenchymal diseases, particularly with those resulting from tubular injury.⁽²⁷⁾

The kidney contains the highest activities of enzymes involved in glutathione synthesis and degradation, with the highest level in the proximal straight tubule as an important site of renal drug metabolism.⁽¹⁴⁾ The pyruvate kinase, an enzyme catalyzing another irreversible step in the glycolytic pathway has the highest activity along the distal part of the nephron.^(11, 14)

As demonstrated in figures 3, 4 and the table, there were significant increases in FENa, FEK and

urinary enzymes of proximal tubular cells. The enhancement of FENa and FEK could be due to the inhibition of reabsorption function of proximal tubule by RVV.⁽¹⁷⁾ The probable damage was supported by the fact that urinary excretion of the proximal tubular cytosolic enzyme FBP and the lysosomal NAG were enhanced drastically. However, the decrement of PK in this experiment may help in the postulation that the activity of specific enzymes of glycolytic pathway is inhibited by some mechanisms.^(9, 14)

The correlation between proteinuria and increased enzymuria was also observed in patients with glomerulonephritis secondary to systemic vasculitis syndromes.⁽¹³⁾ RVV has also been reported to produce arteritis and severe glomerulonephritis.⁽⁴⁾ The increased urinary excretion of these enzymes in response to nephrotoxins may be indicative of various subcellular toxic effects.⁽¹⁵⁾ Nevertheless, it might be pointed out from this study that RVV plays an important role in damaging the kidney directly both at the glomerulus and proximal tubule, with some degrees of indirect toxicity.

Acknowledgement

The author (Dr. Bungorn Chomdej) wishes to thank Dr. Yachai Na Songkhla, the Dean, and Dr. Visith Sitprija, Associate Dean, for allocating the Staff Development Fund, Faculty of Medicine, Chulalongkorn University, that made this study possible. And also authors wish to thank Mr. Methee Visutthivarn for typing of this manuscript.

References

1. Sitprija V, Benyajati C, Boonpucknavig V. Renal involvement in snakebite. In : Ohsaka A, Hayashi K, Sawai Y, eds. *Animal Plant and Microbial Toxins*. New York : Plenum, 1976. 483-495
2. Shastry JCM, Date A, Carman RH, Johny KV. Renal failure following snake bite. A clinicopathological study of nineteen patients. *Am J Trop Med Hyg* 1977 Sep; 26(5) : 1032-1038
3. Chugh KS, Aikat BK, Sharma BK. Acute renal failure following snakebite. *Am J Trop Med Hyg* 1975 Jul; 24(4) : 692-697
4. Sitprija V, Boonpucknavig V. Extracapillary proliferative glomerulonephritis in Russell's viper bite. *Br Med J* 1980 Jun 14 : 2(6229) : 1417
5. Sitprija V, Suvanpha R, Pochanugool C, Chusil S, Tungsanga K. Acute interstitial nephritis in snake bite. *Am J Trop Med Hyg* 1982 Mar; 31(2) : 408-410
6. Sitprija V, Boonpucknavig V. Snake venoms and nephrotoxicity. In : Lee CY, ed. *Snake Venoms*. New York : Springer-Verlag, 1979. 997-1018
7. Tungthanathanich P, Chaiyabutr N, Sitprija V. Effect of Russell's viper (*Vipera Russellii Siamensis*) venom on renal hemodynamics in dogs. *Toxicon* 1986 Apr; 24(4) : 365-371
8. Schrier RW, Conger JD. Acute renal failure : Pathogenesis, diagnosis, and management. In : Schrier RW ed. *Renal and Electrolyte Disorders*. 3rd ed. Boston; Little Brown, 1986. 423-460

9. Burch HB, Narins RG, Chu C, Fagioli S, Choi S, McCarthy W, Lowry OH. Distribution along the rat nephron of three enzymes of gluconeogenesis in acidosis and sepsis. *J Physiol* 1978 Sep; 235(3) : 241-250
10. Guder WG, Auer U. The localization of gluconeogenesis in rat nephron. Determination of phosphoenolpyruvate carboxykinase in dissected tubules. *Hoppe-Seyler's Z Physiol Chem* 1974 Mar; 355 (3) : 273-278
11. Auer U, Guder WG. Site of enzyme activity along the nephron. *Kidney Int* 1976 Mar; 9(3) : 233-242
12. Burch HB, Hays AE, McCreary MD, Cole BR, Chi MM-Y, Dence CN, Lowry OH. Relationships in different parts of the nephron between enzymes of glycerol metabolism and the metabolite changes which result from large glycerol loads. *J Biol Chem* 1982 Apr; 257(7) : 3676-3679
13. Kunin CM, Chesney RW, Craig WA, England AC, DeAngelis C. Enzymuria as a marker of renal injury and disease : Studies of N-acetyl- β -glucosaminidase in the general population and in patients with renal disease. *Pediatrics* 1978 Nov; 62(5) : 751-760.
14. Guder WG, Ross BD. Enzyme distribution along the nephron. *Kidney Int*, 1984 Aug; 26(2) : 101-111
15. Catherine A, Suzuki M, Cherian MG. Renal toxicity of cadmium-metallothionein and enzymuria in rats. *J Pharmacol Expt Ther* 1987 Jan; 240(1): 314-319
16. Gibey R, Dupond JL, Alber D, Leconte des Floris R, Henry J-C. Predictive value of urinary N-acetyl-beta-D-glucosaminidase (NAG), alanine-aminopeptidase (AAP) and beta-2-microglobulin (β_2 M) in evaluating nephrotoxicity of gentamicin. *Clin Chim Acta* 1981 Oct; 116(1): 25-34.
17. Chomdej B, Pfaller W. The acute effect of Russell's viper venom on renal functions in rats. *Chula Med J* 1987 Apr; 31(4): 295-301
18. Huang HC. Effects of phospholipase A, from *Vipera Russellii* snake venom on blood pressure, plasma prostacyclin level and renin activity in rats. *Toxicon* 1984 Feb; 22 (2): 253-264
19. Kotanko P, Keiler R, Knabl L, Aulitzky W, Margreiter R, Gstraunthaler G, Pfaller W. Urinary enzyme analysis in renal allograft transplantation, *Clin Chim Acta* 1986 Oct; 160(2): 137-144
20. Devi A. The protein and nonprotein constituents of snake venoms. In: bucherl W, Buckley E, Deulofeu V, eds. *Venomous Animals and Their Venoms. Vol I. Venomous: Vertebrates*, 1968. 119-216
21. Iwanaga S, Suzuki T. Enzymes in snake venom. In: Lee CY, ed. *Snake Venoms*. New York: Springer-Verlag, 1979. 61-158
22. Schreiner GE, Maher JF. Toxic Nephropathy: review. *Am J Med* 1965 Mar; 38(3): 409-449
23. Linquist PA, Fujikawa K, Davie EW. Activation of bovine factor IX (Christmas factor) by factor XIa (Activated plasma thromboplastin antecedent) and a protease from Russell's viper venom. *J Biol Chem* 1978 Mar; 253(6): 1902-1909
24. Hogue-Angeletti RA, Bradshaw. Nerve growth factors in snake venoms. In: Lee CY, ed. *Snake Venoms*. New York: Springer-Verlag, 1979. 276-294
25. Wilkinson JH. Diagnosis significance of enzyme determinations in urine. In: Dubach UC, ed., *Enzymes in Urine and Kidney. Current Problems in Clinical Biochemistry. Vol 2* Baltimore: Williams and Wilkins, 1968. 207-235.
26. Raab WP. Enzymes and isoenzymes in urine. In: Dubach UC, ed., *Enzymes in Urine and Kidney. Current Problems in Clinical Biochemistry. Vol 2*. Baltimore: Williams and Wilkins, 1968. 17-83
27. Gonick HC, Kramer HJ, Schapiro AE. Urinary β -Glucuronidase activity in renal disease. *Arch Intern Med* 1973 Jul; 132(1): 63-69