

6-1-2020

Isolation and identification of extended spectrum β -lactamases (ESBLs) Escherichia coli from minced camel meat in Eastern province, Saudi Arabia

Waleed Rizk El-Ghareeb

Sherief M. Abdel-Raheem

Theeb M. Al-Marri

Fanan A. Alaql

Mahmoud M. Fayez

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

El-Ghareeb, Waleed Rizk; Abdel-Raheem, Sherief M.; Al-Marri, Theeb M.; Alaql, Fanan A.; and Fayez, Mahmoud M. (2020) "Isolation and identification of extended spectrum β -lactamases (ESBLs) Escherichia coli from minced camel meat in Eastern province, Saudi Arabia," *The Thai Journal of Veterinary Medicine*: Vol. 50: Iss. 2, Article 4.

Available at: <https://digital.car.chula.ac.th/tjvm/vol50/iss2/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Isolation and identification of extended spectrum β -lactamases (ESBLs) *Escherichia coli* from minced camel meat in Eastern province, Saudi Arabia

Waleed Rizk El-Ghareeb¹ Sherief M. Abdel-Raheem¹ Theeb M. Al-Marri²

Fanan A. Alaql³ Mahmoud M. Fayez^{2,4*}

Abstract

Antimicrobial resistance is an increasingly serious threat to global public health that requires action across all government sectors and society. The aim of this study was to determine the rate of extended-spectrum β -lactamases (ESBL)-producing *E. coli* isolation from minced camel meat and identify the phenotype and genotype of the ESBL. A total of 150 samples were collected randomly from butchers' shops in Al-Ahsa, Saudi Arabia. The results indicated that, overall, 17 (11.3 %) *E. coli* isolates were recovered from the minced meat samples. The isolates were classified biochemically at the species level using the VITEK 2 system. The antibiotic susceptibility of *E. coli* isolates was determined based on their MIC profile. The highest resistance was determined to be ampicillin (64.7%), doxycycline (23.5%), cefotaxime (23.5%) and ciprofloxacin (17.6%). Multidrug resistance (MDR) was determined in four isolates. Screening of the 17 isolates for ESBLs revealed that, four strains were resistant to cefotaxime and ceftazidime. A combination disk test (CDT) was used for ESBL phenotype conformation. The ESBL-encoding genes were characterized by PCR. The four isolates produced CTX-M group- 1 ESBLs. The *blaSHV* gene was detected in one isolate and *blaTEM* in two isolates. The *eaeA* gene was detected in 3 isolates, *stx2* gene in two isolates with the *hlyA* gene in one isolate. It can be concluded that there is clear evidence of the circulation of ESBLs producing *E. coli* in the minced camel meat. A high resistance was determined to ampicillin and doxycycline. The molecular detection of virulence genes may suggest the transmission of foodborne illness to consumers.

Keywords: *E. Coli*, β -lactamases, antibiotic resistance, genes, camel meat

¹Department of Public Health, College of Veterinary Medicine, King Faisal University, Saudi Arabia

² Ministry of Environment, Water and Agriculture, Al Ahsa Veterinary Diagnostic Laboratory, Saudi Arabia

³ Department of Microbiology, College of Science, King Saud University, Saudi Arabia

⁴Veterinary Serum and Vaccine Research institute, Ministry of Agriculture, Cairo, Egypt

*Correspondence: mahmoudfayez30@hotmail.com (M. M. Fayez)

Introduction

Inadequate use of antimicrobials in food-producing animals has been documented to be the main source of antimicrobial resistance (AMR) in commensal bacteria and foodborne pathogens (Friedman, 2015; ICF, 2017). Camel production in the KSA is considered as one of the most popular national animal productions over the last decades. In 2015, the total number of live camels in the KSA was estimated to be about 301717 head (MAWE, 2015; FAO, 2018). The concentration of amino acids and inorganic minerals in camel meat is higher, with less fat and higher moisture content than in beef. Moreover, Camel meat has reduced production costs because camels are usually reared by nomads in arid regions which makes camel meat available to consumers at a relatively low price (Kadim *et al.*, 2014).

Escherichia coli is a common bacteria among the intestinal microbiome of both humans and animals. *E. coli* had been isolated from healthy and diseased camels (Moore *et al* 2002, Salehi *et al.*, 2012, Al-Ruwaili *et al* 2012) moreover, the potential role of camels as a reservoir for ESBL *E.coli* has been investigated (Fadlelmula *et al.*, 2016). Multidrug resistant bacteria have been isolated from camels in the KSA due to overuse or prophylactic use of antibiotics (Fayez *et al.*, 2020).

Depending upon the disease's location, pathogenic *E. coli* are mostly divided into two groups: extra intestinal (ExPEC) and intestinal pathogenic *E. coli* (InPEC). Whereas, ExPEC strains are linked mainly with both neonatal meningitis (NMEC) and urinary tract infections (UPEC) in adults, InPEC strains are linked to diarrheal disease. InPEC strains are subdivided into at least 6 well-known path types: enter pathogenic *E. coli* (EPEC), enterohemorrhagic *E. coli* (EHEC), enter invasive *E. coli* (EIEC), enter aggregative *E. coli* (EAEC), diffusely adherent *E. coli* (DAEC) and enter toxigenic *E. coli* (ETEC). The emergence of antimicrobial resistance in *E. coli* has spread worldwide and their treatment has been greatly complicated by resistance to most first-line antimicrobial agents (Sabate *et al.*, 2008).

It has been reported that the ability of *E. coli* to produce broad spectrum beta lactamase is the main reason for the development of multidrug resistance (Zowawi *et al.*, 2013). Extended-spectrum-lactamases (ESBLs) which hydrolyze penicillins and expanded-spectrum cephalosporins appeared in the early 1980s and, recently, beta lactamase that hydrolyse carbapenems have become prominent (Kliebe *et al.*, 1985; Paterson, 2006). TEM, SHV, and CTX-M are the 3 main types of ESBLs. Depending on the dissimilarities of the amino acid sequence, the CTX-M enzymes have been subdivided into five groups (group 1: CTX-M-1, group 2: CTX-M-2, group 8: CTX-M-8, group 9: CTX-M-9, and group 25: CTX-M-25).

In Saudi Arabia, a number of surveys have investigated the prevalence of ESBLs producing bacteria in intensive care units, the community and medical wards in various locations, including Jeddah, Riyadh, Dhahran, Al Khobar and Al-Qassim (Al-Agamy *et al.*, 2009, Tewfik *et al.*, 2011, Zowawi *et al* 2013). Few reports in Saudi Arabia have investigated the microbiological quality of beef meat and poultry (Hemeg, 2018) and to the best of the author's

knowledge, there is no available literature on the prevalence ESBLs producing *E. coli* in minced camel meat. Therefore, the aim of this study was to determine the rate of extended-spectrum β -lactamases (ESBL)-producing *E. coli* isolation from minced camel meat and to identify the phenotype and genotype of the ESBL in Al Ahsa city, eastern province, Saudi Arabia.

Materials and Methods

Sample collection: Between January and September 2018, one hundred and fifty minced camel meat samples were randomly collected from 10 butchers' shops in Al Ahsa city, eastern province, Saudi Arabia. The Samples were collected in sterile, separate plastic bags and labeled. Then they were carried in an ice box within the hour to the Meat Hygiene laboratory, College of Veterinary Medicine, King Faisal University for bacteriological investigation. The samples were stored in a refrigerator at a temperature between 4° C and 8° C until examination within 24 h.

Isolation and identification of *E. coli*: Twenty-five grams (25g) of minced camel meat were weighted and suspended into 225 ml of 0.1% sterilized buffered peptone water (Hi Media, India) in a sterile stomacher bags and homogenized by shaking for 5 minutes. For enrichment purposes, the stomacher bags were incubated at 37 °C for 24 hours. A loop-full (10 μ l) from the overnight growth was streaked on MacConkey agar (Oxoid, England) and incubated at 37 °C for 24 hours. Permissive identification was performed based on colony morphology and gram staining. Five suspected colonies from each sample were purified on brain heart infusion agar (Oxoid, England) and examined for their oxidase activity by oxidase discs (Hi Media, India). Oxidase negative isolates were subjected to further biochemical identification by VITEK 2 compact system using ID-GN cards (bioMérieux, France).

Screening and confirmatory tests for extended-spectrum β -lactamases (ESBLs) producing *E. coli*: Screening for ESBLs was performed by the standard disc diffusion methods according to Clinical and Laboratory Standards Institute (CLSI) guideline (CLSI, 2014) using cefotaxime (30 μ g) and ceftazidime (30 μ g) discs. The bacterial inoculum was adjusted to be equivalent to 0.5 McFarland's standard and inoculated on to Muller-Hinton agar (Oxoid, England) by sterile cotton swabs then the two discs were placed on the inoculated plates and incubated at 35°C for 18 h. The measurements recommended by (CLSI) for isolates inhibition zone is \leq 27 mm for cefotaxime and \leq 22mm for ceftazidime were considered as ESBLs producer. The combination disk test (CDT) method (using both cefotaxime and ceftazidime, separately and in combination with clavulanate) recommended by CLSI was used for phenotypic conformation. The results were interpreted according to CLSI criteria (CLSI, 2014).

Antimicrobial susceptibility testing: The minimum inhibitory concentration (MIC) for nine antimicrobial agents, ampicillin, amoxicillin-clavulanate, cefotaxime,

imipenem, gentamicin, tetracycline, ciprofloxacin, trimethoprim-sulfamethoxazole and chloramphenicol was determined to isolated bacteria. Dilutions and cut off values were performed according to recommendations of (CLSI, 2014). Multidrug resistant (MDR) was considered when a strain demonstrated resistance to three or more antibiotic classes (Schwarz et al., 2010).

Molecular detection of 16SrRNA, virulence and antibiotic resistance genes: Total DNA was extracted from phenotypic identified ESBLs producing strains. QIAamp DNA mini-kit (Qiagen SA, France) was used for isolation and purification of DNA according to the manufacturer's instructions. For molecular conformation of the isolates, the 16SrRNA was amplified using universal primers 27F (5'-AGAGTTTGATCCTGGCTCAG-3') and 1492R (5'TACGGYTACCTTGTACGACTT-3') as described previously by (Lane, 1991). Virulence genes including *eaeA*, *hlyA*, *stx1*, *stx2* genes encoding intimin, enterohemolysin and shiga toxins were detected after the methods of (Gannon et al., 1997; Wang et al., 2002; Gallien, 2003). Screening for *blaCTX-M*, *blaTEM* and *blaSHV* genes was established by PCR using primers and reaction conditions previously described by (Pitout et al., 1998, 2004). The PCR products were purified for partial sequencing using a QIAquick PCR Purification Kit (Qiagen SA, Courtaboeuf, France). Sequencing was conducted by Genetic Analyzer 3500, (Applied Biosystems). Partial nucleotide sequences were analyzed using the BLAST program <https://blast.ncbi.nlm.nih.gov/Blast.cgi>.

Results

Bacterial isolation and screening for ESBLs: Overall, 17 (11.3%) *E. coli* isolates were recovered from the minced meat samples. The isolates were classified biochemically to the species level based on ID-GN database of the VITEK 2 compact system. Screening of the 17 isolates for ESBLs revealed that, four strains had an inhibitory zone less than 27 and 22 for cefotaxime and ceftazidime, respectively. Phenotypical

confirmation was performed for all suspected isolates and a synergy effect when tested with the double disk synergy method was noticed. A ≥ 5 mm increase in zone diameter was determined for either antimicrobial agent tested in combination with clavulanate vs the zone diameter of the agent or when tested alone.

Antimicrobial susceptibility: The antibiotic susceptibility of *E. coli* isolates was determined based on their MIC profile. The percentage of resistance against the 9 antimicrobials is presented in Table (1). Six isolates (35.3%) were susceptible to all antimicrobials. The highest resistance was determined to ampicillin (64.7%), doxycycline (23.5%), and cefotaxime (23.5%). Three isolates (17.6) showed resistance to ciprofloxacin. The resistance profile of all isolates is shown in Table (2). Multidrug resistance (MDR) was determined in four isolates two of which were identified as ESBLs producer.

Molecular detection of 16SrRNA, virulence and antibiotic resistance genes: The 16SrRNA sequence of the 4 ESBLs producer strains showed ≥ 98 % similarity with sequences of their corresponding strains from GenBank. The sequences were submitted to GenBank as accession numbers (GenBank MN658514.1, MN658513.1, MN658485.1 and MN658479.1).

The ESBL-encoding genes were characterized by PCR. The four isolates produced CTX-M group- 1 ESBLs. Sequences from PCR products were submitted to Gene Bank with accession numbers (GenBank MN690604, MN690605, MN690606 and MN690607). Fig (1) shows the molecular phylogenetic analysis by maximum Likelihood method. *blaSHV* gene was detected in one isolate whereas *blaTEM* was in two isolates. The *eaeA* gene was detected in 3 isolates, *stx2* gene in two isolates with the *hlyA* gene in one isolate. The distribution of different genes in the 4 ESBLs producer isolates is presented in Table (2). The different virulence gene sequences were submitted to GenBank with accession numbers (MN708336, MN708337, MN708338, MN708339, MN708340 and MN708341).

Table 1 Resistance proportion, MIC50 and MIC90 of *E. coli* isolates in minced camel meat in Al Ahsa province, Saudi Arabia.

Antibiotic	Total R (%)	MIC (g/ml)		
		Range	50%	90%
AMC	5.90% [95% CI, 0.3-30.8]	1 - 64	2	4
Ampicillin	64.70% [95% CI, 38.6-84.7]	1 - 128	64	128
Chloramphenicol	5.90% [95% CI, 0.3-30.8]	1 - 64	2	4
Ciprofloxacin	17.60% [95% CI, 4.6-44.1]	0.25 - 4	0.5	4
Cefotaxime	23.50% [95% CI, 7.8-50.2]	0.25 - 16	0.5	16
Doxycycline	23.50% [95% CI, 7.8-50.2]	0.5 - 32	2	32
Gentamicin	17.60% [95% CI, 4.6-44.1]	1 - 64	2	32
Imipenem	0.00% [95% CI, 0.0-22.9]	0.125 - 0.5	0.25	0.5
SXT	11.80% [95% CI, 2.1-37.8]	0.5 - 16	1	8

Table 2 Antimicrobial resistance profile, virulence genes and ESBL genotype of *E. coli* isolates in minced camel meat in Al Hsa province, Saudi Arabia.

Isolate ID	ESBL phenotype			ESBL genotype			Antimicrobial resistance profile			Virulence genes			
	ESBL phenotype	<i>bla</i> CTX-M	<i>bla</i> TEM	<i>bla</i> SHV	MDR	<i>eaeA</i>	<i>hlyA</i>	<i>stx1</i>	<i>stx2</i>				
CM1	Positive	Positive	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM2	Positive	Positive	Negative	Positive	MDR	Positive	Positive	Positive	Negative				
CM3	Positive	Positive	Positive	Negative	MDR	Positive	Negative	Negative	Negative				
CM4	Positive	Positive	Positive	Negative	MDR	Positive	Negative	Positive	Negative				
CM5	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM6	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM7	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM8	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM9	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM10	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM11	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM12	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM13	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM14	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM15	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM16	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				
CM17	Negative	Negative	Negative	Negative	MDR	Negative	Negative	Negative	Negative				




Figure 1 Maximum likelihood tree of BlaCTX-M ESBLs nucleotide sequences. A phylogenetic tree of CTX-M group 1

Discussion

The health of humans, animals and ecosystem is interconnected, pointing to the need for a "One Health" approach. Food-borne diseases and antimicrobial resistance are prominent examples of these close inter-linkages. Food becomes an important vehicle for the transmission of commensal as well as pathogenic micro-organisms including zoonotic agents. Pathogen transmission between food and humans occurs during the processing of raw materials as well as cross contamination at production and distribution (MacDonald *et al.*, 2015).

During this work, *E. coli* was isolated from 17 minced camel meat samples (11.3%). The high proportion of protein and water in fresh red meat provides a suitable condition for growth and multiplication of bacteria (ICMSF, 2005). *E. coli* was previously isolated from fresh beef meat and minced beef meat in the KSA (Iyer *et al.*, 2013; Hemeg, 2018). The variation in the isolation frequencies may be attributed to the sample size, isolation techniques, use of selective media animal species, meat handling and sanitary standards of meat retailer shop premises.

In this study, the MIC of nine antibiotics were determined for all isolates. High resistance to ampicillin and doxycycline was observed and the findings are consistent (Hemeg, 2018, Abayneh *et al.*, 2019). Penicillins and tetracycline are widely used in the treatment of animals in the study area. Three isolates (17.6%) showed resistance to ciprofloxacin. The World Health Organization has classified fluoroquinolones as critically important drugs for human medicine due to a strong link between their use and increased resistance, and has advised careful use of fluoroquinolones in both human and veterinary medicine (WHO, 2012). The possibility of transmitting antibiotic resistance genes from commensals to pathogens or between pathogens has been documented elsewhere (Friedman, 2015; ICF, 2017). The isolates showed 7 resistance profiles and 4 profiles

showed MDR. Excessive and inappropriate antimicrobials could be factors enhancing the positive selection of antimicrobial resistance bacteria as a part of commensal flora with an accumulation of antimicrobial resistance genes that encode multiple resistance traits (Nikaido, 2009).

Screening for ESBLs revealed identification of 4 (2.6%) ESBLs producer *E. coli* from minced camel meat. In enterobacteriaceae, the production of beta lactamase is the main mechanism for antibiotic resistance that is a major health issue in the world. The frequency of isolation in this study was lower than other studies elsewhere (Ye *et al.*, 2018, Abayneh *et al.*, 2019, Le *et al.*, 2015). This variation may be attributed to the inappropriate use of fourth and third generations of cephalosporins and the difference in methodology mentioned above.

The molecular characterization of the BlaCTX-M gene revealed that it belongs to the CTX-M group 1. This group includes six plasmid-mediated enzymes; CTX-M-1, CTX-M-3, CTX-M-10, CTX-M-12, CTX-M-15, and FEC-1 (Bonnet 2004). Since there tend to be regional differences in the occurrence of various ESBL variants (Cantòn and Coque, 2006), the current study provides further data relating to minced camel meat. The CTX-M group 1 variants (CTX-M-1 and CTX-M-15) were previously identified in hospitalized people in the Riyadh and Al-Qassim regions (Al-Agamy *et al.*, 2009, Tewfik *et al.*, 2011). In this work, the *eaeA*, *hlyA* and *stx2* genes encoding intimin, enterohemolysin and shiga toxins of *E. coli* were identified by PCR in three ESBLs producer isolates, Table. (2). Intimin is a virulence factor on the outer membrane of pathogenic *E. coli* and is responsible for attachment and adhesion of the bacteria to host cells. Enterohemolysin is a potential virulence factor for pathogenic *E. coli* that is often connected with severe human diseases such as hemorrhagic colitis and hemolytic uremic syndrome. Shiga toxin producing *E. coli* is considered one of the most crucial causes of food-borne illness that can lead to life-threatening complications such as hemolytic-

uremic syndrome (Liptakova *et al.*, 2002). Domestic ruminants, mainly cattle, sheep and goats, have been reported to be major natural reservoirs for shiga toxin producing *E.coli* and play a significant role in the epidemiology of human infections (Islam *et al.*, 2008 and Darwish *et al.*, 2018). Isolation of shiga toxin producing *E.coli* from feces of camels has been reported elsewhere (Martin, and Beutin 2011; Baschera *et al.*, 2019). Contamination of minced meat with pathogenic *E coli* may occur during handling in abattoirs or butcher markets (Kaper *et al.*, 2004; Vincent *et al.*, 2010). Good, widespread sanitation and hygiene is recommended during food processing to prevent fecal contamination of foodstuffs. Clean and convenient restroom facilities and routine cleaning and sanitation of the distribution and transport environment, processing sites and food contact surfaces are essential for preventing cross contamination.

In conclusion, from the obtained results, it can be concluded that there is clear evidence of the circulation of ESBLs producing *E. coli* in the minced camel meat. A high resistance was determined to ampicillin and doxycycline. Resistance of non ESBLs isolates to fluoroquinolones is a serious public health problem. Isolation of multidrug resistance bacteria from minced meat may be a reflection of the improper and excessive use of antibiotics in food producing animals. The ESBLs producing *E. coli* isolates were genotyped as blaCTX-M group 1. The molecular detection of virulence genes may suggest the transmission of foodborne illness to consumers. Therefore, approaches to enhance the knowledge and practice of butchers in the handling and storage of meat should be prepared and enforced. In addition, monitoring the prevalence of antimicrobial resistance among isolates from healthy animals and their food products provides evidence of the need for the design of a strategy for the prevention and control of the spread resistance strain in the community.

Acknowledgements

The authors would like to thank and acknowledge the Deanship of Scientific Research at King Faisal University, Saudi Arabia for supporting this research.

References

- Abayneh M, Tesfaw G, Woldemichael K, Yohannis M, Abdissa A. 2019. Assessment of extended-spectrum β -lactamase (ESBLs)-producing *Escherichia coli* from minced meat of cattle and swab samples and hygienic status of meat retailer shops in Jimma town, Southwest Ethiopia. *BMC Infect Dis.* 19 (1):897.
- Al-Agamy MH, Shibl AM, Tawfik AF. 2009. Prevalence and molecular characterization of extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* in Riyadh, Saudi Arabia. *Ann. Saudi Med.* 29(4):253-7.
- Al-Ruwaili MA, Khalil OM, Selim SA 2012. Viral and bacterial infections associated with camel (*Camelus dromedarius*) calf diarrhea in North Province, Saudi Arabia. *Saudi journal of biological sciences*, 19(1), 35-41. <https://doi.org/10.1016/j.sjbs.2011.10.001>.
- Baschera M, Cernela N, Stevens M, Liljander A, Jores J, Corman VM, Nüesch-Inderbinnen M, Stephan R (2019). Shiga toxin-producing *Escherichia coli* (STEC) isolated from fecal samples of African dromedary camels. *One health (Amsterdam, Netherlands)*, 7, 100087.
- Bonnet R (2004). Growing group of extended-spectrum beta-lactamases: the CTX-M enzymes. *Antimicrobial agents and chemotherapy*, 48(1), 1-14. <https://doi.org/10.1128/aac.48.1.1-14.2004>.
- Cantón R, Coque TM. 2006. The CTX-M β -lactamase pandemic. *Curr Opin. Microbiol.* 9 (5):466-75.
- Clinical and Laboratory Standards Institute (CLSI) 2014. Wayne, pa: Performance standards for antimicrobial susceptibility testing of anaerobic bacteria: informational supplement. Twenty-Fourth Informational Supplement 2014. Approved M100S24.
- Darwish WS, Atia AS, El-Ghareeb WR, Elhelaly AE (2018). Prevalence of multidrug resistant shiga toxin-producing *Escherichia coli* in cattle meat and its contact surfaces. *Journal of Food Quality and Hazards Control.* 5: 146-153.
- Fadlelmula A, Al-Hamam NA, Al-Dughaym AM. A potential camel reservoir for extended-spectrum β -lactamase-producing *Escherichia coli* causing human infection in Saudi Arabia. *Trop Anim Health Prod.* 2016;48 (2):427-433. [doi:10.1007/s11250-015-0970-9](https://doi.org/10.1007/s11250-015-0970-9).
- FAO. 2018. Statistics Reports. Food and Agricultural Organization, Rome. Italy.
- Fayez, M., Elsohaby, I., Al-Marri, T., Zidan, K., Aldoweriej, A., El-Sergany, E., & Elmoslemany, A. (2020). Genotyping and antimicrobial susceptibility of *Clostridium perfringens* isolated from dromedary camels, pastures and herders. *Comp Immunol Microbiol Infect Dis*, 70, 101460 <https://doi.org/10.1016/j.cimid.2020.101460>.
- Friedman M. 2015. Antibiotic-resistant bacteria: prevalence in food and inactivation by food-compatible compounds and plant extracts. *Journal of agricultural and food chemistry.* 9; 63 (15):3805-22.
- Gallien P.2003. Detection and Subtyping of ShigaToxin-Producing *Escherichia coli* (STEC). *Methods Mol Biol.* 216:163-84.
- Gannon VP, D'souza S, Graham T, King RK, Rahn K, Read S. 1997. Use of the flagellar H7 gene as a target in multiplex PCR assays and improved specificity in identification of enterohemorrhagic *Escherichia coli* strains. *J Clin Microbiol.* 35(3):656-62.
- Hemeg HA. 2018. Molecular characterization of antibiotic resistant *Escherichia coli* isolates recovered from food samples and outpatient Clinics, KSA. *Saudi J. Biol. Sci.* 25 (5):928-31.
- ICF I. 2017. EU Insights - Perceptions on the Human Health Impact of Antimicrobial Resistance (AMR) and Antibiotics Use in Animals across the EU. EFSA Supporting Publication EN-1183, N-1162.
- International Commission on Microbiological Specifications for Foods (ICMSF) 2005. Microorganisms in foods 6: Microbial ecology of food commodities. Second Edition. New York: Kluwer Academic/Plenum Publishers; 2005.

- Islam, M. A., Mondol, A. S., de Boer, E., Beumer, R. R., Zwietering, M. H., Talukder, K. A., & Heuvelink, A. E. (2008). Prevalence and genetic characterization of shiga toxin-producing *Escherichia coli* isolates from slaughtered animals in Bangladesh. *Appl Environ Microbiol*, 74 (17), 5414-21 <https://doi.org/10.1128/AEM.00854-08>.
- Iyer A, Kumosani T, Yaghmoor S, Barbour E, Azhar E, Harakeh S. 2013. *Escherichia coli* and *Salmonella* spp. in meat in Jeddah, Saudi Arabia. *J Infect Dev Countr*. 7 (11):812-8.
- Kadim, I.T., Mahgoub, O., Mbagha, M. 2014. Potential of camel meat as a non-traditional high quality source of protein for human consumption. *Animal Frontiers*, 4 (4), 13-17.
- Kaper JB, Nataro JP, Mobley HL. 2004 Pathogenic *Escherichia coli*. *Nat. Rev. Microbiol*. 2 (2):123-140.
- Kliebe C, Nies BA, Meyer JF, Tolxdorff-Neutzling RM, Wiedemann B. 1985. Evolution of plasmid-coded resistance to broad-spectrum cephalosporins. *Antimicrob. Agents Chemother*. 28 (2):302-7.
- Lane DJ. 1991. 16S/23S rRNA Sequencing. In: Stackebrandt, E. and Goodfellow, M., Eds., *Nucleic Acid Techniques in Bacterial Systematic*, John Wiley and Sons, New York, 115-175.
- Le HV, Kawahara R, Khong DT, Tran HT, Nguyen TN, Pham KN, Jinnai M, Kumeda Y, Nakayama T, Ueda S, Yamamoto Y. 2015. Widespread dissemination of extended-spectrum β -lactamase-producing, multidrug-resistant *Escherichia coli* in livestock and fishery products in Vietnam. *Int. j. food contam*. 2 (1):17.
- Liptáková A, Siegfried L, Sabol M, Sehnálková H, Bogyiová E, Rosocha J, Kmeťová M, Kerestešová H, Kotulová D. 2002. Detection of shiga toxins, intimin and enterohemolysin in *Escherichia coli* strains isolated from children in eastern Slovakia. *Folia Microbiol* 47 (2):185.
- MacDonald E, White R, Mexia R, Bruun T, Kapperud G, Lange H, Nygård K, Vold L. 2015. Risk factors for sporadic domestically acquired *Campylobacter* infections in Norway 2010–2011: A national prospective case-control study. *PLoS One* 10 (10):e0139636.
- Martin, A., and Beutin, L. (2011). Characteristics of Shiga toxin-producing *Escherichia coli* from meat and milk products of different origins and association with food producing animals as main contamination sources. *Int J Food Microbiol*, 146 (1), 99-104.
- MAWE. 2015. *Livestock Statistics of 2015 Agriculture Census*. Ministry of Agriculture, Water and Environment, Riyadh, KSA.
- Moore, J. E., McCalmont, M., Jiru, Xu, Nation, G., Tinson, A. H., Crothers, L., et al. (2002). Prevalence of faecal pathogens in calves of racing camels (*Camelus dromedaries*). *Tropical Animal Health and Production*, 34, 283–287.
- Nikaido H. 2009. Multidrug resistance in bacteria. *Annu. Rev. Biochem*. 78:119-46. doi:10.1146/annurev.biochem.78.082907.145923.
- Paterson DL. 2006. Resistance in gram-negative bacteria: Enterobacteriaceae. *Am J Infect Control* 34(5):S20-28, S64 –S73.
- Pitout JD, Hossain A, Hanson ND. 2004. Phenotypic and molecular detection of CTX-M- β -lactamases produced by *Escherichia coli* and *Klebsiella* spp. *J Clin Microbiol*. 42 (12):5715-21.
- Pitout JD, Thomson KS, Hanson ND, Ehrhardt AF, Moland ES, Sanders CC. 1998. β -Lactamases responsible for resistance to expanded-spectrum cephalosporins in *Klebsiella pneumoniae*, *Escherichia coli*, and *Proteus mirabilis* isolates recovered in South Africa. *Antimicrob. Agents Chemother*. 42 (6):1350-1354.
- Sabaté M, Prats G, Moreno E, Ballesté E, Blanch AR, Andreu A. 2008. Virulence and antimicrobial resistance profiles among *Escherichia coli* strains isolated from human and animal wastewater. *Res. Microbiol*. 159(4):288-93.
- Salehi TZ, Tonelli A, Mazza A, Staji H, Badagliacca P, Tamai IA, Jamshidi R, Harel J, Lelli R, Masson L 2012 Genetic characterization of *Escherichia coli* O157:H7 Strains isolated from the one-humped camel *Camelus dromedarius* by using microarray DNA technology. *Molecular Biotechnology* 51:283–288. <https://doi.org/10.1007/s12033-011-9466-7>.
- Schwarz S, Silley P, Simjee S, Woodford N, van Duijkeren E, Johnson AP, Gaastra W. Assessing the antimicrobial susceptibility of bacteria obtained from animals. *J Antimicrob Chemother*. 141 (1-2):1-4-4.
- Tawfik AF, Alswailem AM, Shibl AM, Al-Agamy MH. 2011. Prevalence and genetic characteristics of TEM, SHV, and CTX-M in clinical *Klebsiella pneumoniae* isolates from Saudi Arabia. *Microb Drug Resist*. 17(3):383-8.
- Vincent C, Boerlin P, Daignault D, Dozois CM, Dutil L, Galanakis C, Reid-Smith RJ, Tellier PP, Tellis PA, Ziebell K, Manges AR. 2010. Food reservoir for *Escherichia coli* causing urinary tract infections. *Emerging infectious diseases*. *Emerg Infect Dis*; 16 (1):88–95.
- Wang G, Clark CG, Rodgers FG. 2002. Detection in *Escherichia coli* of the genes encoding the major virulence factors, the genes defining the O157: H7 serotype, and components of the type 2 Shiga toxin family by multiplex PCR. *J. Clin. Microbiol*. 40 (10):3613-9.
- World Health Organization 2012. WHO Advisory Group on Integrated Surveillance of Antimicrobial Resistance (AGISAR). Critically important antimicrobials for human medicine 3rd Revision 2011. WHO Document Production Services, Geneva, Switzerland. *Clin. Infect. Dis*. 55, 712–719.
- Ye Q, Wu Q, Zhang S, Zhang J, Yang G, Wang J, Xue L, Chen M. 2018. Characterization of extended-spectrum β -lactamase-producing Enterobacteriaceae from retail food in China. *Front Microbiol*. 9:1709.
- Zowawi HM, Balkhy HH, Walsh TR, Paterson DL. 2013. β -Lactamase production in key gram-negative pathogen isolates from the Arabian Peninsula. *Clin Microbiol Rev*. 26(3):361-80.