

12-1-2020

Co-circulation of canine chaphamaparvovirus and canine parvovirus 2 in dogs with diarrhea in Turkey

Ender Dinçer

Mehmet Özkan Timurkan

Pelin Fatoş Polat Dinçer

Hakan Aydın

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Dinçer, Ender; Timurkan, Mehmet Özkan; Fatoş Polat Dinçer, Pelin; and Aydın, Hakan (2020) "Co-circulation of canine chaphamaparvovirus and canine parvovirus 2 in dogs with diarrhea in Turkey," *The Thai Journal of Veterinary Medicine*: Vol. 50: Iss. 4, Article 7.

Available at: <https://digital.car.chula.ac.th/tjvm/vol50/iss4/7>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Co-circulation of canine chaphamaparvovirus and canine parvovirus 2 in dogs with diarrhea in Turkey

Ender Dinçer^{1*} Mehmet Özkan Timurkan² Pelin Fatoş Polat Dinçer³ Hakan Aydın²

Abstract

Chaphamaparvoviruses, a divergent group of parvoviruses (family *Parvoviridae*) infect both domestic and wild mammalian and avian species, including bats, chickens and pigs. They have recently been identified using metagenomic analysis from tissue and stool samples of healthy and sick animals. Here, dogs (60 healthy and 43 diarrheal) were investigated for the presence of canine chaphamaparvovirus, canine coronavirus (CCoV), canine adenovirus (CAV), canine distemper virus (CDV) and canine parvovirus 2 (CPV-2) by rectal swabs using polymerase chain reaction (PCR). For chaphamaparvovirus, all the healthy dogs tested negative whereas three of the diarrheal dogs tested positive (3 /103; 2.9%). CPV-2 was also detected in diarrheal dogs (28/43; 65%) but no other viruses were found in the rectal samples of the sick dogs. The three chaphamaparvovirus-positive dogs were also positive for CPV-2. Phylogenetic analysis showed that the chaphamaparvoviruses were all related to American strains, forming a separate clade in the mammalian group. Amino acid sequence comparisons demonstrated that the Turkish strains had one substitution at 743 (Ser→Gly) and substitutions at 688 (Ser→Leu), 716 (Asp→His) and 743 (Cys→Ser/Gly) compared to the American strains. This study provides the first report of chaphamaparvovirus in Turkey. It also documents, for the first time, co-circulation of chaphamaparvovirus and CPV-2 in Turkey's dog population.

Keywords: Parvoviridae, chaphamaparvovirus, CPV-2, Molecular characterization, Turkey

¹Department of Virology, Faculty of Veterinary Medicine, Dokuz Eylül University, İzmir, Turkey

²Department of Virology, Faculty of Veterinary Medicine, Atatürk University, Erzurum, Turkey

³Department of Internal Medicine, Faculty of Veterinary Medicine, Dokuz Eylül University, İzmir, Turkey

*Correspondence: enderdin@gmail.com (E. Dinçer)

Received July 24, 2020.

Accepted September 22, 2020.

Introduction

The global dog population is estimated to be 900 million (Atitwa 2018). Many viral agents infect dogs, causing respiratory, gastrointestinal and nervous system diseases, including canine herpes virus (CHV), canine parainfluenza virus (CPIV), canine distemper virus (CDV), rabies virus, canine oral papillomavirus (CPV1), canine influenza (CIV), canine circovirus (canine CV), canine bocavirus (CBoV) and canine kobuvirus (CaKoV) (Aydin *et al.*, 2018; Caddy 2018; Thaiwong *et al.*, 2018). Canine diarrhea is one of the most significant diseases in dogs, whether caused by viruses, bacteria or parasites (Cave *et al.*, 2002). The main enteric viruses in dogs are canine adenovirus (CAV), canine corona virus (CCoV), mamastrovirus 5 (MAstV5) and canine parvovirus 2 (CPV-2) (De Benedictis *et al.*, 2011; Polat *et al.*, 2019; Timurkan *et al.*, 2018; Yesilbag *et al.*, 2004).

Parvoviruses are small, icosahedral symmetric, non-enveloped, single-stranded linear DNA viruses approximately 5 kb in genome length. Structurally, the genome has two Open Reading Frames (ORF): a non-structural replicase gene (NS) expressing enzyme required for replication and a capsid (VP) gene expressing structural proteins (Calderon *et al.*, 2009; Miranda & Thomson 2016). The family *Parvoviridae* contains three subfamilies, *Parvovirinae*, *Densovirinae* and *Hamaparvovirinae* which infect vertebrate and invertebrate hosts. Chaphamaparvoviruses classified in *Hamaparvovirinae* infect vertebrate hosts. (Cotmore *et al.*, 2014). The subfamily *Parvovirinae* comprises 8 genus: *Dependoparvovirus*, *Copiparvovirus*, *Bocaparvovirus*, *Amdoparvovirus*, *Aveparvovirus*, *Protoparvovirus*, *Tetraparvovirus*, and *Erythroparvovirus*. More recently, two new genus in the subfamily *Hamaparvovirinae* were added: *Marinoparvovirus* and *Chaphamaparvovirus*. *Marinoparvovirus* includes sesavirus, detected from rectal swabs of California sea lion offspring with malnutrition and pneumonia (Phan *et al.*, 2015).

Metagenomic analysis has recently increased the number of identified chaphamaparvoviruses isolated from bat, bird and pig hosts (Palinski *et al.*, 2016; Reuter *et al.*, 2014). Since then, red - crowned crane parvovirus (RcPV) and Turkey parvovirus 1 (TP1) have been found from rectal swabs of both species (Reuter *et al.*, 2014; Wang *et al.*, 2019). *Desmodus rotundus* parvovirus 1 (DrPV- 1) was detected in vampire bat kidney tissue samples (*Desmodus rotundus*) in Brazil (Souza *et al.*, 2017). In China, porcine parvovirus 7 (PPV7) was obtained from rectal swab samples (Palinski *et al.*, 2016) while in the USA, a viral metagenomic analysis identified a new parvovirus, chaphamaparvovirus, (previously named chapparvovirus; cachavirus) in a diarrhea outbreak in dogs (Fahsbender *et al.*, 2019). The other chaphamaparvovirus, peafowl parvovirus-1 (PePV1) and PePV2, were found in peafowls dying of diarrhea in a wildlife rescue center in China. PePV1 DNA was amplified by PCR from various tissues, including the liver, heart, and lungs and identified by immunohistochemical assay. The results indicated that PePV1 and PePV2 may be fatal pathogens in peafowl (Liu *et al.*, 2020).

Chaphamaparvoviruses have been identified over the past decade (Reuter *et al.*, 2014), with recent epidemiological studies revealing new chaphamaparvoviruses in various animal species. Moreover, chaphamaparvoviruses have recently been detected in humans (Fahsbender *et al.*, 2020), indicating that these viruses have a wider host distribution than is currently estimated.

In Turkey, the presence and distribution of chaphamaparvovirus remains unclear. According to our best information, neither serological nor virological studies have been conducted for chaphamaparvoviruses in humans or animals in Turkey. Accordingly, the present study aimed to investigate the presence and distribution of chaphamaparvovirus in Turkey by applying PCR to rectal swabs collected from both healthy and sick dogs from different geographical regions.

Materials and Methods

Study region and samples: A total of 103 dogs (60 healthy and 43 diarrheal) were included in this study. Rectal swabs were collected from Mersin (the Mediterranean region), Sanliurfa (the South Anatolian region) and Erzurum province (the Eastern Anatolia region) of Turkey for the detection of five viruses (CPV-2, CCoV, CDV, Chaphamaparvovirus, and CAV) that cause diarrhea (2018-2019). In addition, rectal samples were tested for chaphamaparvovirus and others viruses' presence in healthy animals. The 60 healthy dogs included 25 females and 35 males while 19 of the 43 diarrheal dogs were females and 24 were males. The age ranges of the healthy and diarrheal dogs were between 1 month and 9 months and 1.5 months and 8 months, respectively. No animals had a history of vaccination for CCoV, CAV, CPV-2, or CDV. Sixty of the dogs were mixed breeds while the other breeds were cocker (n=10), golden retriever (n=20), kangal (n=5), and German shepherd (n=8).

Hematological and Biochemical Analyses: Hematological and biochemical analyses were performed on whole blood and serum diagnostic samples taken from the dogs with diarrhea. The sera from all infected animals were analyzed for ALT, AST, BUN, cholesterol, total bilirubin and glucose using a biochemistry analyzer (Arkray Spotchem SP-4430 EZ, Japan). Whole blood samples of all the infected animals were analyzed for RBC, hemoglobin, PCV, WBC, neutrophils and lymphocytes using an automatic cell counter (Sysmex PocH-100iV Diff, Japan) (Table 1). Because whole blood and serum samples are routinely taken for the diagnosis and treatment of diarrhea in dogs, samples were not collected specifically for this study.

Sample processing, DNA extraction, and PCR: Rectal swabs from the healthy and diarrheal dogs were moved to appropriate storage containers. Each rectal swab was homogenized with PBS (Phosphate Buffer Saline) and centrifuged at 3,000 rpm for 10 minutes. The supernatant taken into the sterile tube was stored at -80 °C until analysis. 200 µl supernatant was used for nucleic acid extraction. For the extraction of nucleic

acids, High Pure Viral Nucleic Acid kit (Roche Diagnostic, Mannheim, Germany) was used. The obtained nucleic acids were stored at -80 °C. Complementary DNA (cDNA) strands were generated using a High-Capacity cDNA Reverse Transcription kit (Applied Biosystem, CA, USA) according to the manufacturer's instructions. For the detection of chaphamaparvovirus, a nested PCR protocol was used as described elsewhere (Fahsbender *et al.*, 2019) while PCR and RT-PCR protocols were used to detect the viruses (CAV, CCoV, CPV-2 and CDV) causing diarrhea and possible co-infections in dogs (Castilho *et al.*, 2007; Timurkan *et al.*, 2018; Timurkan and Oguzoglu, 2015; Yesilbag *et al.*, 2004) (Table 2). After electrophoresis in 1% agarose gel, PCR products were visualized in a ChemiDoc XRS+ imaging system (Bio-Rad Laboratories, Munich, Germany).

Sequence Analysis: Samples identified as positive by PCR were sent to a commercial firm for sequence analysis (Medsantek, Istanbul, Turkey, <http://www.medsantek.com>). The PCR products were sequenced bi-directionally in an ABI PRISM 3500xL Dx Genetic Analyzer (Thermo Fisher Scientific, Hennigsdorf, Germany). The raw data from the analysis was examined for nucleotide and amino acid sequences using BLASTn and BLASTp from the National Center Biotechnology Information (NCBI) service (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) (Altschul *et al.*, 1990). The data was then analyzed with CLUSTAL W program for nucleotide and amino acid similarities and differences (Hall 1999). MEGA 6 was used to create the maximum likelihood tree using the obtained sequences and other sequences in the GenBank (Tamura *et al.*, 2013).

Table 1 Hematological and sera biochemical results of dogs with CPV-2 and CPV-2 + chaphamaparvovirus.

Parameter	CPV-2 infected dogs (n=28) Mean ± SD	CPV-2 + chaphamaparvovirus infected dogs (n=3) Mean ± SD	Reference range*
RBCs ×10 ⁶ /μL	4.1±3.20	4.0±2.0	4.95-7.87
Hemoglobin g/dl	10±0.23	9.8±13.25	11.9-18.9
PCV %	32±1.22	32±0.12	35-57
WBCs ×10 ³ /μL	3.8±2.54	3.6±2.15	5.0-14.1
Neutrophil ×10 ³ /il	2.5±0.41	2.4±3.15	2.9-12.0
Lymphocyte ×10 ³ /il	0.9±0.21	0.9±0.33	1-4.8
ALT u/l	114.3±3.44	116.3±0.41	10-109
AST u/l	15.0±1.92	14.8±0.36	13-15
BUN mg/dl	33.52±7.53	34.10±3.04	8-28
Cholesterol mg/dl	167.12±15.4	173.12±10.2	135-278
Total bilirubin mg/dl	0.61±1.24	0.63±0.32	0-0.3
Glucose mg/dl	73±0.14	72±21.2	76-119

*International reference range (Tvedten, H., et al., 2004; Merck Manual,2020)

Table 2 Information on the primers used for virus detection in the present study (annealing tempure changes with primer).

Virus name	Gene name	Primer sequence (5'→3')	Product (bp)	PCR conditions	References
Canine coronavirus	M	CCoV1-TCCAGATATGTAATGTTCCGG CCoV2-TCIGTTGAGTAATCACCAGCT	409	95 °C for 4 min; 40 cycles	Yesilbag <i>et al.</i> , 2004
Canine distemper	N	F- 5'-ACTGCTCCCTGATACTGC-3' R- 5'-TTCAACACCAAC(T/C)CCC-3'	480	95 °C for 30 sec, X*° C for 45 sec,	Castilho <i>et al.</i> , 2007
Canine parvovirus-2	VP2	Hfor - CAGGTG ATGAATTTGCTACA Hrev - CATTGGATAAACTGGTGGT	610	72 °C for 1 min;	Timurkan and Oguzoglu, 2015
Canine Chaphamaparvo virus	NS1	CPV-625F- CAACTAGCCGAATGCAGGGA	323	min;	Fahsbender <i>et al.</i> , 2019
		CPV- 948R - CGATAACATCCCCGGACTGG CPV18-687FN-AGCTCAGTTGGCCCCAGATC CPV- 911RN - AGAGGGATCGCTGGATCTGT	224	final extension at 72 °C for 7 min	
Canine Adenovirus	E3	HA1- CGCGCTGAACATTACTACCTTGT HA2-CCTAGAGCACTTCGTGTCGCTT	For CAV2 1030 For CAV1 508		Timurkan <i>et al.</i> , 2018

Results

Rectal swab samples from 103 dogs (60 healthy and 43 diarrheal) were investigated for chaphamaparvovirus presence using nested PCR. Chaphamaparvovirus was only detected in 3

unhealthy dog samples. These dogs were aged 3, 5 and 7 months. One was female and the others were male.

The rectal samples were also tested using PCR and RT-PCR to identify four other viruses (CCoV, CAV, CPV-2, and CDV) and possible co-infections causing diarrhea. The PCR analysis demonstrated that 28 of the

43 diarrheal dogs were positive for CPV-2. However, these 43 dogs were negative for CCoV, CDV and CAV (Table 3). Moreover, the healthy dogs were negative for the remaining viruses. The negative samples were investigated no further for etiological factors causing diarrhea. The PCR analysis also showed that 3 dogs were positive for both chaphamaparvovirus and CPV-2.

The sequences of the PCR amplicons of the NS1 partial gene region of chaphamaparvovirus were compared phylogenetically with the American strains. This demonstrated that the Turkish strains (accession numbers; MT076059, MT076060 and MT076061) were located in the same branch as the American strains (accession numbers MK448316 and MK893826) within the mammalian group (Figure 2). The nucleotide sequences of the three chaphamaparvovirus positive samples were 99.5-100% identical with each other and 98.6% identical with the American strains.

Partial amino acid sequences of the NS1 protein were investigated to detect differences between the Turkish and American strains. As shown in Figure 1, the amino acid sequence comparison indicated that the TR-Erz- chaphamaparvovirus strain had a substitution

at 743 (Ser) whereas the TR-Mers- chaphamaparvovirus strain and the TR-Urf- chaphamaparvovirus had one at 743 (Gly). There were also substitutions at 688 (Ser→Leu), 716 (Asp→His) and 743 (Cys→Ser/Gly) compared to the American strains (Figure 1) (Reference gene number, MK448316, using amino acid comparisons).

Hematological and biochemical analyses were performed on whole blood and serum diagnostic samples from the diarrheal dogs. Firstly, both the dogs with CPV-2 and CPV-2 + chaphamaparvovirus had reduced RBC, HB and PCV values compared to the reference values. This may explain the anemia detected in these patients. Secondly, dogs with CPV-2 and CPV-2 + chaphamaparvovirus also had significantly lower WBC, neutrophil and lymphocyte values than the reference values. Thirdly, dogs with CPV-2 and CPV-2 + chaphamaparvovirus had greater ALT and total bilirubin levels than the reference values. Finally, the animals with diarrhea were hypoglycemic, although their AST and cholesterol values were within the reference range (Merck Manual 2020; Tvedten *et al.*, 2004) (Table 1).

Table 3 Result of viruses screened in this study and samples collected from provinces of Turkey (n= number).

	Regions of Turkey (province)		
	Mersin	Sanliurfa	Erzurum
Diarrhea (dogs)	n=15	n=11	n=17
Canine Parvovirus-2	Positive (n=10; 66,6%)	Positive (n=8; 72,7%)	Positive (n=10; 58,8%)
Canine Chaphamaparvovirus	Positive (n=1; 6,6%)	Positive (n=1; 9%)	Positive (n=1; 5,8%)
Canine Coronavirus	Negative	Negative	Negative
Canine Distemper virus	Negative	Negative	Negative
Canine Adenovirus	Negative	Negative	Negative




Figure 1 Amino acid changes in NS1 partial protein of chaphamaparvovirus.

Discussion

The global importance of emerging and re-emerging infections has increased recently due to their potential effects on human and animal health. Infections may be caused by new (emerging) viruses (Zhou *et al.*, 2020) or mutations of circulating viruses (re-emerging) (Polat *et al.*, 2019; Timurkan and Aydin, 2019). Thus, regular epidemiological investigation and identification of viruses are essential to determine replication mechanisms, host ranges and geographical distributions. Until recently, information about

whether or not chaphamaparvoviruses infect vertebrate hosts was uncertain. However, several recent studies have revealed infections of vertebrate hosts. One study demonstrated that MKPV causes kidney disease in laboratory mice (Roediger *et al.*, 2018) while the detection of DrPV-1 DNA in the kidneys of vampire bats caught in Brazilian forest areas indicates this virus may adapt to kidney niche of vampire bats (Souza *et al.*, 2017). Overall, these studies suggest that chaphamaparvoviruses can be pathogenic for various animal species. Specifically, some mammal chaphamaparvoviruses may be nephrotropic and

cause kidney disease in animals exposed to stress (Lee et al., 2020).

A previous study reported chaphamaparvovirus in 4% of dogs with diarrhea, although there was no information about their ages. While this indicates that chaphamaparvovirus causes infections in dogs, the low viral load suggests that chaphamaparvovirus may result from ingesting contaminated foods rather than infection of intestinal tissue (Fahsbender et al., 2019). Thus, it is necessary to investigate whether chaphamaparvovirus can be morbigenous for dogs. There are no previous reports of chaphamaparvovirus presence in Turkey. In the present study, rectal swabs of both healthy and diahrreal dogs were collected from three different geographical regions, including Mersin (the Mediterranean region), Sanliurfa (the South Anatolian region) and Erzurum province (the Eastern Anatolia region) of Turkey. These were screened for the partial-length NS1 gene of chaphamaparvovirus by nested PCR and were also tested for the presence of four other viruses (CCov, CDV, CAV, and CPV-2). Unsurprisingly CPV-2 was detected in the majority of

dogs with diarrhea (28/43; 65%). Previous studies suggest that CPV-2 is widely dispersed in Turkey's dog population (Dincer 2017; Karapınar et al., 2018; Timurkan and Oguzoglu, 2015). Moreover, the CPV-2c genotype has recently been detected in dogs with gastrointestinal symptoms (Polat et al., 2019). In the present study, chaphamaparvovirus was also detected in CPV-2 positive dogs across a wide range, including Mersin (1/15), Erzurum (1/17) and SanlıUrfa (1/11). All chaphamaparvovirus-positive dogs were under one year old. Enteric viruses that cause diarrhea are known to infect susceptible animals together. For example, rotavirus and CPV-2 have been detected in dogs with gastroenteritis in Mexico (Ortega et al., 2017) while co-infections with turkey astrovirus (TAsTV) and turkey coronavirus (TCov) cause more severe morbidity and higher mortality than single infections (Yu et al., 2000). In our study, the co-detection of CPV-2 and chaphamaparvovirus in three dogs with diarrhea indicates that these viruses may circulate together and increase the severity of the disease as in the aforementioned viruses.




Figure 2 Phylogenetic tree based on the partial NS1 sequence (225 nucleotides) of chaphamaparvoviruses. The phylogeny was performed with maximum likelihood analysis based on 1000 replicates by MEGA 6 software. In this study, chaphamaparvovirus sequences are indicated with a 'circle'.

Recent studies have demonstrated that chaphamaparvoviruses infect different organs, such as the kidneys, heart and lungs (Liu et al., 2020; Roediger et al., 2018). Thus, it is unsurprising that chaphamaparvovirus is detected in organs (especially

kidney tissue) other than the intestines. Although all the chaphamaparvoviruses detected in our study were in dogs with diarrhea, recent studies show that chaphamaparvoviruses are found in different organs. Thus, this should be investigated in future

experimental studies of chaphamaparvoviruses. The presence of chaphamaparvovirus in three major regions in Turkey shows the need for large-scale screening followed by detailed epidemiological observation. Since our study was epidemiological, we did not gather information about pathogenesis in these dogs. Nevertheless, we demonstrated that all the detected chaphamaparvovirus strains were found in dogs with diarrhea; hence, they may be associated with gastrointestinal pathogenesis. No significant biochemical or hematological differences were found between CPV-2 positive and CPV-2 + chaphamaparvovirus animals. The absence of a significant difference in serum and whole blood values between these two groups is not sufficient to evaluate possible infection caused by chaphamaparvovirus in dogs. Future studies of more animals and exclusively on dogs with diarrhea caused by chaphamaparvovirus can help to elucidate clinical presentation in dogs.

In the family parvoviridae, the NS1 protein plays a major role in various processes, including viral cytotoxicity and pathogenicity in host cells. Analysis of the NS1 protein in CPV-2 isolates identified several amino acid changes although information about the functions of amino acids on the NS1 protein is limited (Mira *et al.*, 2020). The detected changes may be essential in virus replication and host distribution. We detected amino acid changes in the NS1 protein of chaphamaparvovirus in specimens obtained from dogs in Turkey. These amino acid changes reflected differences between our Turkey strains and the American strains, which indicates that NS1 protein mutations may occur frequently. Thus, more studies are needed to determine the effects of these amino acid changes on chaphamaparvovirus replication and pathogenesis. Chaphamaparvovirus has been very recently reported in human plasma samples in Brazil (Fahsbender *et al.*, 2020). The present study showed that phylogenetically comparisons of partial NS1 gene sequences indicated an identity of 44% between all isolated chaphamaparvovirus (American and Turkish strains) and human chaphamaparvovirus. That is, chaphamaparvoviruses are probably species specific while chaphamaparvoviruses and human chaphamaparvovirus have large genetic distances.

In conclusion, this study provides the first report of chaphamaparvovirus from the dog population in Turkey and also assesses co-circulation of CPV-2 and chaphamaparvovirus for the first time in dogs showing clinical signs of diarrhea. Despite its DNA structure, chaphamaparvovirus may be evolving through amino acid mutations in the NS1 protein gene. Further epidemiological studies are needed to determine the real incidence of the chaphamaparvovirus in dogs in Turkey. Moreover, molecular and serological studies using more samples will help to better understand chaphamaparvovirus's prevalence and kidney cell tropism, identify specific antibodies and determine its replication mechanisms. Finally, new data will help describe chaphamaparvovirus-dog interaction.

Compliance with ethical standards

Author's contributions: All authors contributed equally to this work.

Conflict of interest: The authors declare no conflict of interest.

Ethical approval: This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol.* 215:403-410.
- Atitwa SC 2018. "How Many Dogs Are There In The World?" [Online]. Available: <https://worldatlas.com/articles/how-many-dogs-are-there-in-the-world.html> Accessed April 20, 2020).
- Aydin H, Kirbas A, Timurkan MO, Aktas MS, Acar-Kirmizi G and Aydin O. 2018. Serological evidence of canine influenza virus infection in shelter dogs in Turkey. *Med Weter* 74:791-794.
- Caddy SL. 2018. New viruses associated with canine gastroenteritis. *Veterinary Journal* 232:57-64.
- Calderon MG, Mattion N, Bucafusco D, Fogel F, Remorini P and La Torre J. 2009. Molecular characterization of canine parvovirus strains in Argentina: Detection of the pathogenic variant CPV-2c in vaccinated dogs. *Journal of Virological Methods* 159(2):141-145.
- Castilho JG, Brandão PE, Carnieli Jr P, Oliveira RN, Macedo CL, Peixoto ZMP, Carrieri ML and Kotait I. 2007. Molecular analysis of the N gene of caninedistemper virus in dogs in Brazil. *Arq Bras Med Vet Zootec* 59:654-659.
- Cave NJ, Marks SL, Kass PH, Melli AC and Brophy MA. 2002. Evaluation of a routine diagnostic fecal panel for dogs with diarrhea. *J Am Veterinary Medical Assoc* 221:52-59.
- Cotmore SF, Agbandje-McKenna M, Chiorini JA, Mukha DV, Pintel DJ, Qiu J, Soderlund-Venermo M, Tattersall P, Tijssen P, Gatherer D and Davison AJ. 2014. The family Parvoviridae. *Archives of Virology* 159:1239-1247.
- De Benedictis, Schultz-Cherry S, Burnham A and Cattoli G. 2011. Astrovirus infections in humans and animals - molecular biology, genetic diversity, and interspecies transmissions. *Infection Genetic and Evolution* 11(7):1529-1544.
- Dincer E. 2017. Molecular characterization and phylogenetic analysis of canine parvovirus 2 in dogs, Mersin Province, Turkey. *Etlik Veteriner Mikrobiyoloji Dergisi* 28(2):96-100.
- Fahsbender E, Charlys da-Costa A, Elise Gill D, Augusto de Padua Milagres F, Brustulin R, Julio Costa Monteiro F, Octavio da Silva Rego M, Soares D'Athaide Ribeiro E, Sabino EC and Delwart E. 2020. Plasma virome of 781 Brazilians with unexplained symptoms of arbovirus infection include a novel parvovirus and densovirus. *PLoS One* 15(3), e0229993.
- Fahsbender E, Altan E, Seguin M, Young P, Estrada M, Leutenegger C and Delwart E. 2019. Chapparvovirus DNA Found in 4% of Dogs with Diarrhea. *Viruses* 11(5):398.
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program

- for Windows 95/98/NT. Nucleic Acids Symposium Series 41:95-98.
- Karapınar Z, Dincer E and Ozkan C. 2018. The investigation and phylogenetic analysis of canine parvovirus 2 infection from blood and rectal swab samples from dogs in Van Province, Turkey. *Van Veterinary Journal* 29(2):83-86.
- Lee Q, Padula MP, Pinello N, Williams SH, O'Rourke MB, Fumagalli MJ, Orkin JD, Song R, Shaban B, Brenner O, Pimanda JE, Weninger W, Marciel de Souza W, Melin AD, Wong J, Crim MJ, Monette S, Roediger B and Jolly CJ. 2020. Murine and related chapparvoviruses are nephro-tropic and produce novel accessory proteins in infected kidneys. *PLoS Pathology* 16(1):e1008262.
- Liu X, Wang H, Liu X, Li Y, Chen J, Zhang J, Wang X, Shen S, Wnag H, Deng F, Wang M, Guan, W and Hu Z. 2020. Genomic and transcriptional analyses of novel parvoviruses identified from dead peafowl. *Virology* 539:80-91.
- Merck Manual (2020) Veterinary Manual. <https://www.merckvetmanual.com/special-subjects/reference-guides>. (accessed april 2020).
- Mira F, Canuti M, Purpari G, Cannella V, Di Bella S, Occhiogrosso L, Schiro G, Chiaramonte G, Barreca S, Pisano P, Lastra A, Decaro N and Guercio A. 2020. Molecular Characterization and Evolutionary Analyses of Carnivore Protoparvovirus 1 NS1 Gene. *Viruses* 11(4):308.
- Miranda C, Thompson G and 2016. Canine parvovirus: The world occurrence of antigenic variants. *Journal of General Virology* 97(9):2043-2057.
- Ortega AF, Martínez-Castañeda JS, Bautista-Gómez LG, Muñoz RF and Hernández IQ. 2017. Identification of co-infection by rotavirus and parvovirus in dogs with gastroenteritis in Mexico. *Brazilian Journal of Microbiololgy* 48(4):769-773.
- Palinski RM, Mitra N and Hause BM. 2016. Discovery of a novel Parvovirinea Virus, porcine parvovirus 7, by metagenomic sequencing of porcine rectal swabs. *Virus Genes* 52(4):564-567.
- Phan TG, Gulland F, Simeone, C, Deng X and Delwart E. 2015. Sesavirus: prototype of a new parvovirus genus in feces of a sea lion. *Virus Genes* 50:134-136.
- Polat PF, Şahan A, Aksoy G, Timurkan MO and Dinçer E. 2019. Molecular and restriction fragment length polymorphism analysis of canine parvovirus 2 (CPV-2) in dogs in southeast Anatolia, Turkey. *Onderstepoort Journal of Veterinary Research* 86(1): 1-8.
- Reuter G, Boros Á, Delwart E and Pankovics P. 2014., Novel circular single-stranded DNA virus from turkey faeces. *Archives of Virology* 159(8), 2161-2164.
- Roediger B, Lee Q, Tikoo S, Cobbin JC, Henderson JM, Jormakka M and Wynne M. 2018. An Atypical Parvovirus Drives Chronic Tubulointerstitial Nephropathy and Kidney Fibrosis. *Cell* 175(2):530-543.
- Souza WM, Romeiro MF, Fumagalli MJ, Modha S, de Araujo J, Queiroz LH and Gifford RJ. 2017. Chapparvoviruses occur in at least three vertebrate classes and have a broad biogeographic distribution. *Journal of General Virology* 98(2):225-229.
- Tamura K, Stecher G, Peterson D, Filipski A and Kumar S. 2013. 'EGAA6: molecular evolutionary genetics analysis version 6.0. *Molecular Biology Evolution* 30:2725-2729.
- Thaiwong T, Sledge DG, Wise A. G, Olstad K, Maes RK and Kiupel M. 2018. Malignant transformation of canine oral papillomavirus (CPV1)-associated papillomas in dogs: An emerging concern?. *Papillomavirus Research* 6: 83-89.
- Timurkan MO, Aydin H and Alkan F. 2018. Detection and molecular characterization of canine adenovirus type 2 (CAV-2) in dogs with respiratory tract symptoms in shelters in Turkey. *Veterinarski arhiv* 88(4):467-479.
- Timurkan MO and Aydın H. 2019. Increased genetic diversity of BVDV strains circulating in Eastern Anatolia, Turkey: first detection of BVDV-3 in Turkey. *Tropical Animal Health Prodduction* 51(7): 1953-1961.
- Timurkan MO and Oğuzoğlu TC. 2015. Molecular characterization of canine parvovirus (CPV) infection in dogs in Turkey. *Veterinaria Italiana* 51(1): 39-44.
- Tvedten H. 2004. Appendix II. In: *Small Animal Clinical Diagnosis by Laboratory Methods*. Eds, Willard, M.D. and H. Tvedten, 4 edn, Saunders, USA, pp: 417-419.
- Wang Y, Yang S, Liu D, Zhou C, Li W, Lin Y and Zong M. 2019. The fecal virome of red-crowned cranes. *Archives of Virology* 64(1):3-16.
- Yesilbag K, Yilmaz Z, Torun S and Pratelli A. 2004. Canine coronavirus infection in Turkish dog population. *Journal Vet Med B Infect Dis Vet Public Health* 51(7):353-355.
- Yu M, Tang Y, Guo M, Zhang Q and Saif YM. 2000. Characterization of a small round virus associated with the poult enteritis and mortality syndrome. *Avian Diseases* 44:600-610.
- Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W and Chen HD. 2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579(7798):270-273.