

6-1-2021

Radiographic and ultrasonographic appearances of chronic tenosynovitis of the lateral extensor tendon at the right tarsus in an Argentinian polo pony

Metha Chanda

Ananya Pongpradit

Wutthiwong Theerapan

Attapon Kamlangdee

Chanikarn Puangthong

See next page for additional authors

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Chanda, Metha; Pongpradit, Ananya; Theerapan, Wutthiwong; Kamlangdee, Attapon; Puangthong, Chanikarn; and Jarutummasiri, Thapana (2021) "Radiographic and ultrasonographic appearances of chronic tenosynovitis of the lateral extensor tendon at the right tarsus in an Argentinian polo pony," *The Thai Journal of Veterinary Medicine*: Vol. 51: Iss. 2, Article 24.

Available at: <https://digital.car.chula.ac.th/tjvm/vol51/iss2/24>

This Report is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Radiographic and ultrasonographic appearances of chronic tenosynovitis of the lateral extensor tendon at the right tarsus in an Argentinian polo pony

Authors

Metha Chanda, Ananya Pongpradit, Wutthiwong Theerapan, Attapon Kamlangdee, Chanikarn Puangthong, and Thapana Jarutummasiri

Radiographic and ultrasonographic appearances of chronic tenosynovitis of the lateral extensor tendon at the right tarsus in an Argentinian polo pony

Metha Chanda^{1*} Ananya Pongpradit² Wutthiwong Theerapan^{2,3}

Attapon Kamlangdee⁴ Chanikarn Puangthong⁵ Thapana Jarutummasiri⁶

Abstract

A 19-year-old polo pony presented with mild lameness alongside a lateral protuberance of the right tarsus. Radiographic findings revealed various defects, including an osteochondral lesion of the medial trochlear ridge of the talus, a sclerotic lesion of the tarsometatarsal joint, a radiolucent area on the distal border of the fourth tarsal bone and an enthesophyte at the proximolateral aspect of the metatarsal bone. Ultrasonography demonstrated the interesting finding of synovial sheath thickening of the lateral extensor tendon with synovial fluid accumulation. In addition, the presence of neutrophils and a granular mucinous background, according to cytological analysis of the fluid, indicated inflammation of the synovial sheath of the tendon. All evidence suggested that the pony was likely suffering from degenerative joint disease and chronic tenosynovitis of the lateral extensor tendon at the right tarsus. The imaging analyses benefited the tarsal defect investigation and proper disease management.

Keywords: degenerative joint disease, diagnostic imaging, polo pony, tenosynovitis, tarsus

¹Department of Large Animal and Wildlife Clinical Science, Faculty of Veterinary Medicine, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

²Kasetsart Veterinary Imaging and Radiotherapy Center, Kasetsart University Veterinary Teaching Hospital, Bang Khen Campus, Bangkok 10900, Thailand

³Department of Companion Animals Clinical Sciences, Faculty of Veterinary Medicine, Kasetsart University, Bang Khen Campus, Bangkok 10900, Thailand

⁴Department of Pathology, Faculty of Veterinary Medicine, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

⁵Veterinary Clinical Studies Program, Faculty of Veterinary Medicine, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

⁶TP Equine Ambulatory Service, Bangkok 10240, Thailand

*Correspondence: foetmtcd@ku.ac.th (M. Chanda)

Received: January 13, 2021.

Accepted: February 18, 2021.

doi: <https://doi.org/10.14456/tjvm.2021.49>

Introduction

The tarsus is a complex apparatus of multiple bones, soft tissue and joint structures susceptible to extensive injuries (Clegg, 2003). It acts relatively to stifle, fetlock and coffin joints as a hindlimb shock absorber during the stance phase of motion (Back *et al.*, 1995). Hence, tarsus defects are primarily found in sporting horses engaging in high-intensity exercise such as galloping, reining and jumping (Gabel, 1980). Given the complicated structure of the tarsus and the enormous strain of strenuous exercise, tarsus injuries are the most likely cause of hindlimb lameness in the horse (Axelsson *et al.*, 1998; Kinns and Nelson, 2010).

Various types of tarsus damage, including degenerative joint disease (DJD), osteochondrosis, disease of the synovial structure and tendon and ligament injuries, can lead to hindlimb lameness (Blaik *et al.*, 2000). Amongst the structural defects in the tarsus, tarsal DJD is considered to be the primary cause of hindlimb lameness (Vaughan, 1965). It is characterized by the presence of periarticular osteophytes, subchondral bone lysis or sclerosis and ankylosis (Björnsdóttir *et al.*, 2000; Butler, 2000) and has been reported in several riding populations, including the Swedish Icelandic horse (Axelsson *et al.*, 1998;

Eksell *et al.*, 1998) and the Thoroughbred and American Quarter horse (Taintor *et al.*, 2014).

Tenosynovitis is an inflammation of the synovial membrane and fibrous layer of the tendon sheath, indicated by synovial effusion in the injured area (McIlwraith, 1987). The most common sites of tenosynovitis are the tarsal sheath, digital flexor tendon sheath and extensor tendon sheath of the carpus (McIlwraith, 1987). Peritendinous strain and irritation between the visceral layer and the outer surface of the tendon itself also cause tenosynovitis (Wallace, 1972). This case report demonstrates diagnostic imaging of the affected right tarsus and the cytological properties of synovial fluid from the protuberant lesion in an Argentinian polo pony.

Case description

A 19-year-old Argentinian polo pony (gelding) was presented with an enlargement of the lateral aspect of the right tarsus (Figures 1A and 1B). At the initial examination, a blemished protuberance was observed at the lateral aspect of the right tarsus, which was firm and relatively warm but insensitive to compression. Fundamental gait analysis revealed mild lameness on the right hindlimb and a positive response to the hock flexion test.




Figure 1 The blemished protuberance is present on the lateral aspect of the right tarsus in the dorsal (A) and plantar (B) views.

A radiograph of the right tarsus was performed according to the method described previously (Kinns and Nelson, 2010; Vanderperren *et al.*, 2009b). The radiographic imaging illustrated multiple defects: 1) a lateral protuberance on the right tarsus and the formation of an enthesophyte at the proximolateral third metatarsal bone (MT3) (Figure 2A), 2) a bony spur at the medial trochlear ridge of the talus (Figures 2B),

3) a sclerotic lesion and narrowed space of the joint at the osseous margin of the tarsometatarsal joint (Figures 2B, 2D), 4) various radiolucent areas indicating subchondral bone lysis on the ventral surface of the fourth tarsal bone (Figure 2B), and 5) the presence of enthesopathy characterized by a radiolucent area alongside bony ossification at the dorsal aspect of tuber calcanei (Figures 2B and 2C). Ultrasonography was

subsequently performed at the transverse (Figure 3A) and longitudinal planes (Figure 3B) of the enlargement area to assess soft tissue configuration. The ultrasonographic images (Figures 3C-E) revealed an anechoic fluid accumulation within the fibrous thickening of the lateral extensor tendon sheath. Moreover, synovial proliferation was also observed in the inner lining of the tendon sheath. Approximately 2 mL of accumulated fluid was aspirated from the lesion for cytological analysis. The fluid was a reddish colour with low viscosity and moderate turbidity (Figure 4A) with a $0.88 \times 10^9/L$ total nucleated cell count (TNCC) and 40 g/L of total protein. The aspirated fluid also showed numerous erythrocytes, neutrophils and a granular mucinous background (Figure 4B). The

presence of free erythrocytes might indicate blood contamination during sample collection. Nevertheless, elevated neutrophils alongside the increased expression of a granular mucinous background would suggest inflammation of the synovial tendon sheath. Regarding the case history and clinical examination, the polo pony was diagnosed with degenerative joint disease (DJD) and chronic lateral digital extensor tendon tenosynovitis of the right tarsus. Initially, pain-relieving gel (cold gel containing *Arnica* extract) and oral anti-inflammatory medication (phenylbutazone 4.4 mg/kg body weight) were administered twice daily for 14 days. Although the protuberance still existed, the lameness of the right hindlimb had improved by the end of the therapeutic period.




Figure 2 The radiographic images demonstrated multiple defects, including 1) soft tissue protuberance at the lateral aspect of the joint (sequential white triangles) (A) and a enthesophyte formation at the lateral aspect of the proximolateral angle of the third metatarsal bone (black-white triangles) (A and B), 2) a bone spur at the trochlea ridge of the talus (yellow triangle) (B), 3) a sclerotic lesion of the osseous margin at the tarsometatarsal joint (white arrows) (B and D), 4) various radiolucent areas indicating subchondral bone lysis on the ventral surface of the fourth tarsal bone (B), and 5) a radiolucent area at the dorsal aspect of tuber calcanei (yellow arrow) (B and C).




Figure 3 The ultrasonography was performed at the transverse (A) and longitudinal planes (B). The transverse (C) and longitudinal (D and E) ultrasonographic images demonstrate the hypoechoic soft tissue swelling and synovial thickening with anechoic effusion (*). The images also show a small loss of normal fibre pattern in the lateral extensor tendon (yellow arrows) (D). Synovial proliferation is seen at the inner lining of the affected tendon sheath (D and E). The width of the thickened synovial tendon sheath at the most distended point was 2.33 cm (yellow vertical dots) (D). PROX = proximal part; DIST = distal part; CAU = caudal part; CRA = cranial part; 1 = tibia; 2 = tibial malleolus; 3 = talus; 4 = lateral extensor tendon; 5 = skin; asterisk (*) = anechoic fluid.




Figure 4 Reddish synovial fluid was aspirated from the lesion (A). A captured image of the cytological examination (100x magnification) shows numerous erythrocytes, neutrophils and granular mucinous background (B).

Discussion

Several abnormalities of the tarsal structure, such as degenerative joint disease (DJD), osteochondrosis (OC) and soft tissue injuries, can be revealed by the appropriate imaging technique (Vanderperren *et al.*, 2009a; Vanderperren *et al.*, 2009b). Radiographic and ultrasonographic modalities are still the primarily selected instruments that provide practical advantages for tarsal defect investigation (Kinns and Nelson, 2010). Nonetheless, some convoluted defects may not be correctly identified by these imaging techniques. Consequently, care must be taken to avoid misinterpretation of the tarsal abnormalities. In this case, numerous structural disorders of the right tarsus were discovered by diagnostic imaging analysis.

The radiographic findings demonstrated bone surface irregularity, narrowing of the joint space, subchondral bone lysis, bony sclerosis, enthesophyte formation and periarticular bone proliferation. These symptoms have been associated with DJD of the tarsus in previous studies (Gough and Munroe, 1998; Verschooten and Schramme, 1994). The predisposing cause of DJD at distal intertarsal and tarsometatarsal joints was thought to accompany an unbalanced force distribution on the joint surface (Björnsdóttir *et al.*, 2004). Apart from DJD, the osteochondrosis (OC), a bony defect attributable to abnormal endochondral ossification, was also seen (Donabédian *et al.*, 2008). Although a bony spur was present at the medial trochlear ridge of the talus, in this case, it did not necessarily develop the OC though it is occasionally visible in some horses (Butler, 2000; Clegg, 2003). Hence, the outcome of this imaging addressed the paramount importance of radiography for evaluating tarsal defects. The superimposition of bony structures and the inability to distinguish the architecture of soft tissues are the main limitations of using this modality (Kraft and Gavin, 2001; Park *et al.*, 1987). Therefore, ultrasonography has been subsequently utilized to depict the sonographic appearance of the ligament and tendon structure surrounding the tarsus (Davis *et al.*, 2014; Dik, 1993). Interestingly, the ultrasonography implemented in this case illustrated anechoic fluid

accumulation, fibrous thickening of the lateral digital extensor tendon sheath with synovial proliferation and evidence of the loss of the typical fibre pattern of the tendon. This evidence is likely to indicate chronic tendinitis with tenosynovitis of the tendon as reported previously (McIlwraith, 1987; Raes *et al.*, 2010). The accumulated fluid was aspirated for pathological examination of the tendon architecture. The aspirated fluid was reddish, with reduced viscosity and high turbidity. Although the TNCC of the fluid ($0.88 \times 10^9/L$) was within the normal range (0.2 to $3.5 \times 10^9/L$) (Steel, 2008), an increase in total protein (> 20 g/L) alongside the presence of high neutrophil number and a granular mucinous background support the diagnosis of tenosynovitis of the lateral digital extensor tendon.

The treatment of tenosynovitis of the tendon sheath can be administered by intrathecal injection or surgical resection depending on the severity and location of the affected tendon sheath (Taintor *et al.*, 2013; Wright and McMahon, 1999). Since an improvement in the lameness after oral administration of the anti-inflammatory drug and the application of the pain-relieving gel, the pony has neither undergone a local injection of anti-inflammatory prescription nor surgical treatment even though the lateral protuberance is still present. However, the intrathecal administrations of short-acting corticosteroid may be alternatively considered if the lameness recurred due to DJD and inflammation of the tendon sheath.

References

- Axelsson M, Eksell P, Ronéus B, Broström H, Häggström J and Carlsten J 1998. Relationship between hind limb lameness and radiographic signs of bone spavin in Icelandic horses in Sweden. *Acta Vet Scand.* 39: 349-357.
- Back W, Schamhardt H, Savelberg H, Van Den Bogert A, Bruin G, Hartman W and Barneveld A 1995. How the horse moves: 2. Significance of graphical representations of equine hind limb kinematics. *Equine Vet. J.* 27: 39-45.

- Björnsdóttir S, Axelsson M, Eksell P, Sigurdsson H and Carlsten J 2000. Radiographic and clinical survey of degenerative joint disease in the distal tarsal joints in Icelandic horses. *Equine Vet J.* 32: 268-272.
- Björnsdóttir S, Ekman S, Eksell P and Lord P 2004. High detail radiography and histology of the centrodistal tarsal joint of Icelandic horses aged 6 months to 6 years. *Equine Vet J.* 36: 5-11.
- Blaik MA, Hanson RR, Kincaid SA, Hathcock JT, Hudson JA and Baird DK 2000. Low-field magnetic resonance imaging of the equine tarsus: normal anatomy. *Vet Radiol Ultrasound.* 41: 131-141.
- Butler JA, Colles CM, Dyson SJ, Kold SE and Poulos PW 2000. The tarsus. In: *Clinical Radiology of the Horse.* Oxford: Blackwell Scientific. 247-284.
- Clegg P 2003. Differential diagnosis of a swollen hock in the horse. *In Practice* 25: 328-341.
- Davis W, Caniglia CJ, Lustgarten M, Blackwelder T, Robertson I and Redding WR 2014. Clinical and diagnostic imaging characteristics of lateral digital flexor tendinitis within the tarsal sheath in four horses. *Vet Radiol Ultrasound.* 55: 166-173.
- Dik KJ 1993. Ultrasonography of the equine tarsus. *Vet Radiol Ultrasound.* 34: 36-43.
- Donabédian M, Van Weeren P, Perona G, Fleurance G, Robert C, Leger S, Bergero D, Lepage O and Maerin-rosset W 2008. Early changes in biomarkers of skeletal metabolism and their association with the occurrence of osteochondrosis (OC) in the horse. *Equine Vet J.* 40: 253-259.
- Eksell P, Axelsson M, Broström H, Ronéus B, Häggström J and Carlsten J 1998. Prevalence and risk factors of bone spavin in Icelandic horses in Sweden: a radiographic field study. *Acta Vet Scand.* 39: 339-348.
- Gabel AA 1980. Lameness caused by inflammation in the distal hock. *Vet Clin North Am Large Anim Pract.* 2: 101-124.
- Gough M and Munroe G 1998. Decision making in the diagnosis and management of bone spavin in horses. *In Practice.* 20: 252-259.
- Kinns J and Nelson N 2010. Imaging tarsal trauma. *Equine Vet Educ.* 22: 296-298.
- Kraft SL and Gavin P 2001. Physical principles and technical considerations for equine computed tomography and magnetic resonance imaging. *Vet Clin North Am Equine Pract.* 17: 115-130.
- McIlwraith CW 1987. Disease of Joints, Tendons, Ligaments, and Related Structures. In: *Adams' Lameness in Horses.* 4th ed. Stashak, T.S. (ed). Pennsylvania: Lippincott Williams & Wilkins. 475-479.
- Park RD, Nelson TR and Hoopes PJ 1987. Magnetic resonance imaging of the normal equine digit and metacarpophalangeal joint. *Vet Radiol Ultrasound.* 28: 105-116.
- Raes EV, Vanderperren K, Pille F and Saunders JH 2010. Ultrasonographic findings in 100 horses with tarsal region disorders. *Vet J.* 186: 201-209.
- Steel CM 2008. Equine synovial fluid analysis. *Vet Clin North Am Equine Pract.* 24: 437-454.
- Taintor J, Caldwell F and Almond G 2013. Aseptic tenosynovitis of the carpal flexor sheath caused by rupture of the accessory ligament of the deep digital flexor tendon. *Can Vet J.* 54: 765.
- Taintor JS, Wright J, Caldwell F, Dymond B and Schumacher J 2014. Efficacy of an extract of blue-green algae in the amelioration of lameness caused by degenerative joint disease in the horse. *J Equine Vet Sci.* 34: 1197-1200.
- Vanderperren K, Raes E, Hoegaerts M and Saunders JH 2009^a. Diagnostic imaging of the equine tarsal region using radiography and ultrasonography. Part 1: The soft tissues. *Vet J.* 179: 179-187.
- Vanderperren K, Raes E, Van Bree H and Saunders JH 2009^b. Diagnostic imaging of the equine tarsal region using radiography and ultrasonography. Part 2: Bony disorders. *Vet J.* 179: 188-196.
- Vaughan J 1965. Analysis of lameness in the pelvic limb and selected cases, *Proc Am Ass equine Practnrs.* 223-241.
- Verschooten F and Schramme M 1994. Radiological examination of the tarsus [in horses]. *Equine Vet Educ.* 6: 323-332.
- Wallace CE 1972. Chronic tendosynovitis of the extensor carpi radialis tendon in the horse. *Aust Vet J.* 48: 585-587.
- Wright I and McMahon P 1999. Tenosynovitis associated with longitudinal tears of the digital flexor tendons in horses: a report of 20 cases. *Equine Vet J.* 31: 12-18.