

6-1-2022

Immunochromatographic test (ICT) for the detection of antibody against ALV-J

Guihua Wang

Yanming Wang

Ziqiang Cheng

Follow this and additional works at: <https://digital.car.chula.ac.th/tjvm>



Part of the [Veterinary Medicine Commons](#)

Recommended Citation

Wang, Guihua; Wang, Yanming; and Cheng, Ziqiang (2022) "Immunochromatographic test (ICT) for the detection of antibody against ALV-J," *The Thai Journal of Veterinary Medicine*: Vol. 52: Iss. 2, Article 17. Available at: <https://digital.car.chula.ac.th/tjvm/vol52/iss2/17>

This Short Communication is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in The Thai Journal of Veterinary Medicine by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Immunochromatographic test (ICT) for the detection of antibody against ALV-J

Guihua Wang¹ Yanming Wang¹ Ziqiang Cheng^{1*}

Abstract

ALV-J is an important pathogen in the poultry industry in China. In this study, we developed an immunochromatographic test (ICT) assay for the detection of antibody against of ALV-J to evaluate ALV-J infection. Gp85 protein of ALV-J was employed as a capture antigen. Antigen-antibody reaction was detected by either colloidal-gold conjugated gp85 protein or monoclonal antibody of ALV-J. The high consistency of results in ICT and ELISA indicated that the ICT is reliable. This suggests that the gp85-based ICT can be used as a rapid on-site diagnostic method of ALV-J.

Keywords: immunochromatographic test, antibody, ALV-J, rapid diagnosis

¹Department of Fundamental Veterinary, College of Veterinary Medicine, Shandong Agricultural University, Tai'an, China

*Correspondence: czqsd@126.com (Z.Cheng)

Received December 10, 2021

Accepted February 4, 2022

<https://doi.org/10.14456/tjvm.2022.42>

Introduction

ALV-J is one of the pathogens causing serious harm to the poultry industry in China. Eliminating infected chickens is an effective method to control the spread of ALV-J (Feng and Zhang, 2016). However, coinfection and recombination of ALV-J with other pathogens have been reported (Li *et al.*, 2021; Li *et al.*, 2020; Sun *et al.*, 2020). Therefore, early rapid and accurate detection and diagnosis is very important. Enzyme-linked immunosorbent assay (ELISA) (Yun *et al.*, 2013), indirect fluorescence assay (IFA) and polymerase chain reaction (PCR) (Smith *et al.*, 1998) are usually used in diagnosis. ELISA is a simple, specific but expensive method. IFA is a sensitive and specific method however, it is time consuming. DNA based detection methods are highly sensitive but require specific instruments and trained operators. Thus, a rapid, cheap and simple test is demanded in the detection of clinical ALV-J infection. In this study, a colloidal gold test strip was developed for simple and rapid detection of ALV-J antibody to evaluate ALV-J infection.

Materials and Methods

Protein expression: We compared isolated wild ALV-J strains published on NCBI in the last ten years, and selected 504 bp relatively conservative and higher antigenicity of ALV-J gp85 protein as antigen. The selected gp85 gene sequences were expressed with forward primer (5'-GCGGATCCATCAAGAACGGAA CAACACG-3'), and reverse primer (5'-GCGCGCAAG CTTGTCCCAACAATCAAGAAAATA-3'). A 504 bp target fragment was obtained by PCR amplification and PCR products were transfected into the DH5 α receptor cells. The objective fragment was induced by expression in *Escherichia coli*, then, the thallus was cleaved by ultrasound. The expressed proteins were purified by dialysis. The gp85 antibodies in ALV-J positive serum were used to detect the reactogenicity and specificity of antigen by Western blot.

Preparation of monoclonal antibody against ALV-J: BALB/c mice (6 weeks old) were immunized

subcutaneously by purified protein. The mouse with the highest serum titer was sacrificed and spleen cells were fused with SP2/0 myeloma cells. Positive hybridoma cells were cultured for monoclonal antibody (mAb) production. The animal experiments were approved by the Shandong Agricultural University Animal Care Committee.

Preparation of colloidal gold particles: Colloidal gold particles were prepared as reported (Liu *et al.*, 2012). Briefly, 1% chloroauric acid solution was boiled, mixed with an equal volume of 1% sodium citrate solution and stirred for 10 mins until the color of solution to be stable. Transmission electron microscopy observation revealed that the diameter of colloidal gold particles was about 25 nm.

Preparation of the colloidal gold conjugate: Different amounts of 25 mM K₂CO₃ were used to adjust pH gradient of colloidal gold solution from 7.1 to 7.9. Different concentrations of purified gp85 protein were added to colloidal gold solution (10, 11, 12, 13, 14, 15, 16, 17, 18, 19, and 20 μ g/mL) with gentle stirring at room temperature for 20 mins. The mixture was blocked by different concentrations of BSA (0.5%, 0.8%, 1.0%, 2.0%, and 3.0%) and then centrifuged at 5 000 g for 20 mins. The precipitate was redissolved in 0.01M PBS (pH 7.4) containing 1% BSA and 2.5% sucrose and stored at 4°C.

Preparation of the immunochromatographic strip: The antigens and prepared mAb were diluted to 1.0 mg/mL, 1.5 mg/mL and 2.0 mg/mL with 0.01 M sodium PBS (Ph 7.4). The diluted gp85 protein and mAb were transferred on to the Nitrocellulose membrane to form the test (T) and the control (C) line. Polyester cellulose membrane was soaked in immune colloidal gold solution for 10 minutes and then was dried at 37°C. The absorbent paper and sample pad were cut to an appropriate size for assembling according to Figure 1A. The test strips were stored in a desiccator at room temperature.




Figure 1 The immunochromatographic test strip. (A) Schematic diagram of the immunochromatographic strip. (B) The results of immunochromatographic test.

Sensitivity, specificity and accuracy of the immunochromatographic strip: Serial diluted (1 mg/mL, 100 µg/mL, 10 µg/mL, 1 µg/mL, 100 ng/mL, and 10 ng/mL) ALV-J antibody positive and negative sera were dropped on the sample pad, respectively, to evaluate sensitivity. ALV-J infected chicken serum was used as a positive. Un-infected chicken serum was used as a negative. The specificity was determined using different antibody of Marek's disease virus (MDV), reticuloendotheliosis virus (REV) and avian influenza virus (AIV). To evaluate the accuracy, we compared the detection results of clinical 278 serum samples tested with the test strips and IDEXX commercial ELISA kits.

Results and Discussion

To detect the purified gp85 protein, chicken positive serum which was detected by ALV-J antibody detection kit (IDEXX, America) was used as the primary antibody for gp85, followed by HRP-labeled goat anti-chicken IgG (Bioss, China) as the secondary antibody. Western Blot result showed that the purified protein was capable of binding with gp85 antibody with a good reactivity and specificity (Figure 2A).

Our results demonstrate that the optimal pH of colloidal gold solution was 7.5. The optimal concentration of purified gp85 labelled by colloidal gold was 20 µg/mL. One percent BSA was the optimal concentration of blocking buffer. Following the experimental procedure, the optimal concentrations of coating antigen and antibody were both 2 mg/mL.

ALV-J antibody positive and negative sera were used to determine the validity and sensitivity of the immunochromatographic strip. As shown in Figure 1B, the appearance of both test and control lines was judged to positive results. The negative result was judged by the presence of only the control line. The test was invalid if only the test line appeared or no control line appeared. The antibodies of different antigens were used to determine the specificity of the immunochromatographic strip. We found that the immunochromatographic strip was specific for ALV-J antibody (Figure 2B).

A total of 278 clinical sera samples were detected by the developed test strips and ELISA kit, respectively. The result showed that the positive rate of test strips was 57.6% (Figure 2C) and ELISA was 58.7%. The rate of matching was 98.12%, indicating that the test strip had good specificity, sensitivity and accuracy.




Figure 2 A) The reactivity and specificity of recombinant protein were analyzed by Western blot. (B) Specificity test of immunochromatographic test strip. 1: MDV infection sera sample; 2: REV infection sera sample; 3: AIV infection sera sample; 4: ALV-J infection sera sample; 5: negative sera sample; 6: positive sera sample; (C) Some of the results of clinical sera samples. 1, 2, and 3 from different farms.

ALV-J infection causes enormous economic loss to the poultry industry in China (Qu *et al.*, 2016; Zhou *et al.*, 2019). ELISA, immunofluorescence assays, PCR and virus isolation, which require specialized technologies and equipment, are usually applied to detect ALV-J infection. Immunochromatographic test (ICT) assays are rapid, simple, cheap and have no need of special equipment for detection (Feng *et al.*, 2015), thus they are convenient for primary level and field use.

Conflict of Interest: The authors declare no conflicts of interest.

Acknowledgements

This work was supported by the Key Research and Development Program of Shandong Province (Important Science and Technology Innovation Project) (2019JZZY010735).

References

Feng M, Kong DZ, Wang WB, Liu LQ, Song SS and Xu CL 2015. Development of an Immunochromatographic Strip for Rapid Detection of *Pantoea stewartii* subsp. *stewartii*. *Sensors* (Basel). 15(2): 4291-4301.

- Feng M and Zhang XQ 2016. Immunity to Avian Leukosis Virus: Where Are We Now and What Should We Do? *Front Immunol.* 7:624.
- Li M, Xiong HF, Wu HW, Hu DM, Lin Y, Huang XT, Wang J, Qi KZ and Liu HM 2021. Pathologic Characterization of Coinfection with *Histomonas meleagridis*, Marek's Disease Virus, and Subtype J Avian Leukosis Virus in Chickens. *Avian Dis.* 65(2): 237-240.
- Li QH, Wang K, Li M, Lin LL, Shi MY, Li HJ, Deng QM, Teng H, Mo ML, Wei TC and Wei P 2020. Recombinant subgroup B avian leukosis virus combined with the subgroup J env gene significantly increases its pathogenicity. *Vet Microbiol.* 250:108862.
- Liu X, Xiang JJ, Tang Y, Zhang XL, Fu QQ, Zou JH and Lin L 2012. Colloidal gold nanoparticle probe-based immunochromatographic assay for the rapid detection of chromium ions in water and serum samples. *Anal Chim Acta.* 745:99-105.
- Qu YJ, Liu LT, Niu YJ, Qu Y, Li N, Sun W, Lv CW, Wang PF, Zhang GH and Liu SD 2016. Viral proliferation and expression of tumor-related gene in different chicken embryo fibroblasts infected with different tumorigenic phenotypes of avian leukosis virus subgroup J. *Poultry Sci.* 95(10): 2383-2390.
- Smith EJ, Williams SM and Fadly AM 1998. Detection of avian leukosis virus subgroup J using the polymerase chain reaction. *Avian Dis.* 42(2): 375-380.
- Sun YN, Lu QZ, Zhang JF, Li XX, Zhao JK, Fan WQ, Ji PP, Wang K, Zhou EM and Zhao Q 2020. Coinfection with avian hepatitis E virus and avian leukosis virus subgroup J as the cause of an outbreak of hepatitis and liver hemorrhagic syndromes in a brown layer chicken flock in China. *Poult Sci.* 99(3): 1287-1296.
- Yun BL, Li DL, Zhu HB, Liu W, Qin LT, Liu ZS, Wu G, Wang YQ, Qi XL, Gao HL, Wang XM and Gao YL 2013. Development of an antigen-capture ELISA for the detection of avian leukosis virus p27 antigen. *J Virol Methods.* 187(2): 278-283.
- Zhou DF, Xue JW, Zhang Y, Wang GH, Feng YS, Hu LP, Shang YL and Cheng ZQ 2019. Outbreak of myelocytomatosis caused by mutational avian leukosis virus subgroup J in China, 2018. *Transbound Emerg Dis.* 66(2): 622-626.