

11-1-1994

Loop Electrosurgical Excision Procedure (LEEP) and cervicalintraepithelial neoplasia: A preliminary report

Wichai Termrungruanglert

Wirach Wisawasukmongchol

Kobchitt Limpaphayom

Apichai Vasuratna

Tul Sittisomwong

See next page for additional authors

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Termrungruanglert, Wichai; Wisawasukmongchol, Wirach; Limpaphayom, Kobchitt; Vasuratna, Apichai; Sittisomwong, Tul; Niruttisard, Somchai; and Sirisabya, Nakorn (1994) "Loop Electrosurgical Excision Procedure (LEEP) and cervicalintraepithelial neoplasia: A preliminary report," *Chulalongkorn Medical Journal*: Vol. 38: Iss. 11, Article 4.

Available at: <https://digital.car.chula.ac.th/clmjjournal/vol38/iss11/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Loop Electrosurgical Excision Procedure (LEEP) and cervical intraepithelial neoplasia: A preliminary report

Authors

Wichai Termrungruangleert, Wirach Wisawasukmongchol, Kobchitt Limpaphayom, Apichai Vasuratna, Tul Sittisomwong, Somchai Niruttisard, and Nakorn Sirisabya

นิพนธ์ต้นฉบับ

Loop Electrosurgical Excision Procedure (LEEP) and cervical intraepithelial neoplasia: A preliminary report.

Wichai Termrungruenglert*

Wirach Wisawasukmongchol* Kobchitt Limpaphayom *

Apichai Vasuratna* Tul Sittisomwong*

Somchai Niruttisard* Nakorn Sirisabya*

Termrungruenglert W, Wisawasukmongchol W, Limpaphayom K, Vasuratna A, Sittisomwong T, Niruttisard S, Sirisabya N. Loop Electrosurgical Excision Procedure (LEEP) and cervical intraepithelial neoplasia: A preliminary report. Chula Med J 1994 Nov; 38(11): 653-658

Objective: *To present our early experiences with the loop electrosurgical excision procedure in patients with cervical intraepithelial neoplasia (CIN).*

Design: *Prospective Study.*

Methods: *Thirty one patients who had pap smear class 3-5 and colposcopically confirmed cases of CIN lesions were subjected to LEEP. The procedures were undergone mainly under local anesthesia.*

Results: *The majority of the cases (83.9%), perioperative and postoperative hemorrhage was minimal. No patient needed blood transfusion or subsequent admission to the hospital due to late hemorrhage. Two cases experienced minimal late hemorrhage and responded to AgNO₃ application. The operation time was less than 5 minutes in 22 cases (71.0%). In 28 of the 31 cases (90.3%), this procedure was sufficient for complete treatment.*

Conclusion: *Loop electrosurgical excision procedure tends to be a reliable, well-tolerated, economical and efficient technique for the management of cervical intraepithelial neoplasia with a low incidence of short-term morbidity.*

Key words : *Loop electrosurgical excision procedure, Cervical intraepithelial neoplasia.*

Reprint request : Termrungruenglert W, Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. October 15, 1994.

วิจัย เติมรุ่งเรืองเลิศ, วิรัช วิศวสุขมงคล, กอบจิตต์ ลิ้มพยอม, อภิชัย วสุรัตน์, ดุลย์ สิทธิสมวงศ์, สมชัย นิรุตติศาสตร์, นคร ศิริทรัพย์. การผ่าตัดปากมดลูกด้วยห่วงลวดไฟฟ้าในการรักษาภาวะปากมดลูกมีเซลล์ผิดปกติก่อนเป็นมะเร็งระยะลุกลาม : รายงานเบื้องต้น จุฬาลงกรณ์เวชสาร 2537 พฤศจิกายน 38(11): 653-658

วัตถุประสงค์ เพื่อศึกษาผลการผ่าตัดปากมดลูกด้วยห่วงลวดไฟฟ้า (Loop electrosurgical excision procedure : LEEP) ในการรักษาผู้ป่วยภาวะ cervical intraepithelial neoplasia (CIN) ซึ่งเป็นวิธีการรักษาวิธีใหม่ในเบื้องต้น

รูปแบบการวิจัย การศึกษาไปข้างหน้า

วิธีการวิจัย ผู้ป่วยหญิง 31 คน ซึ่งตรวจ Pap smear เป็นชั้น 3-5 และได้รับการตรวจยืนยันโดยการตัดชิ้นเนื้อปากมดลูกโดยอาศัยกล้อง colposcope เป็น CIN ได้รับการผ่าตัดปากมดลูกด้วยห่วงลวดไฟฟ้า โดยการนัดยาเฉพาะที่

ผลการศึกษา ผู้ป่วยร้อยละ 83.9 มีการเสียเลือดระหว่างและหลังผ่าตัดน้อย (น้อยกว่า 10 มล) ไม่มีผู้ป่วยรายใดเสียเลือดมากจนต้องให้เลือด พบผู้ป่วย 2 รายใน 31 ราย มีเลือดออกเล็กน้อยภายหลังผ่าตัด ซึ่งสามารถหยุดได้โดยการจี้ซิลเวอร์ในเตรด ร้อยละ 71.0 ใช้เวลาในการผ่าตัดน้อยกว่า 5 นาที สามารถให้การรักษาภาวะดังกล่าวโดยการผ่าตัดปากมดลูกด้วยวิธีนี้อย่างเดียว ร้อยละ 90.3

สรุป การผ่าตัดปากมดลูกด้วยห่วงลวดไฟฟ้าเป็นวิธีที่มีประสิทธิภาพในการรักษาภาวะ cervical intraepithelial neoplasia โดยมีภาวะแทรกซ้อนจากการผ่าตัดน้อยในการติดตามเบื้องต้น

The prevalence of cervical intraepithelial neoplasia (CIN) has been increasing over the last 2 decades.⁽¹⁾ Despite the advent of ablative methods of treatment, a large proportion of patients for colposcopic assessment are admitted to the hospital for cold knife conization of the cervix under general or regional anesthesia. Cold knife conization, however, is associated with considerable morbidity, such as hemorrhage, which occasionally requires resuturing of the cervix or even hysterectomy and blood transfusion.^(2,3)

The development of invasive cervical cancer after ablative treatment for precancer^(4,5) has prompted the increased use of excisional methods of treatment. As an alternative to ablative treatment under local anesthesia, laser excision conization of the cervix has been advocated.^(6,7) Recently, several investigators have reported electrosurgical excision of CIN lesions and typical transformation zones using thin wire loop electrodes of either large or small diameter as a safe and highly effective treatment method.⁽⁸⁻¹⁷⁾ Electrosurgical excision of CIN using wire loop electrodes has been referred to by various terms. Cartier⁽⁹⁾ introduced the use of "the diathermy loop excision". Prendiville and Cullimore,⁽¹⁰⁾ who used loops of a larger diameter than did Cartier, referred to their modification of the basic technique as "large loop excision of the transformation zone" (LLETZ). Wright et al⁽¹¹⁾ used the term "loop electrosurgical excision procedure (LEEP)" for these procedures, regardless of the size of the loop electrode used.

The loop electrosurgical excision procedure represents an advanced ability to conservatively manage patients with CIN, as affected tissue is removed in its entirety rather than ablated. The excised tissues are available for histopathologic analysis, allowing both diagnosis

and treatment of selected patients at a single setting and ensuring that colposcopically inapparent invasive or microinvasive cancers are not inadvertently treated by a conservative ablative procedure. In this study, we report our clinical experience using LEEP in the management of CIN.

Materials and Methods

From December 1993 to June 1994, the patients were referred because of Papanicolaou smear class 3-5, and they underwent a standard colposcopic evaluation with site-directed biopsies. All woman included in this study had histologically diagnosed CIN lesions. The entire endocervical limits could be visualized colposcopically.

The subjects were placed in the lithotomy position and a large speculum with suction tube attached was used to expose the cervix. A paracervical block was performed using 3-5 ml. of 2% Xylocaine in each side and intracervically infiltrated 1-2 mm. beneath the cervical surface epithelium at the 12 and 6 o'clock positions. A return electrode was then placed on the patient's thigh (for grounding) and the cervix was painted with Schiller solution. The loop electrodes were made from tungsten wire and were fabricated in various sizes ranging from 1x1 to 2x2 cm. The electrosurgical generator (Birtcher, Bovie) was operated in the cutting current mode at 35-50 W output. The loop was pushed into the cervical tissue perpendicular to the surface, then slowly drawn across the lesion parallel to the surface of the cervix to the opposite side. The entire transformation zone and CIN lesions were excised to a depth of 5-8 mm. in a single pass of the electrode when possible. We attempted to excise the epithelium 4-5 mm. beyond the point at which it became Schiller negative. To prevent post treatment bleeding from

the crater base we cauterized the base and edges of the crater. The specimen was sent for histologic examination.

Patients were advised about potential early and late complications and were advised to avoid sexual intercourse for one month. The subjects returned 1-3 months post-treatment for a Papanicolaou test and a complete colposcopic examination.

Results

Over a 7-month period in 1993-1994 we treated 31 subjects with biopsy-proven CIN using LEEP. The average age of the patients was 35.4 years, with a range of 22-61 years. The procedure was well tolerated by the patients and required only local anesthesia in 29 of the 31 cases. Two cases required general anesthesia due to discomfort and irritability during the procedure. The operation time was less than 5 minutes in 22 cases (71.0%). In the majority of cases (83.9%) perioperative and postoperative hemorrhage was minimal. No patient needed blood transfusion or subsequent admission to the hospital due to late hemorrhage. Two cases experienced minimal late hemorrhage during the first week and responded to AgNo₃ application.

A satisfactory specimen was obtained for histologic examination in the majority of cases. Only in a few cases did the amount of thermal damage in the excised tissue preclude complete histologic evaluation. Four cases showed no histologic evidence of CIN and one showed HPV infection (Table 1). In the patients with CIN, the excision margins were reported as complete in all but one case. The patients with microinvasive cancer were treated subsequently with hysterectomy and no residual disease was detected in the surgical specimens. In 28 of the 31 cases (90.3%),

this procedure was sufficient for complete treatment.

Table 1. Histopathology of the cervix from LEEP (Total = 31)

	No.
- Chronic cervicitis	3
- HPV	1
- CIN	
- mild dysplasia	5
- moderate dysplasia	6
- severe dysplasia, CIS	14
- Microinvasive cancer	2

The healing process after the procedure was also observed. AT 1 week post-treatment, the crater base was filled with soft, tan-yellow, necrotic material. The edges of the crater were well demarcated and had minimal reddening. Uncomplicated cases examined 1-3 months after the procedure were either totally healed or had only a small rim of granulation tissue remaining at the external os.

Discussion

In the treatment of CIN, excisional methods that provide tissue specimens for histopathologic diagnosis have distinct advantages over ablative methods that destroy the affected tissues. Unfortunately, most excisional methods have a number of disadvantages that tend to outweigh their advantages. Cold knife conization is usually performed in a hospital setting under general or spinal anesthesia and has a high rate of complications, even in expert hands. Major bleeding occurs in 13% of the cases, cervical stenosis develops in

17%, and infertility and cervical incompetence are infrequent but well-recognized complications.⁽¹⁶⁾ Because of these disadvantages, many clinicians are now advocating the technique of loop electrosurgical excision procedure.

We have described our early experience using LEEP in the management of CIN. LEEP is a combined diagnostic and therapeutic procedure in which the entire transformation zone is removed. The loop excision procedure was performed in less than 5 minutes in the majority of cases, can be performed on an outpatient basis using local anesthesia, and was well tolerated by the patients. The procedure was easy to perform and required only a relatively inexpensive electrosurgical generator. The complication rate of the procedure was low. Perioperative and postoperative hemorrhage was minimal in the majority of cases. The finding that 2 cases (6.4%) experienced secondary or late hemorrhage during the healing process is a rate similar to previous results reported by Keijser (5.8%).⁽¹⁷⁾ Although there were no immediate operative complications in this study, a larger study series will be required to confirm that operative morbidity with LEEP is decreased in comparison with cold knife conization.

The quality of the specimens obtained in our study permitted an examination of both the cervical abnormalities and excision margins. In a study of 616 patients with abnormal cervical smears, the histologic specimens were adequate in over 90% of the cases.⁽¹⁴⁾ However, some studies show that LEEP provides specimens that are inferior compared with cold knife conization.⁽¹⁹⁾ However, histopathologic results from the specimens are the other interesting issue that we plan to go on further study.

The overall success rate of LEEP in this study was 90.3%. This success rate is similar to other published series using LEEP to treat CIN.^(16,17)

The results of this preliminary study conclude that LEEP tends to be a reliable, well tolerated, economical and efficient technique that offers a good alternative in the management of CIN with a low incidence of short-term morbidity.

References

1. Cancer Facts and Figures-1989. Atlanta : American Cancer Society, 1989.
2. Jones HW 3d, Buller RE. The treatment of cervical intraepithelial neoplasia by cone biopsy. *Am Obstet Gynecol* 1980 Aug 15; 13(8):882-6
3. Jones HW 3d. Treatment of cervical intraepithelial neoplasia. *Clin Obstet Gynecol* 1990 Dec; 33(4): 826-36
4. Benedet JL, Anderson GJ, Boyes DA. Colposcopic accuracy in the diagnosis of microinvasive and occult invasive carcinoma of the cervix. *Obstet Gynecol* 1985 Apr; 65(4): 557-62
5. Townsend DE, Richart RM, Marks E, Nielson J. Invasive cancer following outpatient evaluation and therapy for cervical disease. *Obstet Gynecol* 1981 Feb; 57(2):145-9
6. McIndoe GAJ, Robson MS, Tidy JA, Mason WP, Anderson MC. Laser excision rather than vaporization : the treatment of choice for cervical intraepithelial neoplasia. *Obstet Gynecol* 1989 Aug; 74(2):165-8
7. Partington CK, Turner MJ, Soutter WP, Griffiths M, Krausz T. Laser vaporization versus laser excision conization in the treatment

- of cervical intraepithelial neoplasia. *Obstet Gynecol* 1989 May; 73(5pt 1): 775-9
8. Gunasekera PC, Phipps JH, Lewis BV. Large loop excision of the transformation zone (LLETZ) compared to carbon dioxide laser in the treatment of CIN : a superior mode of treatment. *Br J Obstet Gynaecol* 1990 Nov; 97(11): 995-8
 9. Cartier R. *Practical Colposcopy*. 2nd ed. Paris : Laboratoire Cartier, 1984 : 139-56
 10. Prendiville W, Cullimore NS, Norman S. Large loop excision of the transformation zone (LLETZ). A new method of management for women with cervical intraepithelial neoplasia. *Br J Obstet Gynaecol* 1989 Sep; 96(9): 1054-60
 11. Whiteley PF, Olah KS. Treatment of cervical intraepithelial neoplasia ; experience with the low-voltage diathermy loop. *Am J Obstet Gynecol* 1990 May; 162(5):1272-7
 12. Mot-Yosef S, Lopes A, Pearson S, Monaghan JM, Loop diathermy cone biopsy. *Obstet Gynecol* 1990 May; 75(5):884-6
 13. Bigrigg MA, Codling BW, Pearson P, Read MD, Swingler GR. Colposcopic diagnosis and treatment of cervical dysplasia at a single clinic visit. Experience of low-voltage diathermy loop in 1000 patients. *Lancet* 1990 Jul 28; 336(8709):229-31
 14. Luesley DM, Cullimore J, Redman CWE, Lawton FG, Emens JM, Rollason TP, Williams DR, Buxton EJ. Loop diathermy excision of the cervical transformation zone in patients with abnormal cervical smears. *Br Med J* 1990 Jun 30; 300(6741):1690-3
 15. Murdoch JB, Grimshaw RN, Monaghan JM. Loop diathermy excision of the abnormal cervical transformation zone. *Int J Gynecol Cancer* 1991; 1:105-11
 16. Wright TC JR, Gagnon S, Richart RM, Ferenczy A. Treatment of cervical intraepithelial neoplasia using the loop electrosurgical excision procedure. *Obstet Gynecol* 1992 Feb; 79(2): 173-8
 17. Keijser KG, Kenemans P, Van der Zanden PH, Schiff CP, Vooijs GP, Rolland R. Diathermy loop excision in the management of cervical intraepithelial neoplasia : diagnosis and treatment in one procedure. *Am J Obstet Gynecol* 1992; 166(4):1281-7
 18. Jone HW 3d. Cone biopsy in the management of cervical intraepithelial neoplasia. *Clin Obstet Gynecol* 1983 Dec; 26(4):968-79
 19. Krebs HB, Pastore L, Helmkamp BF. Loop electrosurgical excision Procedures for cervical dysplasia : experience in a community hospital. *Am J Obstet Gynecol* 1993 Aug; 163(2pt 1):289-95