

Chulalongkorn University

Chula Digital Collections

Chulalongkorn University Theses and Dissertations (Chula ETD)

2018

Protective effects of eriodictyol on hydrogen peroxide-induced cell death in SH-SY5Y cells

Saowarose Thongin
Graduate School

Follow this and additional works at: <https://digital.car.chula.ac.th/chulaetd>




Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Thongin, Saowarose, "Protective effects of eriodictyol on hydrogen peroxide-induced cell death in SH-SY5Y cells" (2018). *Chulalongkorn University Theses and Dissertations (Chula ETD)*. 2565.
<https://digital.car.chula.ac.th/chulaetd/2565>

This Thesis is brought to you for free and open access by Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn University Theses and Dissertations (Chula ETD) by an authorized administrator of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

PROTECTIVE EFFECTS OF ERIODICTYOL ON HYDROGEN PEROXIDE-
INDUCED CELL DEATH IN SH-SY5Y CELLS



A Thesis Submitted in Partial Fulfillment of the Requirements
for the Degree of Master of Science in Pharmacology
Inter-Department of Pharmacology
Graduate School
Chulalongkorn University
Academic Year 2018
Copyright of Chulalongkorn University

ฤทธิ์ปกป้องของสารอิลิโอดิกไทออกต่อการตายของเซลล์ที่ถูกเหนี่ยวนำด้วยไฮโดรเจนเปอร์
ออกไซด์ในเซลล์ SH-SY5Y



วิทยานิพนธ์นี้เป็นส่วนหนึ่งของการศึกษาตามหลักสูตรปริญญาวิทยาศาสตรมหาบัณฑิต
สาขาวิชาเภสัชวิทยา สหสาขาวิชาเภสัชวิทยา
บัณฑิตวิทยาลัย จุฬาลงกรณ์มหาวิทยาลัย
ปีการศึกษา 2561
ลิขสิทธิ์ของจุฬาลงกรณ์มหาวิทยาลัย

Thesis Title	PROTECTIVE EFFECTS OF ERIODICTYOL ON HYDROGEN PEROXIDE-INDUCED CELL DEATH IN SH-SY5Y CELLS
By	Miss Saowarose Thongin
Field of Study	Pharmacology
Thesis Advisor	Assistant Professor Ratchanee Rodsiri, Ph.D.
Thesis Co Advisor	Assistant Professor Varisa Pongrakhananon, Ph.D.

Accepted by the Graduate School, Chulalongkorn University in Partial Fulfillment of the Requirement for the Master of Science

..... Dean of the Graduate School
(Associate Professor Thumnoon Nhujak, Ph.D.)

THESIS COMMITTEE

..... Chairman
(Associate Professor Suree Jianmongkol, Ph.D.)

..... Thesis Advisor
(Assistant Professor Ratchanee Rodsiri, Ph.D.)

..... Thesis Co-Advisor
(Assistant Professor Varisa Pongrakhananon, Ph.D.)

..... Examiner
(Chenphop Sawangmake, Ph.D.)

..... External Examiner
(Rawiwan Maniratanachote, Ph.D.)

เสาวรส ทองอินทร์ : ฤทธิ์ปกป้องของสารอีลิโอติกไทออลต่อการตายของเซลล์ที่ถูกเหนี่ยวนำด้วยไฮโดรเจนเปอร์ออกไซด์ในเซลล์ SH-SY5Y. (

PROTECTIVE EFFECTS OF ERIODICTYOL ON HYDROGEN PEROXIDE-INDUCED CELL DEATH IN SH-SY5Y CELLS) อ.ที่ปรึกษาหลัก : ผศ. ญ. ดร.

รัชณี รอดศิริ, อ.ที่ปรึกษาร่วม : ผศ. ญ. ดร.วิรัช พงศ์เรณานนท์

ภาวะเครียดออกซิเดชันและการทำงานของไมโทคอนเดรียที่ผิดปกติ เป็นสาเหตุหลักของการตายของเซลล์ประสาทในกลุ่มโรคความเสื่อมของระบบประสาท อีลิโอติกไทออล เป็นสารกลุ่มฟลาโวนอยด์ สกัดมาจากต้น *Dendrobium ellipsophyllum* (ชื่อไทย : เอื้องทอง) ผลการศึกษาก่อนหน้ารายงานฤทธิ์ต้านอนุมูลอิสระและฤทธิ์ต้านอักเสบของอีลิโอติกไทออล งานวิจัยนี้มีวัตถุประสงค์เพื่อศึกษาฤทธิ์และกลไกปกป้องของอีลิโอติกไทออลต่อการตายของเซลล์ที่ถูกเหนี่ยวนำด้วยไฮโดรเจนเปอร์ออกไซด์ในเซลล์ SH-SY5Y ฤทธิ์ปกป้องของอีลิโอติกไทออลต่อการตายของเซลล์ที่ถูกเหนี่ยวนำด้วยไฮโดรเจนเปอร์ออกไซด์ประเมินโดยวิธี resazurin assay ผลต่อระดับอนุมูลอิสระภายในเซลล์และผลต่อ mitochondria membrane potential ศึกษาโดยวิธี DCFH-DA assay และวิธี TMRE assay ตามลำดับ และผลของอีลิโอติกไทออลต่อการส่งสัญญาณการตายแบบ apoptosis ผ่านทางไมโทคอนเดรีย ศึกษาโดยวิธี Western blot ผลการทดลองพบว่าอีลิโอติกไทออล (0.1, 1 และ 10 μ M) เพิ่มจำนวนเซลล์ที่มีชีวิต ($p < 0.001$) ลดระดับอนุมูลอิสระภายในเซลล์ ($p < 0.05$) และเพิ่มความเข้มแสงฟลูออเรสเซนซ์ของ TMRE ($p < 0.05$) เมื่อเปรียบเทียบกับกลุ่มที่ได้รับไฮโดรเจนเปอร์ออกไซด์เพียงอย่างเดียว ไฮโดรเจนเปอร์ออกไซด์เหนี่ยวนำการส่งสัญญาณการตายแบบ apoptosis ผ่านทางไมโทคอนเดรีย โดยการลดอัตราส่วนของ Bcl-2/BAX เพิ่มการแสดงออกของ cytochrome c, caspase-3 และ cleaved caspase-3 แต่การให้สารอีลิโอติกไทออลจะลดเหตุการณ์เหล่านี้ จึงสรุปการทดลองได้ว่าอีลิโอติกไทออลสามารถป้องกันการตายของเซลล์ SH-SY5Y ที่ถูกเหนี่ยวนำด้วยไฮโดรเจนเปอร์ออกไซด์ ผ่านทางฤทธิ์ต้านอนุมูลอิสระ ซึ่งส่งผลให้ยับยั้งการตายของเซลล์แบบ apoptosis ผ่านทางไมโทคอนเดรีย

สาขาวิชา เกษตรวิทยา

ปีการศึกษา 2561

ลายมือชื่อนิสิต

ลายมือชื่อ อ.ที่ปรึกษาหลัก

ลายมือชื่อ อ.ที่ปรึกษาร่วม

5987222620 : MAJOR PHARMACOLOGY

KEYWORD: Eriodictyol, Hydrogen peroxide, Reactive oxygen species, SH-SY5Y cells

Saowarose

Thongin

:

PROTECTIVE EFFECTS OF ERIODICTYOL ON HYDROGEN PEROXIDE-

INDUCED CELL DEATH IN SH-SY5Y CELLS. Advisor: Asst. Prof. Ratchanee

Rodsiri, Ph.D. Co-advisor: Asst. Prof. Varisa Pongrakhananon, Ph.D.

Oxidative stress and mitochondrial dysfunction are the major causes of neuronal death in neurodegenerative disorders. Eriodictyol is a flavonoid extracted from the whole plant of *Dendrobium ellipsophyllum* (thai name: Ueang Thong). The antioxidant and anti-inflammatory effects of eriodictyol have been widely reported. The present study aimed to investigate the protective effect and mechanisms of eriodictyol on hydrogen peroxide (H_2O_2)-induced cell death in human neuroblastoma SH-SY5Y cells. The protective effect of eriodictyol on cytotoxicity induced by H_2O_2 was evaluated using resazurin assay. The effects on intracellular reactive oxygen species (ROS) and mitochondria membrane potential were evaluated by 2'-7' dichlorodihydrofluorescein diacetate (DCFH-DA) assay and tetramethylrhodamine ethyl ester (TMRE) assay, respectively. The effects of eriodictyol on mitochondria-mediated apoptosis cascade were determined using western blot analysis. The results showed that pretreatment with eriodictyol (0.1, 1 and 10 μM) increased cell viability ($p < 0.001$), reduced intracellular ROS ($p < 0.05$) and increased TMRE fluorescent intensity ($p < 0.05$) compared to H_2O_2 -treated alone. H_2O_2 induced mitochondria-mediated apoptotic signalling by decreasing Bcl-2/BAX ratio, increasing the expressions of cytochrome c, caspase-3 and cleaved caspase-3 while eriodictyol pretreatment reduced these events. In conclusion, eriodictyol can protect against H_2O_2 -induced cytotoxicity in SH-SY5Y cells via an antioxidant effect, and consequently inhibit mitochondria-mediated apoptosis cascade.

Field of Study: Pharmacology

Student's Signature

Academic Year: 2018

Advisor's Signature

Co-advisor's Signature

ACKNOWLEDGEMENTS

The completion of this study could not have been possible without the guidance and support of several people, to whom I am grateful.

First, I would like to express the deepest appreciation to my thesis advisor, Assistant Professor Dr. Ratchanee Rodsiri and my co-advisor, Assistant Professor Assistant Professor Dr. Varisa Pongrakhananon for their invaluable advice, attention, motivation, encouragement and patience throughout this research work. Also, I would like to thank the thesis committees for their invaluable suggestions.

Second, I would like to acknowledge Associate Professor Dr. Boonchoo Sritularak for eriodictyol supported.

Next, I am grateful to all members of Department of Pharmacology and Physiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University for their help, suggestion, encouragement and assistance.

Moreover, This work was financially supported by the 90th Anniversary of Chulalongkorn University Fund (Ratchadaphisedksomphot Endowment Fund)

Finally, I wish to express my infinite thank to my family, especially my mother, for their endless love, understanding and never-ending support throughout my life. Thank you for always being beside me.

จุฬาลงกรณ์มหาวิทยาลัย
CHULALONGKORN UNIVERSITY

Saowarose Thongin

TABLE OF CONTENTS

	Page
ABSTRACT (THAI).....	iii
ABSTRACT (ENGLISH).....	iv
ACKNOWLEDGEMENTS	v
TABLE OF CONTENTS.....	vi
LIST OF FIGURES	ix
ABBREVIATIONS	x
CHAPTER 1 INTRODUCTION	1
1.1 Background and Rationale	1
1.2 Research questions	1
1.3 Objectives.....	2
1.4 Hypothesis	2
1.5 Expected Benefits.....	2
1.6 Conceptual Framework	3
CHAPTER 2 LITERATURES REVIEW	4
2.1 Neurodegenerative disorders.....	4
2.2 Reactive oxygen species and oxidative stress	5
2.3 Mitochondria-mediated apoptosis cascade.....	6
2.4 Eriodictyol.....	8
CHAPTER 3 MATERIALS AND METHODS	10
3.1 Materials.....	10
3.1.1 Cell culture and treatment	10

3.1.2 Chemicals and Reagents	10
3.1.3 Equipment	11
3.1.4 Materials.....	12
3.2 Chemical preparation	12
3.2.1 Preparation of eriodictyol stock solution	12
3.2.2 Hydrogen peroxide (H ₂ O ₂).....	12
3.3 Methods.....	13
3.3.1 Experimental design.....	13
3.3.2 Determination of non-toxic concentrations of eriodictyol.....	14
3.3.3 Determination of toxic concentrations of H ₂ O ₂	14
3.3.4 Determination of the protective effect of eriodictyol on H ₂ O ₂ -induced cell death	15
3.3.5 Determination of the effects of eriodictyol on intracellular ROS levels.....	15
3.3.6 Determination of the effect of eriodictyol on mitochondrial membrane potential	16
3.3.7 Determination of the effects of eriodictyol on mitochondria-mediated apoptosis cascade	17
3.3.8 Statistical Analysis.....	17
CHAPTER 4 RESULTS	18
4.1 The effect of eriodictyol on cell viability	18
4.2 The effect of H ₂ O ₂ on cell viability	19
4.3 The protective effect of eriodictyol on H ₂ O ₂ -induced cell death	20
4.4 The protective effects of eriodictyol on H ₂ O ₂ -induced ROS generation	21
4.5 The effect of eriodictyol and H ₂ O ₂ on mitochondrial membrane potential.....	22
4.6 The effects of eriodictyol and H ₂ O ₂ on mitochondria-mediated apoptosis cascade	24
CHAPTER 5 DISCUSSION AND CONCLUSION	29

REFERENCES	32
VITA	37



LIST OF FIGURES

	Page
Figure 1 Types of reactive oxygen species (ROS).....	5
Figure 2 The generation process of reactive oxygen species (ROS))	6
Figure 3 Apoptotic signaling pathways.....	7
Figure 6 Principle of resazurin assay	14
Figure 7 Principle of 2,7-dichlorodihydrofluorescein diacetate assay	16
Figure 8 The effect of eriodictyol (0.001-100 μ M) on cell viability in SH-SY5Y cells.....	18
Figure 9 The effect of H_2O_2 (100-1000 μ M) on cell viability in SH-SY5Y cells.....	19
Figure 10 The protective effects of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells..	20
Figure 11 The effects of eriodictyol on intracellular ROS levels induced by H_2O_2 ..	21
Figure 12 The effects of eriodictyol and H_2O_2 on mitochondrial membrane potential	23
Figure 13 The effects of eriodictyol and H_2O_2 on BAX and Bcl-2 protein expression.....	25
Figure 14 The effects of eriodictyol and H_2O_2 on cytochrome c expression.	26
Figure 15 The effects of eriodictyol and H_2O_2 on caspase-3 and cleaved caspase-3 expression...	28
Figure 16 The action mechanisms of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells	31

ABBREVIATIONS

α = alpha

β = beta

$^{\circ}\text{C}$ = degree celsius

/ = per

% = percentage

\pm = plus-minus sign

x g = relative centrifugal force

$\mu\text{g/ml}$ = Microgram per milliliter

μl = Microliter

μM = Micromolar

AD = Alzheimer's disease

ANOVA = Analysis of variance

Apaf-1 = Apoptotic protease-activating factor-1

ATCC = American Type Culture Collection

ATP = Adenosine triphosphate

ALS = amyotrophic lateral sclerosis

BAD = Bcl-2 antagonist of cell death

BAK = Bcl-2-antagonist/killer-1

BAX = Bcl-2-associated X protein

Bcl-2 = B-cell lymphoma-2

Caspase = Cysteine aspartic acid specific protease

CAT = Catalase

CO₂ = Carbon dioxide

DCFH-DA = 2', 7'- dichlorodihydrofluorescein diacetate

DMEM-F12 = Dulbecco's Modified Essential Eagle's
Medium-Ham's Nutrient Mixture F-12

DMSO = Dimethyl sulfoxide

DNA = Deoxyribonucleic acid

EDTA = Ethylene diamine tetraacetic acid

ETC = Mitochondrial electron transport chain

FBS = Fetal bovine serum

FDA = Food and drug administration

GAPDH = Glyceraldehyde 3-phosphate dehydrogenase

GSH = Glutathione

GST = Gutathione-S-transferase

GPx = Glutathione peroxidase

h = Hour

H₂O₂ = Hydrogen peroxide

HCl = Hydrochloric acid

HD = Huntington's disease

IC ₅₀	=	50% Inhibition concentration
IL	=	Interleukin
LPS	=	Lipopolysaccharides
mg/ml	=	Milligram per milliliter
MMP	=	Mitochondria membrane potential
MPTP	=	1-methyl-4-phenyl-1,2,3,6- tetrahydropyridine
NaHCO ₃	=	Sodium bicarbonate
NaOH	=	Sodium hydroxide
NF- κ B	=	Nuclear factor kappa-light-chain-enhancer of activated B cells
nm	=	Nanometer
Nrf2	=	The nuclear factor erythroid 2–related factor 2
•O ₂ ⁻	=	Superoxide anion
O ₂	=	Oxygen
•O ₂ ⁻²	=	Peroxide
•OH	=	Hydroxyl radical
HO ² •	=	Hydroperoxyl
PD	=	Parkinson's disease
P53	=	Tumor protein 53
PBS	=	Phosphate buffer saline

pH = The negative logarithm of hydrogen ion
concentration

ROS = Reactive oxygen species

rpm = Round per minutes

S.E.M. = Standard error of mean

SOD = Superoxide dismutase

TNF α = Tumor necrosis factor alpha



CHAPTER 1

INTRODUCTION

1.1 Background and Rationale

Neuronal death in neurodegenerative disorders are mainly caused by oxidative stress and mitochondrial dysfunction (Wang et al., 2014). The current pharmacological treatment of neurodegenerative diseases mainly alleviates the dysfunction of neurotransmission but they do not restore dead neurons or regenerate neurites. The neuroprotective approach is needed to slow down the disease progression.

Oxidative stress is resulted from the accumulation of reactive oxygen species (ROS) and the inadequate protective mechanisms of the cells. Hydrogen peroxide (H_2O_2), one of ROS, can induce neuronal death via an increase of ROS and the decrease of antioxidant enzymes (Feng et al., 2016; Park et al., 2015). In addition, H_2O_2 causes the translocation of cytochrome c from mitochondria to cytosol leading to apoptosis (Cai et al., 2008). Therefore, H_2O_2 is used in this study to induce oxidative stress and neuronal death in human neuroblastoma SH-SY5Y cells.

Eriodictyol is a flavonoid extracted from the whole plant of *Dendrobium ellipsophyllum* or Ueang Thong. Previous studies demonstrated the antioxidant and anti-inflammatory effects of eriodictyol (Zhu et al., 2015). However, the protective effect of eriodictyol has not yet clarified.

The present study aimed to investigate the protective effect of eriodictyol in H_2O_2 -induced SH-SY5Y cell death. The mechanisms underlying the protective effect of eriodictyol involving mitochondria-mediated apoptosis.

1.2 Research questions

1. Does eriodictyol protect against the neurotoxic effect of H_2O_2 in SH-SY5Y cells?
2. What are the mechanisms underlying the protective effect of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells?

1.3 Objectives

1. To investigate the protective effect of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells.
2. To investigate the mechanisms underlying the protective effect of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells.


1.4 Hypothesis

Eriodictyol protects against H_2O_2 -induced neuronal cell death in SH-SY5Y cells via the decrease of ROS production, the protection of mitochondria membrane potential dysfunction and the inhibition of mitochondria-mediated apoptosis cascade.

1.5 Expected Benefits

This research will provide a scientific information regarding the protective effect of eriodictyol on the neuronal cells. The data suggest further study to investigate the neuroprotective effects of eriodictyol in the *in vivo* models of neurodegenerative diseases.

1.6 Conceptual Framework



CHAPTER 2

LITERATURES REVIEW

2.1 Neurodegenerative disorders

Neurodegenerative disorders are characterized by progressive and irreversible loss of neurons in the central and peripheral nervous systems. These disorders include Alzheimer's disease (AD), Parkinson's disease (PD), amyotrophic lateral sclerosis (ALS) and Huntington's disease (HD). The clinical symptoms of neurodegenerative disorders are varied depending on types of neurons and brain areas defected; such as cognitive impairments in AD, and motor dysfunctions in PD, HD and ALS. Nowadays, there is no treatment for recovering neuronal death. The available treatments are to reduce the symptoms by altering neurotransmission. Therefore, the protection of neuronal death is considered as an important therapeutic approach to delay the progression of neuronal loss in neurodegenerative disorders.

The pathology of neurodegenerative disorders involves oxidative stress (Thanan et al., 2014). The increase of lipid peroxidation and the decreases of glutathione (GSH), glutathione peroxidase (GPx), glutathione reductase, glutathione-S-transferase (GST), glucose-6-phosphate dehydrogenase, superoxide dismutase (SOD) and catalase (CAT) have been reported in the frontal cortex of AD patients (Ansari & Scheff, 2010). In addition, the transgenic mouse model of AD presented mitochondria dysfunction and the increase of ROS in the brains (David et al., 2005). To produce an animal model of PD, a mitochondrial complex I inhibitor, 1-methyl-4-phenyl-1,2,3,6- tetrahydropyridine (MPTP), caused the elevation of malondialdehyde level an oxidative stress marker, and the depletion of GSH leading to parkinsonism behaviors in rats (Xu et al., 2013). Moreover, a transgenic mouse model of HD showed the increases of hydroxyguanosine (8-OHG) and 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels which are the markers oxidative DNA damage (Acevedo-Torres et al., 2009). Taken together, oxidative stress is an important cause of neuronal death in the neurodegenerative disorders.

2.2 Reactive oxygen species and oxidative stress

Reactive oxygen species (ROS) are defined as a group of reactive molecules derived from oxygen (O_2). ROS include peroxide ($\bullet O_2^{-2}$), superoxide anion ($\bullet O_2^-$), hydroxyl radical ($\bullet OH$), hydroperoxyl (HO_2^\bullet) and hydrogen peroxide (H_2O_2) (Figure 1). ROS is generally short-lived and highly reactive because of their unpaired valence electrons (Patten et al., 2010). The major source of intracellular ROS is mitochondrial electron transport chain (ETC) (Patten et al., 2010). It is estimated that 1-2% of electrons are leaked during ETC resulting in the production of superoxide anion in the mitochondrial matrix (Circu & Aw, 2010; Murphy, 2009). Superoxide anion can be transformed into H_2O_2 by superoxide dismutase (SOD). H_2O_2 can be further transformed into the more toxic forms; hydroxyl radicals and hydroxyl anions, by Fenton reaction. Moreover, superoxide anion can be transformed into hydroperoxyl (Birben et al., 2012; Kim et al., 2015) (Figure 2). ROS cause cellular lipid, protein and DNA damage leading to oxidative stress, mitochondrial dysfunction, and cell death (Blesa et al., 2015; Kim et al., 2015; Valko et al., 2007). In normal condition, ROS production and protective mechanisms in the cells; such as antioxidants and antioxidative enzymes, are balanced. However, the excessive ROS generation and the depletion of the antioxidant system in the pathological condition resulted in oxidative stress in the cells (Kwon et al., 2014).




Figure 1 Types of reactive oxygen species (ROS) (Kim et al., 2015)




Figure 2 The generation process of reactive oxygen species (ROS) (Kwon et al., 2014)

Hydrogen peroxide (H_2O_2) has been used in various neurotoxic models *in vitro*. Hydrogen peroxide diffuses into plasma membrane (Sies, 2017) and is readily converted into the highly toxic hydroxyl radicals via Fenton reaction (Kim et al., 2015). H_2O_2 induced the increase of ROS, lipid peroxidation and mitochondria dysfunction leading to mitochondrial mediated apoptosis in human neuroblastoma SH-SY5Y cells (de Oliveira et al., 2018; Yang et al., 2016). Moreover, H_2O_2 stimulated apoptotic cell death in cardiomyocytes, associated with PARP cleavage, DNA fragmentation, upregulation of p53, loss of mitochondria membrane potential, BAX/BAD mitochondrial translocation, mitochondrial cytochrome c release and caspase-3 activation (Cook et al., 1999). In addition, H_2O_2 inhibit expression of Bcl-2 protein in human promyelocytic leukemia HL-60 cell line (Lee et al., 2000). Hence, H_2O_2 will be used to generate the *in vitro* neurotoxic model in this study.

2.3 Mitochondria-mediated apoptosis cascade

Apoptosis is the process of programmed cell death. Apoptosis normally occurs to maintain cell populations in tissues and to eliminate damaged cells (Elmore, 2007). However, progressive neuronal death via apoptosis leads to neurodegenerative diseases.

Apoptosis is triggered by extrinsic or intrinsic signaling pathways. Mitochondria plays an important role in the intrinsic signaling pathways of apoptosis. DNA damage, ER stress, hypoxia and metabolic stress can activate BH3-only proteins. BH3-only proteins then activates BAX and BAK resulting in translocation of BAX and BAK to the outer membrane of mitochondria.

This event leads to pore formation, mitochondrial membrane potential alteration and mitochondrial damage. Cytochrome c is subsequently released from mitochondria and then associates with adaptor protein apoptotic protease activating factor 1 (Apaf-1) to form apoptosome. Apoptosome sequentially activates caspase-9, caspase-3 and caspase-7. On the other hand, an anti-apoptotic protein, Bcl-2 proteins, inhibit cell death by inhibiting BAX and BAK (Circu & Aw, 2010; Ichim & Tait, 2016) (Figure 3).




Figure 3 Apoptotic signaling pathways (Ichim & Tait, 2016)

2.4 Eriodictyol

A flavonoid, eriodictyol ($C_{15}H_{12}O_6$) (Figure 4), is extracted from the whole plant of *Dendrobium ellipsophyllum* (thai name: Ueang Thong) (Figure 5). Eriodictyol is colorless powder with molecular weight 258 g/mol, and Log P 2.02 (U.S. Environmental Protection Agency, 2018). It exhibits beneficially pharmacological properties including anti-metastatic (Tanagornmeatar et al., 2014), antioxidant and anti-inflammatory effects (Zhu et al., 2015). Eriodictyol attenuated LPS-induced acute lung injury by inhibiting the inflammatory cytokine such as TNF- α , IL-6 and IL-1 β (Zhu et al., 2015). In addition, eriodictyol attenuated β -amyloid 25–35-induced oxidative cell death in primary cultured neuron by activating Nrf2 (Jing et al., 2015). Moreover, eriodictyol protected against linoleic acid hydroperoxide and H_2O_2 -induced PC12 cell death (Lou et al., 2012; Sasaki et al., 2003)




Figure 4 Chemical structure of eriodictyol



Figure 5 *Dendrobium ellipsophyllum* or Ueang Thong

CHAPTER 3

MATERIALS AND METHODS

3.1 Materials

3.1.1 Cell culture and treatment

The human neuroblastoma SH-SY5Y cell line was purchased from ATCC (Manassas, VA, USA). SH-SY5Y cells were cultured in Dulbecco's Modified Essential Eagle's Medium-Ham's Nutrient Mixture F-12 (DMEM-F12) supplemented with 10% heat-inactivated fetal bovine serum (FBS) and 0.1% penicillin/streptomycin at 37 °C in a humidified atmosphere of 5% CO₂ and 95% air. The medium was changed every 2 days. Cells were sub-cultured at 80–90% confluent.

3.1.2 Chemicals and Reagents

Anti-cytochrom c antibody (Santa Cruz Biotechnology, CA, USA)

Anti-GAPDH antibody (Millipore, Billerica, MA, USA)

BCA protein assay kit (Thermo Scientific, Rockford, IL, USA)

2'-7' Dichlorodihydrofluorescein diacetate (DCFH-DA) (Sigma, USA)

Dimethyl sulfoxide (DMSO) (Fisher Scientific, Loughborough, UK)

Dulbecco's Modified Essential Eagle's Medium-Ham's Nutrient Mixture F-12 (DMEM-F12) (Gibco, USA)

Ethanol (Merk, Germany)

Fetal bovine serum (Gibco, New Zealand)

Horseradish peroxidase (HRP)-conjugated goat anti-rabbit IgG antibody (Millipore, Billerica, MA, USA)

Hydrogen peroxide (H₂O₂) (Ajax Finechem, Auckland, New Zealand)

Nonfat Dry Milk (Cell Signaling Technology, Inc. USA)

Penicillin/streptomycin (Gibco, USA)

Polyvinylidene difluoride transfer membrane (Immobilon, Bedford, MA, USA)

Resazurin (Sigma, USA)

Tetramethylrhodamine ethyl ester (Sigma, USA)

Phosphate buffered saline (Gibco, USA)

Protein ladder (BLUEstain[™], Gold Biotechnology, Inc. US)

Rabbit monoclonal anti-BAX antibody (Cell Signaling Technology, Inc. USA)

Rabbit monoclonal anti-Bcl-2 antibody (Sigma-Aldrich, St. Louis, MO. USA)

Rabbit monoclonal anti-caspase-3 antibody (Cell Signaling Technology, Inc. USA)

Tris-HCl (Sigma-Aldrich, St. Louis, MO. USA)

0.4% Trypan blue dye (Sigma, USA)

0.25% Trypsin-Ethylenediaminetetraacetic acid (EDTA) (Gibco, USA)

3.1.3 Equipment

Analytical balance (Mettler Toledo, Switzerland)

Autoclave (Hirayama, Japan)

Autopipette (Gilson, USA)

Centrifuge (Hettich, USA)

Controller pipette (Gilson, USA)

CO₂ incubator (Thermo Fisher Scientific, USA)

Light microscope (Nikon, Japan)

Micro centrifuge (Thermo Fisher Scientific, USA)

Microplate fluorescence reader (CLARIOstar®, BMG LABTECH, Germany).

pH meter (Schott, UK)

Power supply (PowerPac[™], USA)

Trans-Blot SD sem-dry electrophoretic transfer cell (Trans-Blot®, USA)

Vortex mixer (Scientific Industries, USA)

3.1.4 Materials

25 and 75 ml cell culture flask (Gibco, USA)
15 and 50 ml centrifuge tube (Corning Inc., USA)
1.7 ml micro centrifuge tube (Accumax, USA)
6-well plate (Corning Inc., USA)
96-well plate (Corning Inc., USA)
96-well plate black flat bottom polystyrene (Corning Inc., USA)
200 and 1250 μ l universal grad tip (Accumax, USA)

3.2 Chemical preparation

3.2.1 Preparation of eriodictyol stock solution


Eriodictyol was prepared as previously described (Tanagornmeatar et al., 2014) and provided by Associate Professor Dr. Boonchoo Sritularak, Department of Pharmacognosy and Pharmaceutical Botany, Faculty of Pharmaceutical Sciences, Chulalongkorn University (Tanagornmeatar et al., 2014). Fifty-mM eriodictyol stock solution was prepared in 100% DMSO and stored at 20 °C until use. In the experiments, the stock solution was diluted in culture medium to final concentrations. The 0.2% DMSO was used as a vehicle control.

3.2.2 Hydrogen peroxide (H₂O₂)

One hundred mM H₂O₂ stock solution was freshly prepared in deionized water. The stock solution was then diluted in medium to final concentrations.

3.3 Methods

3.3.1 Experimental design



3.3.2 Determination of non-toxic concentrations of eriodictyol

SH-SY5Y cells were seeded in 96-well plates at a density of 5×10^4 cells/well and incubated overnight at 37°C and 5% CO_2 . Cells were treated with eriodictyol 0.001, 0.01, 0.1, 1, 10 and 100 μM , and incubated for 24 h. Cell survival was evaluated by resazurin assay.

In resazurin assay, 10 μL of resazurin (0.05 mg/ml) was added to each well 4 h prior to the end of incubation time. Viable cells with active metabolism can reduce resazurin into the resorufin product (Figure 6) detected by a microplate fluorescent reader with at 530 nm excitation wavelength and 590 nm emission wavelength. The percentage of cell viability was calculated as follows:

$$\text{Cell viability (\% control)} = \frac{\text{Fluorescence intensity of sample}}{\text{Fluorescence intensity of control}}$$




Figure 4 Principle of resazurin assay (Riss et al., 2016)

3.3.3 Determination of toxic concentrations of H_2O_2

SH-SY5Y cells were seeded in 96-well plates at a density of 5×10^4 cells/well and incubated for 48 h. Cells were treated with H_2O_2 100, 200, 300, 400, 500, 600 and

1000 μM . Cell survival was evaluated by resazurin assay. The 50% inhibitory concentration (IC_{50}) was calculated and selected as a toxic concentration of H_2O_2 .

3.3.4 Determination of the protective effect of eriodictyol on H_2O_2 -induced cell death

SH-SY5Y cells were seeded in 96-well plates at a density of 5×10^4 cells/well and incubated overnight at 37°C and 5% CO_2 . Cells were treated with eriodictyol 0.1, 1, and 10 μM for 24 h. Then, the medium was removed. Cells were then treated with 200 μM H_2O_2 for 12 h. Cell viability was measured using resazurin assay.

3.3.5 Determination of the effects of eriodictyol on intracellular ROS levels

SH-SY5Y cells were seeded in 96-well plates at a density of 5×10^4 cells/well and incubated overnight at 37°C and 5% CO_2 . Cells were treated with eriodictyol 0.1, 1, and 10 μM for 24 h. Then, the medium was removed and cells were treated with 200 μM H_2O_2 for 1 hour. After incubation period, cells were washed twice with PBS and 100 μL of DCFH-DA was added (Hu et al., 2015; Kwon et al., 2014). The fluorescence intensity was measured by microplate reader.

In the DCFH-DA assay, 100 μL of DCFH-DA (25 μM) was added to each well and incubated for 30 minutes. DCFH-DA is the nonfluorescent dye. Oxidation of DCFH-DA produces a fluorescent molecule, dichlorofluorescein (DCF) (Figure 7). DCF was quantified using a microplate reader, with 488 nm excitation and 575 nm emission wavelengths. Intracellular ROS levels were expressed as fluorescence intensity and further calculated as a percentage of control.

$$\text{Intracellular ROS levels (\% control)} = \frac{\text{Fluorescent intensity of sample}}{\text{Fluorescent intensity of control}}$$




Figure 5 Principle of 2,7-dichlorodihydrofluorescein diacetate (DCFH-DA) assay (Gomes et al., 2005)

3.3.6 Determination of the effect of eriodictyol on mitochondrial membrane potential

SH-SY5Y cells were seeded in 96-well plates at a density of 5×10^4 cells/well and incubated overnight at 37°C and 5% CO_2 . Cells were treated with eriodictyol 0.1, 1, and 10 μM for 24 h. Then, the medium was removed and cells were treated with 200 μM H_2O_2 for 12 h. Cells were washed with PBS once time. Mitochondrial membrane potential was detected using tetramethylrhodamine ethyl ester (TMRE) assay.

In TMRE assay, 100 μL of TMRE (100 nM) was added to each well and incubated for 15 min. Fluorescent intensity of TMRE was measured using a microplate reader at excitation wavelength 585 nm and emission wavelength 535 nm. TMRE, a red-orange positive charge dye, can accumulate inside a healthy mitochondrion which exhibits negative membrane potential.

3.3.7 Determination of the effects of eriodictyol on mitochondria-mediated apoptosis cascade

SH-SY5Y cells were seeded in 6-well plates at a density of 2×10^6 cells/well and incubated with eriodictyol 0.1, 1, and 10 μM for 24 h. Then, medium was removed and cells were treated with 200 μM H_2O_2 for 12 h. Cell lysates were then collected. A proapoptotic protein, BAX, an anti-apoptotic protein, Bcl-2 and the markers of apoptosis, cytochrome c, cleaved caspase-3 and caspase-3 expression levels were detected using western blot analysis.

In the western blot analysis, cells are washed with PBS, harvested and centrifuged at 16,000 g for 15 min at 4 °C. The protein concentrations were determined using a BCA protein assay kit. Eighty- μg protein were separated on an 15% SDS–polyacrylamide gel, and then transferred onto a polyvinylidene difluoride membrane. The nonspecific proteins were blocked with 5% skim milk containing 0.5 mM Tris–HCl (pH 7.5), 150 mM NaCl, and 0.1% Tween-20 for 1 h at room temperature. The membrane was subsequently incubated with the primary antibodies (1:1000 anti-Bax, anti-Bcl-2, anti-cytochrome c, and anti-caspase-3 antibodies) overnight at 4 °C. After washes with TBST (Tris-buffered saline with 0.1% Tween-20), the blots were incubated with a secondary antibody (1:1000 in TBST with 5% skim milk) for 2 hours at room temperature. The blots were then washed and developed using the enhanced chemiluminescence detection method by a luminescence-image analyzer. Protein bands were quantified by densitometric analysis using Image J software.

3.3.8 Statistical Analysis

Each experiment was repeated three times. All results were presented as mean \pm S.E.M. Data were analyzed by one-way analysis of variance (ANOVA), followed by LSD post-hoc for comparisons between group means. Differences were considered statistically significant if p value was lesser than 0.05 ($p < 0.05$).

CHAPTER 4

RESULTS

4.1 The effect of eriodictyol on cell viability

Eriodictyol (0.001-100 μM) did not affect cell viability (Figure 8). The highest concentration of eriodictyol (100 μM) slightly decreased cell viability by 17.36% but this was not significantly different from control. Therefore, three concentrations of eriodictyol, 0.1, 1 and 10 μM , were selected for further study.





Figure 6 The effect of eriodictyol (0.001-100 μM) on cell viability in SH-SY5Y cells. Data are presented as mean \pm S.E.M of three independent experiments.

4.2 The effect of H₂O₂ on cell viability

To determine the effects of H₂O₂ on cell viability in SH-SY5Y cells, cells were exposed to H₂O₂ (100-1000 μ M) for 12 h. H₂O₂ (100-1000 μ M) significantly reduced cell viability ($p < 0.01$ VS control) (Figure 9). The IC₅₀ was 163 ± 0.12 μ M. Since H₂O₂ 200 μ M decreased cell viability to $52.56 \pm 3.60\%$, this concentration was selected to induce cytotoxicity.



CHULALONGKORN UNIVERSITY

Figure 7 The effect of H₂O₂ (100-1000 μ M) on cell viability in SH-SY5Y cells. Data are presented as mean \pm S.E.M of three independent experiments. ** $p < 0.01$ compared to control.

4.3 The protective effect of eriodictyol on H₂O₂-induced cell death

H₂O₂ (200 μ M) significantly decreased cell viability compared to control group ($p < 0.01$). Pretreatment with eriodictyol (0.1, 1 and 10 μ M) significantly increased cell viability compared to H₂O₂-treated alone ($p < 0.01$) (Figure 10), indicating the protective effect of eriodictyol.




Figure 8 The protective effects of eriodictyol on H₂O₂-induced cell death in SH-SY5Y cells. Data are presented as mean \pm SEM of three independent experiments. ** $p < 0.01$ compared to control; ## $p < 0.01$ compared to H₂O₂-treated alone.

4.4 The protective effects of eriodictyol on H₂O₂-induced ROS generation

H₂O₂ significantly increased intracellular ROS levels ($p < 0.05$) VS control. Eriodictyol pretreatment significantly decreased intracellular ROS levels induced by H₂O₂ compared to H₂O₂-treated alone ($p < 0.05$) (Figure 11).





Figure 9 The effects of eriodictyol on intracellular ROS levels induced by H₂O₂. Data are presented as mean \pm S.E.M. of three independent experiments. * $p < 0.05$ compared to control; # $p < 0.05$ compared to H₂O₂-treated alone.

4.5 The effect of eriodictyol and H₂O₂ on mitochondrial membrane potential

Mitochondrial membrane potential was determined by TMRE fluorescent intensity. Figure 12A showed the reduction of TMRE intensity in H₂O₂-treated group while eriodictyol (0.1, 1 and 10 μ M) pretreated cells increased TMRE intensity. The quantitative analysis by a microplate reader showed a significant decrease of TMRE intensity in H₂O₂-treated cells compared to control ($p < 0.05$). Eriodictyol (0.1, 1 and 10 μ M) pretreatment significantly increased TMRE intensity ($p < 0.05$ vs H₂O₂-treated alone), indicating the protection of mitochondria dysfunction (Figure 12B).



B





Figure 10 The effects of eriodictyol and H_2O_2 on mitochondrial membrane potential presented as TMRE fluorescent imaging (A) and TMRE fluorescent intensity (B). The fluorescent intensity are presented as mean \pm S.E.M. of three independent experiments. * $p < 0.05$ compared to control; # $p < 0.05$ compared to H_2O_2 -treated alone.

4.6 The effects of eriodictyol and H₂O₂ on mitochondria-mediated apoptosis cascade

Mitochondria-mediated apoptosis cascade was determined by protein expression levels of a proapoptotic protein, Bcl-2, an anti-apoptotic protein, Bax, and the markers of apoptosis, cytochrome c, cleaved caspase-3 and caspase-3. H₂O₂ significantly reduced ratio of Bcl-2/BAX expression ($p < 0.05$) compared to the control while eriodictyol pretreatment significantly increased ratio of Bcl-2/BAX expression compared to H₂O₂-treated alone ($p < 0.05$) (Figure 13). Moreover, H₂O₂ increased cytochrome c expression ($p < 0.05$ vs control) while eriodictyol pretreatment significantly decreased cytochrome c expression ($p < 0.05$ vs H₂O₂-treated alone) (Figure 14). In addition, H₂O₂ increased caspase-3 and cleaved caspase-3 expression levels ($p < 0.05$ vs control) while eriodictyol pretreatment significantly decreased caspase-3 and cleaved caspase-3 expression levels ($p < 0.05$ vs H₂O₂-treated alone) (Figure 15).

A



B





Figure 11 The effects of eriodictyol and H_2O_2 on Bcl-2 and BAX protein expression (A). Bcl-2/BAX ratio of each experimental groups were calculated and presented as mean \pm S.E.M. of three independent experiments (B). * $p < 0.05$ compared to control; # $p < 0.05$ compared to H_2O_2 -treated alone.

A



B






Figure 12 The effects of eriodictyol and H₂O₂ on cytochrome c expression (A). The expression levels were calculated and presented as cytochrome c/GAPDH of three independent experiment (B). **p* < 0.05 compared to control; #*p* < 0.05 compared to H₂O₂-treated alone.

A



B



C




Figure 13 The effects of eriodictyol and H_2O_2 on caspase-3 and cleaved caspase-3 expression (A). The expression levels were calculated and presented as caspase-3/GAPDH (B) and cleaved caspase-3/GAPDH of three independent experiment (C). * $p < 0.05$ compared to control; # $p < 0.05$ compared to H_2O_2 -treated alone.

CHAPTER 5

DISCUSSION AND CONCLUSION

Oxidative stress and mitochondrial dysfunction are the important causes of neuronal death in neurodegenerative disorders (Kim et al., 2015). The excessive ROS generation and the depletion of the antioxidant system result in oxidative stress in neurons (Kwon et al., 2014). Therefore, the neuroprotective approaches in the treatment of neurodegenerative disorders, such as a decrease of ROS production, an increase of antioxidant system and the protection of mitochondria function have gained more interested. In this study, pretreatment of eriodictyol can increase cell viability in H₂O₂-treated cells as concentration, indicating the protective effect of eriodictyol in H₂O₂-induced cytotoxicity in SH-SY5Y cells.

The protective effect of eriodictyol was concentration-independent as similar percentage cell viability was observed in cells pretreated with 0.1, 1 and 10 μ M of eriodictyol (Figure 8). This might be due to the maximum effect of eriodictyol presented at a concentration of 0.1 μ M. Either pretreatment with the lower concentrations of eriodictyol or increasing H₂O₂ concentration can be used to clarify this point. In addition, as resazurin assay detects mitochondrial reductase activity in living cells (Riss et al., 2004), this assay cannot distinguish between apoptosis and necrosis cell death. Using Annexin V-FITC and PI staining together with flow cytometry can elucidate whether the maximum effect of eriodictyol is caused by the maximum protection of either death pathway.

Flavonoids exert a direct ROS scavenging effect by reacting with $\bullet\text{O}_2^-$, HO_2^+ , nitric oxide (NO \cdot), alkoxyl and peroxy radicals (Bors et al., 1990). The ROS scavenging-mediated antioxidant activity of flavonoid is primarily attributed to the benzene ring-bound hydroxyl groups that are capable of donating either one hydrogen atom or a single electron to the ROS, and consequently stabilize ROS (Amic et al., 2007; Bors et al., 1990). Scavenging activity of eriodictyol was reported in the *in vitro* DPPH assay (Chanput et al., 2016). Taken together, eriodictyol, as a flavonoid, can reduce intracellular ROS partly due to its direct scavenging effect. Previous studies also demonstrated that eriodictyol activated Nrf2 transcription factor (Johnson et al., 2009; Xie et al., 2017). Nrf2 is responsible for the induction of various genes encoding the protective antioxidant proteins, and thus promoting the regulation of an optimal intracellular

redox environment (Ma, 2013). Therefore, the protective mechanisms of eriodictyol involving the increase of antioxidant enzyme expression should be further investigated.

Mitochondria dysfunction has been implicated in neurodegenerative diseases as the reductions of mitochondria complex I and IV activity have been reported in patients with Alzheimer's disease and Parkinson's disease, respectively (Sullivan & Brown, 2005). Moreover, mitochondria complex I inhibitors, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), and rotenone have been used to produce parkinsonism behaviors in animal models (Betarbet & Greenamyre, 2008; Bové et al., 2005). Excessive ROS generation directly impairs mitochondrial function as it causes an decrease of Bcl-2/BAX ratio resulting in the impairment of mitochondria membrane potential and the decrease of adenosine triphosphate (ATP) levels (Alvarino et al., 2017; Redza-Dutordoir & Averill-Bates, 2016; Waseem et al., 2017). In addition, ROS induces Bcl-2 down-regulation via the ubiquitin-proteasomal degradation pathway (Chanvorachote et al., 2009). This study showed that H₂O₂ markedly decreased Bcl-2/BAX ratio, and consequently decreased mitochondria membrane potential while pretreatment with eriodictyol can reverse these events. These results suggest that the mitochondria protection effect of eriodictyol is partly due to a decrease of an intracellular ROS. Further study should focus on the direct effect of eriodictyol on mitochondria function, for example the protection of mitochondria electron transport complex and the effect on mitophagy.

The decrease of mitochondria membrane potential causes cytochrome c release, and consequently induces apoptosis cascade (Green & Kroemer, 1998). Cytochrome c and caspase-3 expressions were used as indicators of mitochondria-mediated apoptosis in this study, H₂O₂ increased cytosolic cytochrome c, caspase-3 and cleaved caspase-3 expression while pretreatment with eriodictyol decreased the expressions of proteins. This result is in agreement with previous study showing that eriodictyol decreased pro-caspase-3 and cytochrome c expression in human keratinocyte HaCaT cells (Lee et al., 2011).

In conclusion, eriodictyol protect H_2O_2 -induce cell death by reducing an intracellular ROS leading to the inhibition of mitochondria-mediated apoptosis cascade (Figure 16). The effect is partly due to the direct ROS scavenging effect of eriodictyol. The effect of eriodictyol on the increase of cellular defensive mechanism and the protection of mitochondria should be further investigated. Overall, this study suggests the possible benefit of eriodictyol in the protection neuronal death in neurodegenerative disorders.




Figure 14 The action mechanisms of eriodictyol on H_2O_2 -induced cell death in SH-SY5Y cells

REFERENCES

- Acevedo-Torres, K., Berrios, L., Rosario, N., Dufault, V., Skatchkov, S., Eaton, M. J., . . . Ayala-Torres, S. (2009). Mitochondrial DNA damage is a hallmark of chemically induced and the R6/2 transgenic model of Huntington's disease. *DNA Repair (Amst)*, 8(1), 126-136. doi:10.1016/j.dnarep.2008.09.004
- Alvarino, R., Alonso, E., Tribalat, M. A., Gegunde, S., Thomas, O. P., & Botana, L. M. (2017). Evaluation of the Protective Effects of Sarains on H₂O₂-Induced Mitochondrial Dysfunction and Oxidative Stress in SH-SY5Y Neuroblastoma Cells. *Neurotox Res*, 32(3), 368-380. doi:10.1007/s12640-017-9748-3
- Amic, D., Davidovic-Amic, D., Beslo, D., Rastija, V., Lucic, B., & Trinajstic, N. (2007). SAR and QSAR of the antioxidant activity of flavonoids. *Curr Med Chem*, 14(7), 827-845.
- Ansari, M. A., & Scheff, S. W. (2010). Oxidative stress in the progression of Alzheimer disease in the frontal cortex. *J Neuropathol Exp Neurol*, 69(2), 155-167. doi:10.1097/NEN.0b013e3181cb5af4
- Betarbet, R., & Greenamyre, J. T. (2008). Chapter 14 - Complex I Inhibition, Rotenone and Parkinson's Disease. In R. Nass & S. Przedborski (Eds.), *Parkinson's Disease* (pp. 195-206). San Diego: Academic Press.
- Birben, E., Sahiner, U. M., Sackesen, C., Erzurum, S., & Kalayci, O. (2012). Oxidative stress and antioxidant defense. *World Allergy Organ J*, 5(1), 9-19. doi:10.1097/WOX.0b013e3182439613
- Blesa, J., Trigo-Damas, I., Quiroga-Varela, A., & Jackson-Lewis, V. R. (2015). Oxidative stress and Parkinson's disease. *Front Neuroanat*, 9, 91. doi:10.3389/fnana.2015.00091
- Bors, W., Heller, W., Michel, C., & Saran, M. (1990). Flavonoids as antioxidants: determination of radical-scavenging efficiencies. *Methods Enzymol*, 186, 343-355.
- Bové, J., Prou, D., Perier, C., & Przedborski, S. (2005). Toxin-Induced Models of Parkinson's Disease. *NeuroRX*, 2(3), 484-494. doi:<https://doi.org/10.1602/neurorx.2.3.484>
- Cai, L., Wang, H., Li, Q., Qian, Y., & Yao, W. (2008). Salidroside inhibits H₂O₂-induced apoptosis in PC12 cells by preventing cytochrome c release and inactivating of caspase cascade. *Acta Biochim Biophys Sin (Shanghai)*, 40(9), 796-802.

- Chanput, W., Krueyos, N., & Ritthiruangdej, P. (2016). Anti-oxidative assays as markers for anti-inflammatory activity of flavonoids. *Int Immunopharmacol*, 40, 170-175.
doi:10.1016/j.intimp.2016.08.038
- Chanvorachote, P., Pongrakhananon, V., Wannachaiyasit, S., Luanpitpong, S., Rojanasakul, Y., & Nimmannit, U. (2009). Curcumin sensitizes lung cancer cells to cisplatin-induced apoptosis through superoxide anion-mediated Bcl-2 degradation. *Cancer Invest*, 27(6), 624-635.
doi:10.1080/07357900802653472
- Circu, M. L., & Aw, T. Y. (2010). Reactive oxygen species, cellular redox systems, and apoptosis. *Free Radic Biol Med*, 48(6), 749-762. doi:10.1016/j.freeradbiomed.2009.12.022
- Cook, S. A., Sugden, P. H., & Clerk, A. (1999). Regulation of bcl-2 family proteins during development and in response to oxidative stress in cardiac myocytes: association with changes in mitochondrial membrane potential. *Circ Res*, 85(10), 940-949.
- David, D. C., Hauptmann, S., Scherping, I., Schuessel, K., Keil, U., Rizzu, P., . . . Gotz, J. (2005). Proteomic and functional analyses reveal a mitochondrial dysfunction in P301L tau transgenic mice. *J Biol Chem*, 280(25), 23802-23814. doi:10.1074/jbc.M500356200
- de Oliveira, M. R., de Bittencourt Brasil, F., & Furstenu, C. R. (2018). Sulforaphane Promotes Mitochondrial Protection in SH-SY5Y Cells Exposed to Hydrogen Peroxide by an Nrf2-Dependent Mechanism. *Mol Neurobiol*, 55(6), 4777-4787. doi:10.1007/s12035-017-0684-2
- Elmore, S. (2007). Apoptosis: a review of programmed cell death. *Toxicol Pathol*, 35(4), 495-516.
doi:10.1080/01926230701320337
- Feng, C., Luo, T., Zhang, S., Liu, K., Zhang, Y., Luo, Y., & Ge, P. (2016). Lycopene protects human SHSY5Y neuroblastoma cells against hydrogen peroxide-induced death via inhibition of oxidative stress and mitochondria-associated apoptotic pathways. *Mol Med Rep*, 13(5), 4205-4214. doi:10.3892/mmr.2016.5056
- Gomes, A., Fernandes, E., & Lima, J. L. F. C. (2005). Fluorescence probes used for detection of reactive oxygen species. *Journal of Biochemical and Biophysical Methods*, 65(2), 45-80.
doi:<https://doi.org/10.1016/j.jbbm.2005.10.003>
- Green, D., & Kroemer, G. (1998). The central executioners of apoptosis: caspases or mitochondria? *Trends Cell Biol*, 8(7), 267-271.
- Hu, X. L., Niu, Y. X., Zhang, Q., Tian, X., Gao, L. Y., Guo, L. P., . . . Zhao, Q. C. (2015).

- Neuroprotective effects of Kukoamine B against hydrogen peroxide-induced apoptosis and potential mechanisms in SH-SY5Y cells. *Environ Toxicol Pharmacol*, 40(1), 230-240. doi:10.1016/j.etap.2015.06.017
- Ichim, G., & Tait, S. W. (2016). A fate worse than death: apoptosis as an oncogenic process. *Nat Rev Cancer*, 16(8), 539-548. doi:10.1038/nrc.2016.58
- Jing, X., Shi, H., Zhu, X., Wei, X., Ren, M., Han, M., . . . Lou, H. (2015). Eriodictyol Attenuates beta-Amyloid 25-35 Peptide-Induced Oxidative Cell Death in Primary Cultured Neurons by Activation of Nrf2. *Neurochem Res*, 40(7), 1463-1471. doi:10.1007/s11064-015-1616-z
- Johnson, J., Maher, P., & Hanneken, A. (2009). The flavonoid, eriodictyol, induces long-term protection in ARPE-19 cells through its effects on Nrf2 activation and phase 2 gene expression. *Invest Ophthalmol Vis Sci*, 50(5), 2398-2406. doi:10.1167/iovs.08-2088
- Kim, G. H., Kim, J. E., Rhie, S. J., & Yoon, S. (2015). The Role of Oxidative Stress in Neurodegenerative Diseases. *Exp Neurobiol*, 24(4), 325-340. doi:10.5607/en.2015.24.4.325
- Kwon, S.-H., Ma, S.-X., Lee, S.-Y., & Jang, C.-G. (2014). Sulfuretin inhibits 6-hydroxydopamine-induced neuronal cell death via reactive oxygen species-dependent mechanisms in human neuroblastoma SH-SY5Y cells. *Neurochemistry International*, 74, 53-64. doi:<https://doi.org/10.1016/j.neuint.2014.04.016>
- Lee, E. R., Kim, J. H., Choi, H. Y., Jeon, K., & Cho, S. G. (2011). Cytoprotective effect of eriodictyol in UV-irradiated keratinocytes via phosphatase-dependent modulation of both the p38 MAPK and Akt signaling pathways. *Cell Physiol Biochem*, 27(5), 513-524. doi:10.1159/000329973
- Lee, J. E., Sohn, J., Lee, J. H., Lee, K. C., Son, C. S., & Tockgo, Y. C. (2000). Regulation of bcl-2 family in hydrogen peroxide-induced apoptosis in human leukemia HL-60 cells. *Exp Mol Med*, 32(1), 42-46. doi:10.1038/emm.2000.8
- Lou, H., Jing, X., Ren, D., Wei, X., & Zhang, X. (2012). Eriodictyol protects against H₂O₂-induced neuron-like PC12 cell death through activation of Nrf2/ARE signaling pathway. *Neurochem Int*, 61(2), 251-257. doi:10.1016/j.neuint.2012.05.013
- Ma, Q. (2013). Role of nrf2 in oxidative stress and toxicity. *Annu Rev Pharmacol Toxicol*, 53, 401-426. doi:10.1146/annurev-pharmtox-011112-140320
- Murphy, M. P. (2009). How mitochondria produce reactive oxygen species. *Biochem J*, 417(1), 1-

13. doi:10.1042/bj20081386

Park, H. R., Lee, H., Park, H., Jeon, J. W., Cho, W. K., & Ma, J. Y. (2015). Neuroprotective effects of *Liriope platyphylla* extract against hydrogen peroxide-induced cytotoxicity in human neuroblastoma SH-SY5Y cells. *BMC Complement Altern Med*, 15, 171.

doi:10.1186/s12906-015-0679-3

Patten, D. A., Germain, M., Kelly, M. A., & Slack, R. S. (2010). Reactive oxygen species: stuck in the middle of neurodegeneration. *J Alzheimers Dis*, 20 Suppl 2, S357-367. doi:10.3233/jad-2010-100498

Redza-Dutordoir, M., & Averill-Bates, D. A. (2016). Activation of apoptosis signalling pathways by reactive oxygen species. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*, 1863(12), 2977-2992. doi:<https://doi.org/10.1016/j.bbamcr.2016.09.012>

Riss, T. L., Moravec, R. A., Niles, A. L., Duellman, S., Benink, H. A., Worzella, T. J., & Minor, L. (2004). Cell Viability Assays. In G. S. Sittampalam, N. P. Coussens, K. Brimacombe, A. Grossman, M. Arkin, D. Auld, C. Austin, J. Baell, B. Bejcek, J. M. M. Caaveiro, T. D. Y. Chung, J. L. Dahlin, V. Devanaryan, T. L. Foley, M. Glicksman, M. D. Hall, J. V. Haas, J. Inglese, P. W. Iversen, S. D. Kahl, S. C. Kales, M. Lal-Nag, Z. Li, J. McGee, O. McManus, T. Riss, O. J. Trask, Jr., J. R. Weidner, M. J. Wildey, M. Xia, & X. Xu (Eds.), *Assay Guidance Manual*. Bethesda (MD): Eli Lilly & Company and the National Center for Advancing Translational Sciences.

Sasaki, N., Toda, T., Kaneko, T., Baba, N., & Matsuo, M. (2003). Protective effects of flavonoids on the cytotoxicity of linoleic acid hydroperoxide toward rat pheochromocytoma PC12 cells. *Chem Biol Interact*, 145(1), 101-116.

Sies, H. (2017). Hydrogen peroxide as a central redox signaling molecule in physiological oxidative stress: Oxidative eustress. *Redox Biol*, 11, 613-619. doi:10.1016/j.redox.2016.12.035

Sullivan, P. G., & Brown, M. R. (2005). Mitochondrial aging and dysfunction in Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry*, 29(3), 407-410. doi:10.1016/j.pnpbp.2004.12.007

Tanagornmeatar, K., Chaotham, C., Sritularak, B., Likhitwitayawuid, K., & Chanvorachote, P. (2014). Cytotoxic and anti-metastatic activities of phenolic compounds from *Dendrobium ellipsophyllum*. *Anticancer Res*, 34(11), 6573-6579.

- Thanan, R., Oikawa, S., Hiraku, Y., Ohnishi, S., Ma, N., Pinlaor, S., . . . Murata, M. (2014). Oxidative stress and its significant roles in neurodegenerative diseases and cancer. *Int J Mol Sci*, 16(1), 193-217. doi:10.3390/ijms16010193
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol*, 39(1), 44-84. doi:10.1016/j.biocel.2006.07.001
- Wang, X., Wang, W., Li, L., Perry, G., Lee, H. G., & Zhu, X. (2014). Oxidative stress and mitochondrial dysfunction in Alzheimer's disease. *Biochim Biophys Acta*, 1842(8), 1240-1247. doi:10.1016/j.bbadis.2013.10.015
- Waseem, M., Sahu, U., Salman, M., Choudhury, A., Kar, S., Tabassum, H., & Parvez, S. (2017). Melatonin pre-treatment mitigates SHSY-5Y cells against oxaliplatin induced mitochondrial stress and apoptotic cell death. *PLoS One*, 12(7), e0180953. doi:10.1371/journal.pone.0180953
- Xie, G., Meng, X., Wang, F., Bao, Y., & Huo, J. (2017). Eriodictyol attenuates arsenic trioxide-induced liver injury by activation of Nrf2. *Oncotarget*, 8(40), 68668-68674. doi:10.18632/oncotarget.19822
- Xu, C. L., Qu, R., Zhang, J., Li, L. F., & Ma, S. P. (2013). Neuroprotective effects of madecassoside in early stage of Parkinson's disease induced by MPTP in rats. *Fitoterapia*, 90, 112-118. doi:10.1016/j.fitote.2013.07.009
- Yang, R., Wei, L., Fu, Q. Q., Wang, H., You, H., & Yu, H. R. (2016). SOD3 Ameliorates H₂O₂-Induced Oxidative Damage in SH-SY5Y Cells by Inhibiting the Mitochondrial Pathway. *Neurochem Res*, 41(7), 1818-1830. doi:10.1007/s11064-016-1897-x
- Zhu, G. F., Guo, H. J., Huang, Y., Wu, C. T., & Zhang, X. F. (2015). Eriodictyol, a plant flavonoid, attenuates LPS-induced acute lung injury through its antioxidative and anti-inflammatory activity. *Exp Ther Med*, 10(6), 2259-2266. doi:10.3892/etm.2015.2827

VITA

NAME	Saowarose Thongin
DATE OF BIRTH	24 January 1994
PLACE OF BIRTH	Saraburi
INSTITUTIONS ATTENDED	Bachelor of Applied Thai Traditional Medicine
HOME ADDRESS	88/14-15 Subantad Road T. Pakpiew Muang Saraburi 18000
PUBLICATION	Thongin S, Teerapattarakon N, Benya-aphikul H, Pongrakhananon V, Sritularak B, Rodsiri R. Eriodictyol attenuates hydrogen peroxide-induced cell death in neuronal SH-SY5Y cells. (Submitted)
AWARD RECEIVED	The Best Poster Presentation Award (Molecular Research) at the 41st Pharmacological and Therapeutic Society of Thailand Meeting "Herbal and Complementary Medicine Innovation: East Meets West"