

12-1-2000

Characteristics and attitude of contraceptive acceptors among adolescents attending the Family Planning clinic at King Chulalongkorn Memorial Hospital

S. Bunyavejchevin

W. Wisawasukmongchol

U. Jaisamran

D. Reinprayoon

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Bunyavejchevin, S.; Wisawasukmongchol, W.; Jaisamran, U.; and Reinprayoon, D. (2000) "Characteristics and attitude of contraceptive acceptors among adolescents attending the Family Planning clinic at King Chulalongkorn Memorial Hospital," *Chulalongkorn Medical Journal*: Vol. 44: Iss. 12, Article 4. Available at: <https://digital.car.chula.ac.th/clmjjournal/vol44/iss12/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Characteristics and attitude of contraceptive acceptors among adolescents attending the Family Planning clinic at King Chulalongkorn Memorial Hospital

Suvit Bunyavejchevin * Wirach Wisawasukmongchol *

Unnop Jaisamran * Damrong Reinprayoon *

Bunyavejchevin S, Wisawasukmongchol W, Jaisamran U, Reinprayoon D. Characteristics and attitude of contraceptive acceptors among adolescents attending the Family Planning clinic at King Chulalongkorn Memorial Hospital. Chula Med J 2000 Dec; 44(12): 917 - 25

Objective : *To study the characteristics, choice and attitude towards selected contraceptive methods of contraception among female adolescents attending the Family Planning clinic at King Chulalongkorn Memorial Hospital, Bangkok.*

Design : *Descriptive study.*

Materials and methods : *200 female contraceptive acceptors aged 10 -19 years attending a Family planning clinic at King Chulalongkorn Memorial Hospital during January 1995 to December 1996 were randomly selected and interviewed by trained nurses about their attitude towards the selected contraceptive methods. Routine gynecological outpatient data were recorded.*

Results : *The mean age was 15.50 ± 1.55 years. Most of them were housewives (50 per cent) and factory workers (48 per cent). Fifty-eight per cent had graduated from primary school. The contraceptive choices were pills (43.0 per cent), injectables*

(38.0 per cent), intrauterine devices (12.0 per cent), subdermal implants (5.0 per cent), condoms (1.5 per cent), and tubal sterilization (0.5 per cent). The main reason for choice of the method was the confidence in the effectiveness of the contraception. Most of them obtained contraceptive information from friends and relatives, but chose the methods by themselves. Most of them had no premarital sex education (56.5 per cent).

Conclusion : Adolescent acceptors made the contraceptive decision by themselves using their own available information. Contraceptive counseling should be directed to the acceptors themselves and adjusted to the background of the individual. Special education programs about sex and contraception should be encouraged.

Key words : Adolescent, Contraception.

Reprint request : Bunyavejchevin S, Department of Obstetrics and Gynecology, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. July 5, 2000.

สุวิทย์ บุญยะเวชชีวิน, วิรัช วิศวสุขมงคล, อรรณพ ใจสำราญ, ดำรง เจริญประยูร. ลักษณะ
และทัศนคติต่อการคุมกำเนิดในผู้รับบริการวัยรุ่นที่มารับบริการที่คลินิกวางแผนครอบครัว
โรงพยาบาลจุฬาลงกรณ์. จุฬาลงกรณ์เวชสาร 2543 ธ.ค; 44(12): 917 - 25

วัตถุประสงค์ : เพื่อศึกษาลักษณะของผู้รับบริการและทัศนคติต่อวิธีการคุมกำเนิดที่เลือกใน
วัยรุ่นที่มารับบริการที่คลินิกวางแผนครอบครัว โรงพยาบาลจุฬาลงกรณ์

รูปแบบการศึกษา : วิจัยเชิงพรรณนา

วัตถุประสงค์และวิธีการ : ทำการเก็บข้อมูลทั่วไปและข้อมูลทางนรีเวชในผู้รับบริการอายุระหว่าง
10 -19 ปี ที่มารับบริการที่คลินิกวางแผนครอบครัว โรงพยาบาลจุฬาลงกรณ์
ระหว่างเดือน มกราคม พ.ศ. 2548 ถึง เดือนธันวาคม พ.ศ. 2549 และทำ
การสุ่มเลือก ผู้รับบริการจำนวน 200 คน เพื่อสอบถามถึงทัศนคติต่อวิธีการ
คุมกำเนิดที่เลือกใช้โดยพยาบาลที่ฝึกอบรมพิเศษ

ผลการศึกษา : อายุเฉลี่ยของผู้รับบริการคือ 15.50 ± 1.55 ปี ส่วนใหญ่ผู้รับบริการมีอาชีพ
แม่บ้าน (ร้อยละ 50) และทำงานโรงงาน (ร้อยละ 48) มีการศึกษาระดับประถม
ต้น (ร้อยละ 58) เลิกยาเม็ดคุมกำเนิด (ร้อยละ 43.0) ยาฉีดคุมกำเนิด
(ร้อยละ 38.0) ห่วงอนามัย (ร้อยละ 12.0) ยาฝังคุมกำเนิด (ร้อยละ 5.0)
ถุงยางอนามัย (ร้อยละ 1.5) และทำหมัน (ร้อยละ 0.5) สาเหตุที่เลือกวิธี
คุมกำเนิดคือเชื่อมั่นในประสิทธิภาพของวิธีคุมกำเนิด ได้รับความรู้จากเพื่อน
และญาติ แต่ส่วนใหญ่ผู้รับบริการจะตัดสินใจด้วยตนเอง ส่วนใหญ่ไม่ได้รับ
ความรู้เรื่องเพศก่อนแต่งงาน (ร้อยละ 56.5)

สรุปผลการศึกษา : ส่วนใหญ่ของผู้รับบริการจะตัดสินใจด้วยตนเอง โดยข้อมูลของตัวเอง การ
ให้คำปรึกษาในวัยรุ่นควรจะมีตรงที่ตัวผู้รับบริการ และปรับตามลักษณะ
พื้นฐานของผู้รับบริการ ควรเน้นที่การให้ความรู้เรื่องเพศศึกษาและการคุม
กำเนิดในผู้รับบริการกลุ่มนี้

Young people, aged 10 to 19 years, number more than one billion, comprise nearly one-fifth of the world population and are growing in number.⁽¹⁾ Projections indicate that the number of young adults in Southeast Asian countries will increase by over 20 % over the next 15 years.^(1,2) In many parts of the world, premarital sexuality is common among young people.⁽³⁾ In Asia, data from Hongkong, South Korea, and Thailand show that fewer than 10% of unmarried women under the age of 24 have experienced intercourse.⁽⁴⁾ In Thailand, boys reported having first sexual intercourse by age 18, often with commercial sex workers.⁽⁵⁾ One study show that many sexually active youths have had their first experience of intercourse by the at age of 13 years or younger.⁽⁶⁾ Adolescent relationships tend to be more ephemeral than at older age and generally constitute a serial monogamy.⁽⁷⁾ Early sexual intercourse results in high rates of adolescent pregnancy and abortion , as well as in increased risk of sexually transmitted disease (STD).⁽⁸⁻¹⁰⁾ In India and Bangladesh, sexually active young people are less likely to use contraception than adults, even within marriage.^(11,12) In Africa, there is about 8% of unmet need for contraception in unmarried women aged 15 to 19.⁽¹³⁾ In Ecuador, there is low use of contraception at the first intercourse (11% in females and 15 % in males).⁽¹⁴⁾ Even in United Kingdom, 46 per cent (of non-virgins) were found to have had intercourse without contraception and 18 % used condoms 'rarely' or 'never'.⁽¹⁵⁾

In Thailand, the adolescent pregnancy rate is 52 per 1,000 births and the percentage of women in 1987 who ever used contraception during adolescents is 62 %.⁽¹¹⁾ In a study of Thai adolescents, 41 % of males reported using condoms during sexual

intercourse while only 23 % in females reported using condoms.⁽¹⁶⁾ Up to now there has been a lack of information in Thailand about the choices of contraception and the attitude of adolescents to the selected methods. The present study was conducted to examine the characteristics and attitude of adolescents who seeked contraceptive services in King Chulalongkorn Memorial Hospital.

Materials and Methods

From January 1995 to December 1996, 200 contraceptive acceptors aged 10-19 years attending the Family Planning clinic at King Chulalongkorn Memorial Hospital were randomly selected into the study. They were interviewed by well-trained nurses about their attitudes towards the selected method, history of illegal abortion, premarital sex education and their sources of information about contraceptive methods. Demographic and gynecological data were also obtained.

Results

The mean age of the contraceptive acceptors was 15.50 years (Table 1). Most of these attended for a postpartum check up and contraceptive service (67 %) and the remainder came for contraceptive service only (23 %). Pills and injectables were the two most commonly chosen techniques (43 % and 38 %) (Table 2). Most women got contraceptive information from friends or relatives (68.0 %) (Table 2). Most acceptors decided on the method by themselves (150,75.0 %) and only 24 (12.0 %) of acceptors depended on their husbands' decision. (Table 3).

One hundred and two (51%) of women reported a history of ever-use of contraception. Ninety

Table 1. Acceptors' characteristics.

	N = 2240
	Mean ± SD
Age (years)	15.50 ± 1.55
Weight (kgs)	49.5 ± 22.00
Height (cms)	150.5 ± 21.50
	N (%)
<u>Educational levels</u>	
Primary school	116 (58)
Secondary school	40 (20)
Vocational	40 (20)
Bachelor degree or higher	4 (2)
<u>Occupation</u>	
House wife	100 (50)
Employee	56 (48)
Business	2 (1)
Student	2 (1)
<u>Marital status</u>	
Single	4 (2.0)
Married	195 (97.5)
Widow/Divorce	1 (0.5)

(45.5 %) acceptors desired a change of methods from pill to the other methods. The most common reason was forgetting to take the pills (52.0 %). Twenty acceptors (10 %) reported they had used the postcoital contraception, most of which were combined pill (75 %). One hundred and nine (54.5 %) acceptors had husband accompanied them to the hospital. Eighty-seven (43.5 %) of acceptors reported receiving sex education before marriage, mainly from their school (89.6 %) (Table 4)

Eighteen (9 %) of acceptors reported a previous illegal abortion. Most were not married at that time and had undergone the operation at a

the private clinic (66.7 %) with a cost of US \$ 41-120 (83.3 %) and the cost of less than US \$40 (16.7 %). Thirty nine per cent of the sources of information came from friends and 33 % from advertisement. Fifteen per cent of women decided to get an abortion if their current contraceptive had failed.

Table 2. Contraceptive choices, reasons for choosing and sources of contraceptive information.

	N = 200
	n (%)
<u>Choices</u>	
Pills	86 (43.0)
Injectables	76 (38.0)
IUD	24 (12.0)
Subdermal implants	10 (5 0)
Condom	3 (1.5)
Tubal sterilization	1 (0.5)
<u>Reasons for choosing the methods</u>	
a. Belief in the efficacy	58 (29)
b. Influenced by a previous user	48 (24)
c. Less expensive	2 (1)
d. Forced by husband or other partners	24 (12)
e. Others:	
Desire no more children	6 (3)
Incorrect beliefs about methods chosen	14 (7)
Need the non -contraceptive benefit effects of pill.	10 (5)
Switching from poor compliance method (pills to another)	20 (10)
The technique chosen was accessible	16 (8)
Want to try the other methods	2 (1)
<u>Sources of contraceptive information</u>	
Friends or relatives	136 (68)
Mass media (TV or Radio or Newspapers)	8 (4)
Health personnel	38 (19)
None	18 (9)

Table 3. Person who influenced the decision.

	N = 200
	n (%)
Made the decision herself	150 (75)
Husband	24 (12)
Health personnel	4 (2)
Friends or relatives	20 (10)
Teachers	2 (1)

Table 4. Premarital sexual relationship, premarital sex education and source of information.

	(N = 200)
	n (%)
Premarital sexual relationship	
• Yes	26 (13)
• No	174 (87)
Premarital sex education (N = 200)	
• Yes	87 (43.5)
• No	113 (56.5)
Source of information (N = 87)	
- Television	2 (2.3)
- School	78 (89.5)
- Journal / Magazine	2 (2.3)
- Friend	1 (1.2)
- Hospital	3 (3.5)
- Special training program	1 (1.2)

Discussion

From this study, most of the adolescent acceptors chose pills and injectables as their contraception (Table 2). Injectables were well accepted in the study group and in other studies, most of the adolescents viewed amenorrhea as a positive feature of Depo Medroxy Progesterone Acetate (DMPA)

with a consequently high continuation rate.^(17, 18) As expected, few adolescent acceptors chose sterilization because they had enough children (more than two). Subdermal implants are a good choice for adolescents and adults,⁽¹⁹⁻²²⁾ but few women chose this method.

Few adolescents in this hospital-based study were using condoms when compared to other reports^(23,24) This group of patients coming to hospital for a postpartum contraceptive service requested more effective methods.

Although previous users and husbands influenced most acceptors, the majority decided on a method by themselves. (Table 2 and 3). Many acceptors reported forgetting to take the pills (Table 4), as observed in other studies which report a lack of access to contraceptive information and poor compliance with contraceptive recommendations among adolescents.⁽²⁵⁻²⁷⁾ Other factors that contribute to nonuse of contraceptives by adolescents include the fear of parental discovery and waiting for a closer relationship with their partners.⁽²⁷⁾ We also found evidence of poor information and compliance from our data (Table 2). The use of postcoital pills was noted in only 20 cases (10 %), and this method had the low incidence rate in many other studies.^(28, 29) More information and counseling about emergency contraception and increasing access of this to the adolescent can provide them a second chance to prevent pregnancy.⁽³⁰⁾

Husbands may play an important role in contraceptive counseling. We found 54.5% of women had husbands accompanying them. The couple's counseling programs was encouraged if they came together to the clinic. Only few acceptors had received previous premarital sex education. We have found that

a premarital counseling program which had included sex education and family planning was helpful in preventing unintended teenage pregnancy and marital problems in young couples.⁽³¹⁾

In Thailand, abortion for unintended pregnancy is illegal, but 9 % of acceptors reported prior abortion. Some acceptors would choose abortion if contraception failed. To prevent unwanted pregnancy and abortion, the adolescent should be encouraged to use the high effective methods of contraception.

Due to the limitation of the hospital-based design, generalizing the finding of this study to the general population is a problem. However, these are the first preliminary data of adolescent contraception use in Thailand, which will be useful for both acceptor care and the planning for further community-based studies.

From this study, contraceptive counseling should be adjusted to the background of individual acceptors. Education programs about sex and contraception for Thai adolescents still need to be encouraged.

Acknowledgement

The authors wish to thank Professor Ronald H. Gray, School of Public Health, Johns Hopkins University, for his advice in this study .

References

1. United Nations (UN). Department of international economic and social information and policy analysis. Population division. The Sex and Age Distribution of the World Populations: The 1994 revision. New York, United Nations, 1994: 858
2. United nations (UN) Department of international economic and Social affairs. Adolescent reproductive behavior: evidence from developing countries. (vol.1) New York, United Nations, 1988 (Population studies No.109): 187
3. Morris L. Sexual Experience and Contraceptive Use Among Young Adolescent in Central America. Presented at the symposium on Population in Central America, San Jose, Costa Rica, Oct.16-18 1995: 25
4. United Nations (UN) Department of international economic and Social affairs. Adolescent reproductive behavior: evidence from developing countries. (vol.2) New York, United Nations, MN, 1989 (Population studies No.109/ Add.1): 128
5. Xenos P, Pitaktepsombati P, Sittitrai W. Partner patterns in the sexual behavior of unmarried, rural Thai men. Asia Pacific Popul Forum 1993; 6(4): 104 - 17
6. Center for Disease Control and Prevention: Miami youth risk behavior survey, 1993 Sexual behavior and drug use among youth in dropout-prevention programs – Miami, 1994. MMWR Morb Mortal Wkly Rep 1994 Dec 2; 43(47): 873 - 6
7. Brown RT, Cromer BA. The pediatrician and the sexually active adolescent. Sexual activity and contraception. Pediatr Clin North Am 1997 Dec; 44(6): 1379 - 90
8. Kaunitz AM. Contraception for the adolescent patient. Int J Fertil Womens Med. 1997 Jan - Feb; 42(1): 30 - 8
9. Newcomer S, Baldwin W. Demographics of adolescent sexual behavior, contraception,

- pregnancy, and STDs. *J Sch Health*. 1992 Sep; 62(7): 5 - 70
10. Creatsas GK. Sexuality: Sexual activity and contraception during adolescence. *Curr Opin Obstet Gynecol* 1993 Dec; 5(6); 774 - 83
 11. International Institute for Population Sciences (IIPS). International family health survey 1992-3. Bombay, IIPS, 1995: 402
 12. Mahmud M, Islam MM. Adolescent contraceptive use and its determinants in Bangladesh: evidence from Bangladesh Fertility Survey 1989. *Contraception*. 1995 Sep; 52(3): 181 - 6
 13. Westoff CF. Age at marriage at first birth, and fertility in Africa. Washington D.C., World Bank, (World Bank Technical Paper 169) 1992; 63
 14. Eggleston E. Use of family planning at first sexual intercourse among young adults in Ecuador. *J Biosoc Sci* 1998 Oct; 30(4): 501 - 10
 15. Coleman L, Ingham R. Attenders at young people's clinics in Southampton: variations in contraceptive use. *Br J Fam Plann* 1998 Oct; 24(3): 101 - 4
 16. Sonenstein FL, Pleck JH, Ku LC. Sexual activity, condom use and AIDS awareness among adolescent males. *Fam Plann Perspect* 1989 Jul-Aug; 21(4): 152 - 8
 17. Tyrer LB. Oral contraception for the adolescent. *J Reprod Med* 1984 Jul; 29(7 Suppl): 551 - 9
 18. Davis AJ. Use of depot medroxyprogesterone acetate contraception in adolescents. *J Reprod Med* 1996 May; 41(5 Suppl): 407 - 13
 19. Polaneczky M, Slap G, Forke C, Rappaport A, Sondheimer S. The use of levonorgestrel implants (Norplant) for contraception in adolescent mothers. *N Engl J Med* 1994 Nov 3; 331(18): 1201 - 6
 20. Cromer BA, Berg-Kelly KS, Van Groningen JP, Seimer BS, Ruusuvaara L. Depot medroxyprogesterone acetate (Depo-Provera) and levonorgestrel (Norplant) use in adolescents among clinicians in Northern Europe and the United States. *J Adolesc Health* 1998 Aug; 23(2): 74 - 80
 21. Dabrow SM, Merrick CL, Conlon M. Adolescent girls' attitudes toward contraceptive subdermal implants. *J Adolesc Health* 1995 May; 16(5): 360 - 6
 22. Cullins VE, Remsburg RE, Blumenthal PD, Huggins GR. Comparison of adolescent and adult experiences with Norplant levonorgestrel contraceptive implants. *Obstet Gynecol* 1994 Jun; 83(6): 1026 - 32
 23. Harbin RE. Female adolescent contraception. *Pediatr Nurs* 1995 May-Jun; 21(3): 221 - 6
 24. Polaneczky M. Adolescent contraception. *Curr Opin Obstet Gynecol* 1998 Jun; 10(3): 213 - 9
 25. Goldstuck ND, Hammar E, Butchart A. Use and misuse of oral contraceptives by adolescents attending a free-standing clinic. *Adv Contracept* 1987 Dec; 3(4): 335 - 9
 26. Davis AJ. The role of hormonal contraception in adolescents. *Am J Obstet Gynecol* 1994 May; 170(5 pt 2): 1581 - 5
 27. Zabin LS, Stark HA, Emerson MR. Reason for delay in contraceptive clinic utilization. Adolescent clinic and nonclinic populations compared. *J Adolesc Health Care* 1991 May; 12(3): 225 - 32
 28. Gold MA, Schein A, Coupey SM. Emergency contraception: a national survey of adolescent

- health experts. Fam Plann Perspect 1997 Jan-Feb; 29(1): 15 - 9
29. Seamark CJ, Pereira Gray DJ. Teenage's use of emergency contraception in a general practice. J R Soc Med 1997 May; 90(8): 443-4
30. Gold MA. Emergency contraception: a second chance at preventing adolescent unintended pregnancy. Curr Opin Pediatr 1997 Aug; 9(4): 300-9
31. Reinprayoon D, Bunyavejchevin S. Premarital counseling clinic at Chulalongkorn Hospital. J Med Assoc Thai 1998 Dec; 81(12): 993 - 7