

7-1-2006

Correlation of tumor grading and cellular proliferation in colorectal adenocarcinomas

M. Tawevisit

N. Chirakalwasan

M. Chaipipat

U. Phumsuk

N. Wisedopas

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Tawevisit, M.; Chirakalwasan, N.; Chaipipat, M.; Phumsuk, U.; and Wisedopas, N. (2006) "Correlation of tumor grading and cellular proliferation in colorectal adenocarcinomas," *Chulalongkorn Medical Journal*: Vol. 50: Iss. 7, Article 2.

Available at: <https://digital.car.chula.ac.th/clmjjournal/vol50/iss7/2>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Correlation of tumor grading and cellular proliferation in colorectal adenocarcinomas

Mana Taweevisit*

Narida Chirakalwasan* Mookda Chaipipat*

Ubon Phumsuk*

Naruemon Wisedopas*

Taweevisit M, Chirakalwasan N, Chaipipat M, Phumsuk U, Wisedopas N. Correlation of tumor grading and cellular proliferation in colorectal adenocarcinomas. Chula Med J 2006 Jul; 50(7): 461 - 70

Problem/Background : *Several values including patient status, tumor biomarkers and molecular aspect are used as predictors in the patient follow-up and management of colorectal cancer. Tumor grading or differentiation is a prognostic factor which is practically identifiable in routine histologic sections. However, decision making of histologic grade in some cases is difficult.*

Objective : *To study the correlation between cell proliferation, using Ki67 immunohistochemical staining, and grading of colorectal adenocarcinoma.*

Design : *Retrospective analytic study.*

Setting : *Department of Pathology, Faculty of Medicine, Chulalongkorn University*

Materials and Methods : *One hundred and sixty-two cases, first diagnosed as colorectal adenocarcinoma, were recruited in our study. Tumor grading was classified into well-differentiated, moderately-differentiated, poorly-differentiated and undifferentiated type, reviewing on hematoxylin and eosin stained slides in each case, according to the World Health Organization's criteria 2000. Neoplastic cell proliferation is determined by Ki67 immunostain. The correlation was analyzed by means of the Spearman's rank correlation coefficient.*

- Results** : *Correlation between neoplastic cell proliferation and tumor differentiation of colonic carcinoma was found statistically significant ($P = 0.014$).*
- Conclusions** : *This study demonstrated a significant correlation between cell proliferation and differentiation of colorectal cancer. The poorer tumor differentiation is, the higher is the proliferation rate. The Ki67 may have a role in some tumors, which are difficult to grade in routine histologic sections.*
- Keywords** : *Cellular proliferation, Colorectal cancer, Tumor differentiation.*

Reprint request : Tawevisit M. Department of Pathology, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. December 16, 2005.

มานะ ทวีวิศิษฐ์, นฤตา จิรกาลวสาน, มุกดา ชัยพิพัฒน์, อุบล พุ่มสุข, นฤมล วิเศษโอภาส.
ความสัมพันธ์ระหว่างเกรดและการแบ่งตัวของเซลล์ในมะเร็งลำไส้ใหญ่. จุฬาลงกรณ์เวชสาร
2549 ก.ค; 50(7): 461 - 70

เหตุผลของการทำวิจัย : ค่าต่าง ๆ ไม่ว่าจะเป็นสภาวะของผู้ป่วย รวมถึงตัวชี้วัดทางชีวภาพ และเชิงโมเลกุล ถูกนำมาใช้เพื่อติดตามผลการรักษาผู้ป่วยมะเร็งลำไส้ใหญ่เพื่อการดูแลผู้ป่วยอย่างมีประสิทธิภาพ เกรด หรือการเปลี่ยนแปลงของมะเร็งก็ถูกนำมาใช้ในการทำนายโรคด้วยเช่นกัน โดยอาศัยการตรวจจากสไลด์เนื้อเยื่อ ถึงกระนั้น ในเนื้อมะเร็งบางราย ก็เป็นการยากที่จะตัดสินใจในการให้เกรดของมัน

วัตถุประสงค์ : เพื่อศึกษาความสัมพันธ์ระหว่างอัตราการแบ่งตัวกับการเปลี่ยนแปลงของเซลล์มะเร็งลำไส้ใหญ่

รูปแบบการวิจัย : การศึกษาเชิงวิเคราะห์ย้อนหลัง

สถานที่ทำการวิจัย : ภาควิชาพยาธิวิทยา คณะแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย

ตัวอย่างและวิธีการศึกษา : อาศัยชิ้นเนื้อจากผู้ป่วยโรคมะเร็งลำไส้ใหญ่จำนวน 162 ราย โดยแบ่งเกรดหรือระดับการเปลี่ยนแปลงของเนื้องอกเป็นเจริญเปลี่ยนแปลงได้ดี ปานกลาง แย่ และแทบจะไม่เจริญเปลี่ยนแปลงเลย ซึ่งอ้างอิงตามเงื่อนไขการแบ่งขององค์การอนามัยโลกเมื่อปี พ.ศ.2543 สำหรับดัชนีบ่งชี้การเจริญของเซลล์มะเร็ง ศึกษาโดยอาศัยการย้อมชิ้นเนื้อทางกระบวนการอิมมูโนฮิสโตเคมี แล้วคำนวณความสัมพันธ์ระหว่างระดับการเปลี่ยนแปลงของเนื้อมะเร็งกับอัตราการเจริญของเซลล์มะเร็งลำไส้ใหญ่ด้วยกระบวนการทางสถิติ

ผลการศึกษา : พบความสัมพันธ์ระหว่างระดับการเปลี่ยนแปลงของเนื้อมะเร็งกับการเจริญของเซลล์อย่างมีนัยสำคัญทางสถิติ ($P = 0.014$)

สรุป : ถ้าอัตราการแบ่งตัวของมะเร็งยิ่งสูงขึ้น การเปลี่ยนแปลงของเซลล์มะเร็งลำไส้ใหญ่ก็ยิ่งแย่ลงตามไปด้วย ดังนั้นอาจนำอัตราการแบ่งตัวของมะเร็งมาใช้ทดแทนได้ในมะเร็งบางรายที่ยากต่อการให้เกรด

คำสำคัญ : การแบ่งตัวของเซลล์, มะเร็งลำไส้ใหญ่, การเปลี่ยนแปลงของเนื้องอก.

The distribution of colorectal cancer is worldwide; between 20 % and 50 % of patients with colorectal cancer will pass away within five years after their diagnosis, generally because of extensive metastasis.⁽¹⁾ During the past decade this malignancy has been studied in various facets, e.g., epidemiology, etiology or pathogenesis.⁽²⁾ One of the most significant influent factors for predicting the patient outcome is biological markers of neoplastic behavior.^(3, 4) Apart from conventional stage and grading system, according to tumor differentiation, the proliferation factors at biomolecular levels have an important role in patient management and biological classification of tumors.⁽⁴⁾

Neoplastic cellular proliferation will act autonomically without control. Ki67 is a cell cycle associated antigen of nuclear protein which expresses in all cell growth phases except the resting phase (G_0).^(5,6) The number of cells stained positively for monoclonal Ki67 antibody correlates with tissue proliferation rate and degree of tumor differentiation.⁽⁷⁾ Our study is designed to prove this concept and to evaluate the utility of Ki67 as a prognostic biomarker of colorectal adenocarcinoma.

Materials and Methods

One or two paraffin blocks per case from 162 cases, first diagnosed as colorectal adenocarcinoma at the King Chulalongkorn Memorial Hospital from 2002 to 2003 were selected. Representative blocks always comprised the deep invasive portion and in the majority of cases superficial parts of the tumor were also included. Subjects who have had a history of polyposis syndrome, recurrent tumor and previous treatment, either chemotherapy or radiation, were

excluded from the study. Four- micrometer-thick haematoxylin and eosin stained slides were reviewed and classified the tumor differentiation, according to the World Health Organization (WHO) histological classification of tumors of the colon and rectum. They consisted of well-differentiated (grade 1), moderately -differentiated (grade 2), poorly-differentiated (grade 3) and undifferentiated (grade 4) when glandular structures occurred >95 %, 50 – 95 %, 5 – 50 % and <5 %, respectively. In addition some variants, composed of mucinous adenocarcinoma and signet-ring cell carcinoma, were classified as poorly differentiated whereas medullary carcinoma was classified as undifferentiated. Those specimens were noted the depth of tumor penetration, number of regional lymph node metastasis and distant organ spreading, according to TNM staging as well as their location, including ascending, transverse, descending and rectosigmoid regions.

In the determination of neoplastic cell proliferation, formalin-fixed, paraffin wax-embedded blocks were deparaffinized in xylene, rehydrated through graded alcohols and employed Ki67 antibody (mouse antihuman, clone MIB1, 1/300 dilution; Dako), a biotin-conjugated secondary antibody, avidin-horseradish peroxidase, and the chromogen diaminobenzidine tetrahydrochloride (DAB) as a detection agent on the tumor tissue sections. Ki67 immunohistochemical stains were evaluated semiquantitatively and counted the number of positive cells expressing nuclear brown stain among the neoplastic cells. Field selection used highest Ki67 expression areas (hot spots) by lower power scanning and scored as: negative, <10 %; 1+, 10 - 25 %; 2+, 26 – 50 %; 3+, 51-75 %; and 4+, >75 %.

Discussion

Colorectal carcinoma still remains a leading cause of cancer death in most countries without evidence of any decline in its incidence.^(1,2) Regardless of surgical resection, which is the principle management, most oncologists need effective marker for patient follow-up and predicting the outcome. Recently, there is not only one ideal marker, but it is used as a panel and its reliability remains varied among different geographic areas.^(3,4,8)

Similar to other malignancies, tumor grade or degree of differentiation is depended on tumor molecular kinetic and considered to be an important independent prognostic factor besides the staging system.⁽³⁾ Differentiation of colorectal adenocarcinomas are usually graded into, namely: well- differentiated (grade 1), moderately- differentiated (grade 2) and poorly-differentiated (grade 3), when glandular architectures appear >95 %, 50 - 95 % and 5 - 50 %, respectively, and the colonic tumor is classified as undifferentiated type (grade 4) when glandular formation occurs <5 %, according to the WHO classification.⁽⁹⁾

However, histologic grading is subjective. Some authors have recommended a five-grade system, including well, well to moderately, moderately, moderately to poorly and poorly differentiated in order to complete the spectrum.⁽¹⁰⁾ On the other hand, some authors have suggested that colonic tumor should be divided simply into low-grade, consisting of well and moderately-differentiated adenocarcinomas, and high-grade, including poorly-differentiated and undifferentiated carcinomas. The reason for this classification is that in multivariate analyses there is no difference between well and moderately differentiated tumors.⁽⁹⁾

Therefore, grading is problematic in some situations.⁽¹⁰⁾ When a carcinoma possesses variegated differentiation, grading should be based on the least differentiated element and avoided the advancing edge.^(10,11) Small foci of less differentiated are common at the deepest part, but this feature is insufficient to classify the tumor as poorly differentiated.⁽¹²⁾ However, another way is to assess the growing tumor edge separately from the overall grade because this tumor fraction is considered to be most aggressive and more related tumor behavior.⁽¹³⁾ In addition, in case of tiny biopsy specimen, gradation is difficult to report and tumor grades may not represent the whole tumor.

Ki67 is the antibody for detecting the intranuclear matrix protein, correlating with cell proliferation. In tumorigenesis turnover rate of malignant cells is significantly higher than normal because reversible G_1-G_0 pathway is mostly skipped.⁽⁵⁻⁷⁾ Then Ki67 expression in malignancy is higher than in normal.⁽¹⁴⁻¹⁶⁾ Backus HH and colleagues have found that the progression of primary colonic adenocarcinoma to liver metastasis is associated with an increased proliferation rate, using Ki67 measurement.⁽¹⁷⁾

In the present study, we found that the higher Ki67 is, the worse is the neoplastic differentiation. This result is statistically significant. Pathologists may add Ki67 as a prognostic biomarker, particularly in problematic cases such as heterogeneous differentiation of the tumor. For adenocarcinoma of the colon, there is a literature, that has prove it; according to which the result of Ki67 expression is similar between the central and peripheral parts of the tumor.⁽¹⁴⁾

In conclusion, Ki67 immunostain for determining cell proliferation is related to tumor differentiation or tumor grading in colorectal adenocarcinoma. This marker is sensitive and can probably be used for monitoring and predicting the stage of the disease and therefore helps planning the therapy.

References

1. Prall F, Gringmuth U, Nizze H, Barten M. Microvessel densities and microvascular architecture in colorectal carcinomas and their liver metastases: significant correlation of high microvessel densities with better survival. *Histopathology* 2003 May;42(5):482-91
2. Vassilopoulos PP, Kelessis N, Plataniotis G, Gondikakis E, Galanos A. Colorectal cancer trends by anatomic sides, age and staging. A twenty-year study of 1412 Greek cases. *Anticancer Res* 2000 Nov-Dec;20(6C):4773-6
3. Nguyen VN, Mirejovsky P, Mirejovsky T, Melinova L, Mandys V. Expression of cyclin D1, Ki-67 and PCNA in non-small cell lung cancer: prognostic significance and comparison with p53 and bcl-2. *Acta Histochem* 2000 Aug; 102(3):323-38
4. Chapusot C, Assem M, Martin L, Chalabreysse L, Benhamiche AM, Lignier MA, Chauffert B, Teyssier JR, Faivre J, Piard F. Expression of p21 WAF1/CIP1 protein in colorectal cancers: study of its relation to p53 mutation and Ki67 antigen expression. *Pathol Biol (Paris)* 2001 Mar;49(2):115-23
5. Hui EP, Poon TC, Teo PM, Mo F, Zee B, Leung SF, Ho S, Mok TS, Kwan WH, Johnson PJ, et al. A prospective study of pre-treatment cell kinetics and clinical outcome in nasopharyngeal carcinoma. *Radiother Oncol* 2003 Oct;69(1):53-62
6. Al-Moundhri MS, Nirmala V, Al-Hadabi I, Al-Mawaly K, Bumey I, Al-Nabhani M, Thomas V, Ganguly SS, Grant C. The prognostic significance of p53, p27 kip1, p21 waf1, HER2/neu, and Ki67 proteins expression in gastric cancer: a clinicopathological and immunohistochemical study of 121 Arab patients. *J Surg Oncol* 2005 Sep 15;91(4):243-52
7. Kanavaros P, Stefanaki K, Valassiadou K, Vlachonikolis J, Mavromanolakis M, Vlychou M, Kakolyris S, Gorgoulis V, Tzardi M, Georgoulis V. Expression of p53, p21/waf, bcl-2, bax, Rb and Ki67 proteins in colorectal adenocarcinomas. *Med Oncol* 1999 Apr;16(1): 23-30
8. Alexander J, Watanabe T, Wu TT, Rashid A, Li S, Hamilton SR. Histopathological identification of colon cancer with microsatellite instability. *Am J Pathol* 2001 Feb;158(2):527-35
9. Hamilton SR, Vogelstein B, Kudo S, Riboli E, Nakamura S, Hainaut P, Rubio CA, Sobin LH, Fogt F, Winawer SJ, et al. Carcinoma of the colon and rectum. In: Hamilton SR, Aatonen LA, eds. *Pathology & Genetics of Tumours of the Digestive System*. World Health Organization Classification of Tumours. Lyon: International Agency for Research on Cancer (IARC) Press, 2000: 103-19
10. Riddell RH, Petras RE, Willis GT, Sobin LH. Colorectal carcinoma. In: Rosai J, Sobin LH, eds. *Atlas of Tumor Pathology. Tumors of the*

- Intestine. 3rd series. Fascicle 32. Washington, DC: Armed Forces Institute of Pathology, 2003: 145-52
11. Ikeda K, Pant B, Mishiro A, Ozawa K, Masujima T, Sugiyama M. A convenient method for the evaluation of anti-tumor agents affecting the cell cycle. *J Biosci Bioeng* 2000;90(5): 574-6
 12. Greenson JK, Bonner JD, Ben-Yzhak O, Cohen HI, Miselevich I, Resnick MB, Trougouboff P, Tomsho LD, Kim E, Low M, et al. Phenotype of microsatellite unstable colorectal carcinomas: Well-differentiated and focally mucinous tumors and the absence of dirty necrosis correlate with microsatellite instability. *Am J Pathol* 2003 May;27(5): 563-70
 13. Ueno H, Murphy J, Jass JR, Mochizuki H, Talbot IC. Tumor 'budding' as an index to estimate the potential of aggressiveness in rectal cancer. *Histopathology* 2002 Feb;40(2): 127-32
 14. Berenzi A, Benetti A, Bertalot G, Rodolfi A, Portolani N, Giulini SM, Pulcini G, Vinco A, Tiberio G. Ki67 immunohistochemical evaluation in colorectal cancer and normal colonic mucosa. Possible clinical applications. *Pathologica* 1992 Mar-Apr;84 (1090):155-63
 15. Jaskiewicz K, Lancaster E, Banach L, Karmolinski A. Proliferative activity of normal and neoplastic colonic mucosa in population groups with high and low risk for colorectal carcinoma. *Anticancer Res* 1998 Nov-Dec; 18(6B):4641-4
 16. Risio M, Rossini FP. Cell proliferation in colorectal adenomas containing invasive carcinoma. *Anticancer Res* 1993 Jan-Feb;13(1):43-7
 17. Backus HH, Van Groeningen CJ, Vos W, Dukers DF, Bloemena E, Wouters D, Pinedo HM, Peters GJ. Differential expression of cell cycle and apoptosis related proteins in colorectal mucosa, primary colon tumors, and liver metastases. *J Clin Pathol* 2002 Mar; 55(3):206-11