

11-1-2003

CK-MB isoenzyme exceeding creatine kinase, a type of macro creatine kinase phenomenon in Thai hospitalized patients, retrospective data from King Chulalongkorn Memorial Hospital

V. Wiwanitkit

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Wiwanitkit, V. (2003) "CK-MB isoenzyme exceeding creatine kinase, a type of macro creatine kinase phenomenon in Thai hospitalized patients, retrospective data from King Chulalongkorn Memorial Hospital," *Chulalongkorn Medical Journal*: Vol. 47: Iss. 11, Article 4.

Available at: <https://digital.car.chula.ac.th/clmjjournal/vol47/iss11/4>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

CK-MB isoenzyme exceeding creatine kinase, a type of macro creatine kinase phenomenon in Thai hospitalized patients, retrospective data from King Chulalongkorn Memorial Hospital*

Viroj Wiwanitkit **

Wiwanitkit V. CK-MB isoenzyme exceeding creatine kinase, a type of macro creatine kinase phenomenon in Thai hospitalized patients, retrospective data from King Chulalongkorn Memorial Hospital. Chula Med J 2003 Nov; 47(11): 727 - 34

Objective : *To determine the prevalence, etiology and clinical significance of macro creatine kinase (CK) defined as CK-MB isoenzyme exceeding creatine kinase, in hospitalized Thai patients.*

Methods : *All data from laboratory requests for total CK and CK-MB in the inpatient unit sent to Clinical Chemistry Unit, Chulalongkorn University Hospital during the year 1999 were studied. Cases were CK-MB exceeded total CK, defined as a type of macro CK phenomenon, were included. Review of the medical records of these inpatients was performed.*

Results : *From this study, 5 cases with macro CK were detected. There were 4 female and 1 male patients with age ranging from 34 to 70 years old (average age = 54.8 ± 13.5 years old). Diseases of many systems including brain, lung and liver diseases were found to be the causes of macro CK. No deaths were found in this study.*

* Presented at 41st Annual Scientific Meeting, Faculty of Faculty of Medicine, Chulalongkorn University, 2000.

** Department of Laboratory Medicine, Faculty of Medicine, Chulalongkorn University

Conclusion : *Like other previous studies, the prevalence of macro creatine kinase is about 1.5 % and usually found in older female patients.*

Keyword : *Macro creatine kinase*

Reprint request : Wivanitkit V. Department of Laboratory Medicine, Faculty of Medicine,
Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. May 15, 2002.

วิโรจน์ ไวรานิขกิจ. CK-MB isoenzyme ระดับสูงกว่า creatine kinase, ปะการุกการณ์ macro creatine kinase ชนิดหนึ่งในผู้ป่วยชาวไทย การศึกษาย้อนหลัง ณ โรงพยาบาลจุฬาลงกรณ์. จุฬาลงกรณ์เวชสาร 2546 พ.ย; 47(11): 727 - 34

- วัตถุประสงค์** : เพื่อศึกษาความชุกของปะการุกการณ์ macro creatine kinase (CK) ในรูปแบบที่ CK-MB isoenzyme มีระดับสูงกว่า creatine kinase ในผู้ป่วยในชาวไทย
- วิธีการ** : ทำการศึกษาย้อนหลังจากบันทึกการส่งตรวจ total CK และ CK-MB ในกลุ่มผู้ป่วยใน ณ หน่วยเคมีคลินิกโรงพยาบาลจุฬาลงกรณ์ ในช่วงปีพ.ศ. 2542 ในกรณีที่ CK-MB มีระดับสูงกว่า total CK ซึ่งจัดได้ว่าเป็นปะการุกการณ์ macro CK ชนิดหนึ่งได้ทำการคัดเลือกไว้เพื่อทำการศึกษาประวัติต่อไป
- ผลการศึกษา** : ได้จำนวนกรณีศึกษาปะการุกการณ์ macro CK จำนวน 5 กรณี โดยเป็นผู้ป่วยหญิง 4 ราย ผู้ป่วยชาย 1 ราย อายุระหว่าง 34 และ 70 ปี (อายุเฉลี่ย = 54.8 ± 13.5 ปี) พบว่าโรคทางสมอง, ปอด และตับเป็นสาเหตุที่เกี่ยวข้องกับการเกิดปะการุกการณ์ macro CK ทั้งนี้ไม่พบผู้ป่วยที่ตายในโรงพยาบาลในกลุ่มผู้ป่วยที่ทำการศึกษานี้
- สรุป** : ความชุกของปะการุกการณ์ macro creatine kinase (CK) ในการศึกษาครั้งนี้ได้เท่ากับ 1.5 % ใกล้เคียงกับการศึกษาที่ผ่านมา และมักพบปะการุกการณ์นี้ในผู้ป่วยหญิงชรา
- คำสำคัญ** : Macro creatine kinase

Creatine kinase (CK) is an important enzyme in medicine. Determination of levels of total CK and its isoenzymes is a widely used laboratory test in the present day. CK-MB is a frequently requested CK isoenzyme due to its usefulness as a marker of myocardial infarction.

The total CK level and CK-MB isoenzyme level is determined in the laboratory. Sometimes spurious results due to total and partial determination of unknown values can be found.⁽¹⁾ The macro creatine kinase is an interesting cause for aberrant determination for total CK and CK-MB levels. A higher CK-MB level than total CK level result makes the interpretation of results difficult. This extremely excessive CK-MB is known as one form of macro creatine kinase.

This study was performed in order to determine the prevalence, etiology and clinical significance of such cases in hospitalized Thai patients.

Materials and methods

All data on 323 laboratory requests for total CK and CK-MB in the inpatient unit sent to Clinical Chemistry Unit, Chulalongkorn University Hospital during the year 1999 were studied. Both total CK and CK-MB levels in this study were determined by the NAC - activated method (Boehringer Mannheim). The NAC- activated method is an enzymatic based chemical determination. The unknown CK or CK-MB will catalyze the reaction between creatine phosphate and ADP in the reagent to form ATP. The resulting ATP then reacts with glucose in the reagent to provide glucose-6-P. In the last step of the cascade, glucose-6-P will react with NADP in the reagent to produce NADPH. Then NADPH is measured by UV photometry.

Cases where CK-MB exceeded total CK, defined as a type of macro CK phenomenon, were included. Review of medical records of these inpatients was performed.

Table 1. Detail of case with higher CK-MB level than total CK level in this series.

No	Sex	Age (years)	Total CK(U/l)	CK-MB (U/l)	Definite diagnosis	EKG abnormality
1	Female	68	21	22	Temporal lobe epilepsy, cryptococcal meningitis	No
2	Female	70	50	58	Unstable angina, respiratory failure	No
3	Female	56	103	137	Acute pyelonephritis, Cirrhosis,	No
4	Male	34	26	61	Sepsis	No
5	Female	46	38	45	Pulmonary congestion due to congestive heart failure Lobar pneumonia	No

Results

From this study, 5 cases with macro CK were detected. There were 4 female and 1 male patients with ages ranging from 34 to 70 years (average age = 54.8 ± 13.5 years old). The prevalence of the macro creatine kinase was 1.5 % (total requests equaled to 323 tests).

From medical record review, all 5 cases were suspected for acute coronary syndrome, therefore, the blood samples were collected and sent to the laboratory for analysis of total CK and CK-MB. However, the electrocardiogram examinations of all cases showed no evidence of acute coronary syndrome. Hence, they were not diagnosed to have acute coronary syndrome and not performed follow up total CK and CK-MB tests. Baseline details of each case in this series were presented in Table 1. No death case was detected in this study.

Discussion

These cases are examples of problematic cases in total and partial determination of chemical substances in body fluids, resulting in abnormal laboratory results. The sum quantity of components exceeded the total quantity. Interpretation of these abnormal laboratory results was very difficult.

All cases were those in which the CK-MB level exceeded the CK level. Usually an isoenzyme level should not exceed the total enzyme level.⁽²⁾ There must be some error if the isoenzyme level exceeds total enzyme level. What is the reason? Both total CK and CK-MB levels were measured by the same NAC-activated method. But in the total CK determination all forms of CK were measured. Therefore, the quantity of total CK by this method is

implied directly by determination of product of chemical reaction. While CK-MB determination is based on the principle of immunoassay. A specific antibody is used to inhibit the CK-M moiety without affecting the CK-B moiety. In this method, conclusion that CK-B fraction accounts for one half the activity of CK-MB is accepted. The CK-B fraction is determined by the NAC-activated method. The CK-MB level is assumed to be equal to two times the CK-B level. Therefore, the presence of isoenzyme CK-BB may stimulate an increased proportion of CK-MB in the immunologic CK-MB assay. CK-BB can stimulate falsely elevated CK-MB exceeding 35 % and even up to twice the total CK activity. CK-BB can be measured in the serum after damage to many organs such as brain, liver, kidney and lung.⁽³⁻⁴⁾

An atypical CK isoenzyme result, called macro CK,⁽⁵⁾ can be found in 3 - 6 % of hospitalized patient especially in elderly females. In this situation, the total CK is usually in the normal range but a high CK-MB level can be observed. In these patients, the reasons relating to macro creatine kinase are good examples for why CK-MB level exceeds total CK level. The limitation of biochemistry tests for determination of isoenzyme has been discussed. Another alternative method to measure isoenzymes, such as electrophoresis, should be used in these problematic cases.

Macro creatine kinase (Macro CK) is a major cause of elevations of the creatine kinase MB isoenzyme (CK), which may cause mistakes in the evaluation of patients suspected of ischemic cardiopathy.⁽⁶⁻⁸⁾ Its presence in serum interferes with the immunoinhibition methods normally used in emergency room laboratories causing false positive

results in the evaluation of the total creatine kinase level (CK), compared to the CK-MB levels. Usually, the CK-MB activity exceeds total CK activity by more than 25 %.^(6,8 - 10) Increasing concentrations of various types of creatine kinase isoenzymes mimic myocardial infarction or injury.⁽¹¹⁾ Probably a maximum of 1% of patients with suspected acute myocardial infarction,^(11 - 13) have macro CK in concentrations causing diagnostic errors.

Of interest, in Thailand, there had been no reported about the prevalence of macro CK and its clinical correlation among the Thais, therefore, we performed this study. Like the previous studies,^(11 - 13) our study showed that the macro CK was common in elderly female patients, with an approximate prevalence of 1 %. In clinical practice, markedly elevated levels of CK-MB, or increased levels of CK-MB in combination with CK-BB may point away from a myocardial origin and toward the existence of a malignancy. Macro CK can be detected as an abnormal band in creatine kinase electrophoresis. However, in the setting with lack of electrophoresis, the misinterpretation can be performed without the knowledge on macro CK.

In general, macro CK can be distinguished as two types, type I and type II.⁽¹⁴⁾ Presence of both variant CK isoenzymes may lead to diagnostic or therapeutic errors due to an altered CK:CK-MB ratio.⁽¹⁵⁾ The serum total CK is significantly higher, and an increased CK-MB proportion is also significantly more common in patients with macro CK type 1 than in those with type 2. Macro CK types 1 and 2 are much more heat stable than CK-MB and CK-BB, and thus, by heating samples for 20 min at 45 degrees C, the presence of thermostable macro types can be

demonstrated.⁽¹²⁾ In addition, macro CK type 2 has a much higher activation energy than macro CK type 1. The phenomenon described as CK-MB exceeded total CK like in this study is the common presentation of macro CK type II.⁽¹⁶⁾ Although this type of macro CK is not as clinically important as macro CK type I, it is always misinterpreted as laboratory analysis error and resulted in repeated request for retest and also result in unnecessary lost of money.

Nevertheless, like other macro enzymes⁽¹⁷⁾ such as macroamylase and macro lactate dehydrogenase, macro CK type II can be detected in advanced age and malignant lesions.⁽¹⁸⁾ It can be found in diseases of many organ systems such as brain,⁽¹⁹⁾ thyroid,⁽²⁰⁾ GI,^(21 - 22) prostate,⁽²²⁾ breast⁽²²⁾ and kidney.⁽²³⁾ However, macro CK isoenzyme type II in the serum of apparently healthy individuals has also been reported.^(24 - 27) In conclusion, like other previous studies, the prevalence of macro creatine kinase is about 1.5 % and usually found in the old female patients.

Acknowledgement

The author would like to thank all medical personnel who perform the CK test. We are also thankful to Associate Professor Narudee Bhokaisawan who give the first idea for us on this topic.

References

1. Wiwanitkit V. Total and partial determination in biochemistry laboratory tests: problematic cases. *Chula Med J* 1999 Feb; 43(2): 109 - 13
2. Binder L. The CK-MB isoenzyme is higher than creatine kinase. *Internist* 1996 Apr; 37(4): 398-9

3. Vrbica Z, Durovic O, Oreb N. Interference of CK-BB isoenzyme in the determination of CK-MB using the immunoinhibition method in patients with pulmonary diseases. *Lijec Vjesn* 1997 Nov; 119(10): 263 - 5
4. Shoji S. Creatine kinase. *Nippon Rinsho* 1995 May; 53(5): 1136 - 40
5. Urdal P, Landaas S. Macro creatine kinase BB in serum and some data on its prevalence. *Clin Chem* 1979 Mar; 25(3): 461 - 5
6. Rosa-Jimenez F, Gasso de Campos M, Camacho Reina MV, Vazquez de Castroviejo E, Hernandez Burruezo JJ, Pousibet Sanfeliu H. Macro creatine kinase type 1 as a cause of increase of CF-MB isoenzyme. Apropos of 7 cases. *Rev Esp Cardiol* 1997 Mar; 50(3): 166 - 72
7. Hulting J, Waldenlind L, Onica D, Wallinder H. Creatine kinase -MB mass concentration versus creatine kinase-B activity for the detection of acute myocardial infarction in patients with slightly elevated total creatine kinase activity in serum. *J Intern Med* 1994 Mar; 235(3): 211 - 6
8. Punt JM. Macro-creatine-kinase: not every increased CK-MB activity points to a heart infarct. *Ned Tijdschr Geneesk* 1989 Oct 28;133(43): 2142 - 3
9. Wang HC, LU JY, Ting YM. Elevated serum CK-MB and CK-BB isoenzymes in a patient with small cell lung cancer. *Chung Hua I Hsueh Tsa Chih (Taipei)* 1995 Mar; 55(3): 270 - 3
10. Raga X, Vilanova MA, Bardaji A, Pastor MR, Richart C. Macrocreatine kinase, a cause of MB isoenzyme levels falsely elevated. *Arch Mal Coeur Vaiss* 1992 Nov; 85(11): 1601 - 3
11. Antonsen S, Botker HE, Horder M. Macro creatine kinase. A cause of diagnostic problems in acute myocardial infarction. *Ned Tijdschr Geneesk* 1989 Jun 12; 151(24): 1526 - 8
12. Tsung JS. Creatine kinase and its isoenzymes in neoplastic disease. *Crit Rev Clin Lab Sci* 1986; 23(1): 65 - 75
13. Devine JE. Macro-creatine kinase as an interference in CK isoenzyme determinations. *Enzyme* 1983; 30(2): 139 - 44
14. Lee KN, Csako G, Bernhardt P, Elin RJ. Relevance of macro creatine kinase type 1 and type 2 isoenzymes to laboratory and clinical data. *Clin Chem* 1994 Jul; 40(7 pt 1): 1278 - 83
15. Bayer PM, Wider G, Unger W, Kohn H, Hajdusich P, Hotschek H. Variant creatine kinase isoenzymes: frequency and clinical significance. *Klin Wochenschr* 1982 Apr 1; 60(7): 365 - 9
16. Delanghe J, De Scheerder I, De Buyzere M, Algoed L, Robbrecht J. Macro CK type 1 as a marker for autoimmunity in coronary heart disease. *Atherosclerosis* 1986 Jun; 60(3): 215 - 9
17. Kuint J, Pipano S, Linder N, Reichman B, Sela BA. Serum macro creatine kinase type 2 in asphyxiated newborn infants. *Clin Biochem* 1993 Apr; 26(2): 117 - 20
18. de Almeida Camarozano AC, Gaspar Henriques LM. A macromolecule capable of modifying CK-MB results and of inducing false acute myocardial infarction diagnosis. *Arq Bras Cardiol* 1996 Mar; 66(3): 143 - 7
19. Shimoda K, Hirata Y, Terai Y, Usui K, Morofuji M, Umei T, Sumida I, Hanada M. A case of

- hepatic encephalopathy with high blood level of macro creatine kinase. *Nippon Naika Gakkai Zasshi* 1990 May 10; 79(5): 680 - 1
20. Giampeitro O, Clerico A, Buzzigoli G, Lucchetti L, Boni C, Del Chicca MG, Mariani G. Macro serum CK-BB in a woman with severe primary hypothyroidism. *Am J Med Sci* 1985 Apr; 289(4): 160 - 3
21. Hirata RD, Hirata MH, Strufaldi B, Possik RA, Asai M. Creatine kinase and lactate dehydrogenase isoenzymes in serum and tissues of patients with stomach adenocarcinoma. *Clin Chem* 1989 Jul; 35(7): 1385 - 9
22. Wong SS, Wu AH, Fritsche HA. Reassessment of creatine kinase BB as a marker for cancer of the prostate, breast, and lung. *Clin Chem* 1987 Jun; 33(6): 809 - 11
23. Nakashima A, Toyama T, Okuda K, Kuroda H. Two cases of chronic renal failure on maintenance hemodialysis, presenting macro creatine kinase. *Nippon Jinzo Gakkai Shi* 1986 Jun; 28(6): 829 - 35
24. Maire I, Artur Y, Sanderink GJ. Macroenzymes in human plasma. 1: Macroamylase, macrocreatin kinase, macrolactate dehydrogenase. *Ann Biol Clin (Paris)* 1987; 45(3): 269 - 76
25. Tsung SH. Total CK activity and isoenzyme patterns in normal and neoplastic tissue of gastrointestinal tract. *J Clin Pathol* 1982 Feb; 35(2): 204 - 6
26. Whelan PV, Malkus H. A macro creatine kinase isoenzyme in serum of apparently healthy individuals. *Clin Chem* 1983 Jul; 29(7): 411 - 4
27. Kanemitsu F, Okigaki T. Clinical application of subforms of creatine kinase MM and macro creatine kinases. *J Chromatogr* 1990 Apr 6; 526(2): 423 - 38