

1-1-2019

Pediatric respiratory severity score evaluates disease severity of respiratory tract infection in children

Janjira Thokngaen

Wissaroot Karoonboonyanan

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjournal>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Thokngaen, Janjira and Karoonboonyanan, Wissaroot (2019) "Pediatric respiratory severity score evaluates disease severity of respiratory tract infection in children," *Chulalongkorn Medical Journal*: Vol. 63: Iss. 1, Article 7.

DOI: 10.58837/CHULA.CMJ.63.1.6

Available at: <https://digital.car.chula.ac.th/clmjournal/vol63/iss1/7>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact ChulaDC@car.chula.ac.th.

Original article

Pediatric respiratory severity score evaluates disease severity of respiratory tract infection in children

Janjira Thokngaen, Wissaroot Karoonboonyanan*

Department of Pediatrics, Bhumibol Adulyadej Hospital, Bangkok, Thailand

Background: Acute respiratory infections (ARIs) are commonly found causes of morbidity and mortality in children aged below 5 years old. The pediatric respiratory severity (PRESS) score is a simple severity scoring system. Healthcare providers can apply it as a preliminary patient assessment for proper treatments.

Objective: To evaluate the pediatric respiratory severity score as a severity assessment for pediatric patients with acute respiratory infections.

Methods: This is a prospective study from 1st September 2016 to 31st October 2017. The study group includes 120 patients aged 3 months to 14 years old. The use of the PRESS score as an assessment tool has 5 parameters: respiratory rate, wheezing, accessory muscle use, peripheral oxygen saturation (SpO₂), and feeding difficulties, and can be classified into 3 groups: mild (score of 0 or 1), moderate (score of 2 or 3), and severe (score of 4 or 5). The primary outcomes are sensitivity, and specificity for hospitalization. The secondary outcomes are sensitivity, specificity of intensive care unit (ICU) admission, mean and standard deviation of the duration of oxygen therapy and nebulized bronchodilator.

Results: The admitted regular patients in the moderate and severe groups had a sensitivity of 0.94 and a specificity of 0.88, whilst the ICU patients in the severe group had a sensitivity of 0.75 and a specificity of 0.66, longer duration for oxygen treatment, and longer duration for nebulized bronchodilator treatment. Furthermore, this system is more reliable than previous respiratory severity score.

Conclusions: PRESS scores could predict condition severity and might guide a proper treatment of acute respiratory infection in children.

Keywords: Pediatric severity score, respiratory tract infection, PRESS score.

Acute respiratory infections (ARIs) in children are one of the most common cause of morbidity and mortality in children aged younger than 5 years old.^(1, 2) Lower respiratory tract infections (LRI) are the leading causes of death in children. The death toll reached 10.8 million casualties per year. There were 1.9 million deaths from ARIs and the World Health Organization (WHO) found approximately 2 million children under five died yearly from pneumonia in 2005.⁽³⁾ In Thailand, the Pediatric Respiratory and Critical Care Medicine Society of Thailand (PRCS) published data which indicate that in 2013 the prevalence of pneumonia was about

45-50 percent of children under 5 years of age with acute respiratory tract infections.⁽⁴⁾ Data from a weekly epidemiological surveillance report shows pneumonia in Thailand, published in 2012, during 2003 – 2010, the highest occurrence of pneumonia was in the age group between zero to four years old. In 2010, the age group 0 - 4 years old had a rate of 1,939.49 per 100,000 population.⁽⁵⁾ From the pediatric patients record of Bhumibol Adulyadej Hospital, the Royal Thai Air Force, the number of admitted pediatric patients with LRI in the past 5 years reached about 200 patients per year.

At multiple community hospitals, it is found that in addition to a doctor, there are operational nurses and trained public health center officers who can provide basic medical treatment. Additionally, it is also found that some patients living in remote areas, and consequently having difficulties visiting any provincial hospital, prefer to receive treatments at

*Correspondence to: Wissaroot Karoonboonyanan, Department of Pediatrics, Bhumibol Adulyadej Hospital, Bangkok 10220, Thailand.

Received : February 9, 2018

Revised : March 19, 2018

Accepted : May 9, 2018

community hospitals or public health centers nearest to their residences.

In community hospitals, there are many pediatric patients with fevers, coughs, and dyspnea, hence, several cases are found, e.g., some patients have a respiratory failure that requires immediate treatment and even urgent transfer. Some are with moderate symptoms and not with severe dyspnea, and they require oxygen treatment, nebulization, and need an admission. Some with less severe symptoms can be treated with a nebulization and go home, etc. Therefore, these characteristics change their assessment. The assessments practically depend on the individual's experience, which is not standardized nor equal. If there is a simple assessment scoring system, that everyone can utilize, it can be of great use to properly evaluate the severity of the case.⁽⁶⁻⁹⁾ The patients could be better treated promptly and more appropriately. As a result, it may reduce complications and ultimately mortality rates.

The Department of Pediatrics, National Hospital of Yokohama, published a research article: Pediatric Respiratory Severity Score (PRESS) for Respiratory Tract Infections in Children in July 10, 2015, showing a new and simpler bedside scoring system.⁽¹⁰⁾ It is for the assessment of severity of pediatric patients with respiratory infections classified into three groups:

mild (0 - 1), moderate (2 - 3), severe (4 - 5) based on five parameters: respiratory rate, wheezing, accessory muscle use, SpO₂, and feeding difficulties (Table 1).

Based on the review, the PRESS score is easy and simple for assessing a patient. Each parameter has either a score of 0 or 1, where 1 is present, and 0 is otherwise. A summation of these parameters classifies the group of the patient as mild (with score 0 to 1), moderate (with score 2 to 3), and severe (with score 4 to 5). When the classification identifies condition as severity, this consequently indicates a proper treatment with urgency. Thus, patients can be treated promptly and appropriately.

Methods

The objective of this study is to evaluate the PRESS score as a severity assessment for pediatric patients with acute respiratory infections.

The research design follows the prospective study scheme. The duration of this research was from 1st September 2016 to 31st October 2017. The location of the data is from the pediatric outpatient and inpatient clinic, Bhumibol Adulyadej Hospital, the Royal Thai Air Force. The target population for the research is pediatric patients aged between 3 months to 14 years diagnosed with respiratory infections, that includes both outpatients and inpatients.

Table 1. PRESS scoring system.

Score component	Operational definition		Scoring
Respiratory rate	Respiratory rate at rest, on room air*		0 or 1
Wheezing	High-pitch expiratory sound heard by auscultation		0 or 1
Accessory muscle use	Any visible use of accessory muscles		0 or 1
SpO ₂	Oxygen saturation <95% on room air		0 or 1
Feeding difficulties	Refusing feedings		0 or 1
	Sum of five components		
PRESS score	0 - 1: mild, 2 - 3: moderate, 4 - 5: severe		0 - 5
Criteria of tachypnea*	Month	Respiratory rate	
	<12	>60	1
	≤12, <36	>40	1
	≤36, <156	>30	1
	≥156	>20	1

*Respiratory rate evaluated according to American Heart Association guideline.

PRESS: Pediatric Respiratory Severity score, SpO₂: Peripheral oxygen saturation.

Statistical analysis of the results from data collected from 120 cases, with primary outcomes as sensitivity and specificity of hospitalization, and secondary outcomes as sensitivity and specificity of ICU (Intensive Care Unit) admission, was performed using the receiver operating characteristics (ROC) curve to analyse the area under the curve (AUC) of PRESS scores, mean \pm SD of the duration of oxygen therapy and nebulized bronchodilator. The categorical data were assessed using Chi-square tests. The continuous data were expressed as a mean \pm standard deviation (SD) with 95% confidence intervals (CI) where applicable, and the significant level was set as 0.05 on SPSS (version 24).

Results

In this research, the population size was 120 patients, 61 females (50.8%), and 59 males (49.2%) (Table 2). The average age was 37.2 ± 35.9 months old (mean \pm SD). There were 33 upper ARIs patients (27.5%), and another 87 patients (72.5%), diagnosed as lower ARIs. There were 86 admitted patients (71.6%), who consisted of 82 general pediatric admitted patients (68.3%), and 4 ICU admitted patients (3.3%). The rest were 34 non-admitted patients (28.3%). The results show that there were 35 mild patients (29.2%), 43 moderate patients (35.8%), and 42 severe patients classified in accordance to PRESS scores severity levels (Figure 1).

Table 2. Characteristics of the patients.

Characteristics		All patients n = 120	%
Age (months)		37.2 \pm 35.9	
Gender			
	Male	61	50.8
	Female	59	49.2
Acute respiratory tract infection			
	Upper	33	27.5
	Lower	87	72.5
Treatment			
- Hospitalization	General ward	82	68.3
	ICU admission	4	3.3
	Total	86	71.6
- No admission		34	28.3

ICU: intensive care unit




Figure 1. Numbers of admission classified according to their severity.

The statistical results of admitted patient treatments show the area under curve of admitted patient treatments is 0.97 (95%CI: 0.95 - 0.99) with a cutoff point of 1.5 (scores >1.5 predicting admission), giving a sensitivity of 0.94 and a specificity of 0.88 (Figure 2). As for ICU patients, the area under curve of this group is 0.81 (95%CI: 0.64 - 0.98) with a cutoff point of 2.5 (scores >2.5 predicting ICU admission), giving a sensitivity of 1.00 and a specificity of 0.46, and with a cutoff point of 3.5 (scores >3.5 predicting ICU admission), giving a sensitivity of 0.75 and a specificity of 0.66 (Figure 2).

The duration of oxygen therapy results classified from the severity are 0.23 ± 0.73 days (mean \pm SD) for mild, 1.60 ± 1.69 days for moderate, and 3.00 ± 2.83 days for severe. This shows a statistically significant difference (Figure 3). Moreover, the nebulized bronchodilator results also show significant differences between each group: 0.66 ± 1.63 days for mild, 3.79 ± 2.56 days for moderate, and 5.64 ± 3.16 days for the severe group (Figure 4). Of note, no mortality occurred from this research.




Figure 2. Receiver operating characteristic curve of hospitalization (A), and ICU admission (B).




Figure 3. Duration of oxygen therapy results classified from the severity.




Figure 4. Duration of nebulized bronchodilator results classified from the severity.

Discussion

In community hospitals, there are several child patients with fever, cough, and dyspnea. Some patients have a respiratory failure condition that requires an immediate treatment and an urgent referral. Some other patients may not be excessively tired, but still need to be treated with oxygen therapy, a nebulized bronchodilator, and admission. As for those with less severe cases, a simple nebulizer treatment without any admission is sufficient.

PRESS scores are a severity assessment to categorize patients with respiratory infections into 3 groups: mild (0 - 1), moderate (2 - 3), and severe (4 - 5) based on five parameters: respiratory rate, wheezing, accessory muscle use, SpO₂, and feeding difficulties.

According to this study, the admitted regular patients in the moderate, and severe group results have a sensitivity of 0.94 and a specificity of 0.88, statistically significant, whilst the ICU patients in the severe group had a sensitivity of 0.75 and a specificity of 0.66, also statistically significant. This shows a relationship between PRESS scores and the predictability of admission to a regular ward, and ICU.

If the cutoff point is at >2.5, which includes both moderate (score 3), and severe (score 4 - 5) groups, the sensitivity and the specificity become 1.00, and 0.46, respectively. It may yield an over treatment condition for ICU admissions that covers all severe condition patients, which is suitable for a hospital with sufficient ICU capacity. However, if the cutoff point is at >3.5, which includes only a severe (score 4 - 5) group, it yields 0.75 sensitivity, and 0.66 specificity. That excludes 25% of severe condition patients. This is suitable for a hospital that has lower ICU capacity.

Additionally, severe patients required statistically significantly longer oxygen therapy, and a nebulized bronchodilator more than the mild, and moderate patients. This is similar to a PRESS score study of the Department of Pediatrics, National Hospital of Yokohama, which showed that the rates of hospitalization for moderate, and severe patients were higher than the mild cases, and the severe group required a longer duration of oxygen treatment than others. Furthermore, this system is more reliable than the respiratory distress assessment Instrument (RDAI), and the Children's Hospital of Wisconsin Respiratory Score (CHWRS). The area under the curve for the CHWRS was 0.68 with a cutoff point of 7.5 (scores >7.5 predicting admission), giving a sensitivity of 0.65 and a specificity of 0.65. The area under the curve for the RDAI was 0.51.⁽¹¹⁾

As there is no guidelines of treatment for participant doctors, it raises concern about the inconsistencies of receiving treatment among patients that could be effected following such as rate of general ward or PICU admission, times to start and finish oxygen therapy and nebulized bronchodilators. Hence this may reduce the reliability of these results. The recommendation outlook is to regulate a treatment guideline to improve reliability of these results.

Conclusion

In addition to its simplicity and ease of use, PRESS scores can predict the severity of acute respiratory infections and could guide a proper treatment for pediatric patients.

Conflict of interest

None of the authors has any potential conflict of interest to disclose.

References

1. Prapphal N, Limudomporn S, Chumdermpadetsuk S. Risk factors of acute respiratory infection in Thai children. *Chula Med J* 1993;37:13-20.
2. Prapphal N. Respiratory infection and allergy in children. *Chula Med J* 2002;46:459-60.
3. Simoes EAF, Cherian T, Chow J, ShahidSalles S, Laxminarayan R, John TJ. Acute respiratory infections in children. In: Jamison DT, Breman JG, Measham AR, Alleyne G, Claeson M, Evans DB, et al, editors. *Source Disease Control Priorities in Developing Countries*. 2nd edition. Washington (DC): World Bank; 2006. Chapter 25.
4. Suwanjutha S, Sunakorn P, Chantarojanasiri T, Siritantikorn S, Nawanoparatkul S, Rattanadilok Na Bhuket T, et al. Respiratory syncytial virus associated lower respiratory tract infection in under-5-year-old children in a rural community of central Thailand, a population-based study. *J Med Assoc Thai* 2002; 85 Suppl 4:S1111-9.
5. Kaewnokkhao W. Situation of pneumonia, Thailand, 2005 - 2010. *W Epidemiol Surveil Rep* 2012;43 Suppl: S90-8.
6. Feldman AS, Hartert TV, Gebretsadik T, Carroll KN, Minton PA, Woodward KB, et al. Respiratory severity score separates upper versus lower respiratory tract infections and predicts measures of disease severity. *Pediatr Allergy Immunol Pulmonol* 2015;28:117-20.
7. Ducharme FM, Chalut D, Plotnick L, Savdie C, Kudirka D, Zhang X, et al. The pediatric respiratory assessment measure : a valid clinical score for assessing acute asthma severity from toddlers to teenagers. *J Pediatr* 2008;152:476-80.
8. Gold DL, Mihalov LK, Cohen DM. Evaluating the Pediatric Early Warning Score (PEWS) system for admitted patients in the pediatric emergency department. *Acad Emerg Med* 2014;21:1249-56.
9. Beyeng RTD, Purniti PS, Naning R. Validity of bacterial pneumonia score for predicting bacteremia in children with pneumonia. *Paediatr Indones* 2011;2: 322-6.
10. Miyaji Y, Sugai K, Nozawa A, Kobayashi M, Niwa S, Tsukagoshi H. Pediatric Respiratory Severity Score (PRESS) for Respiratory Tract Infections in Children. *Austin Virol and Retrovirology* 2015;2:1009.
11. Destino L, Weisgerber MC, Soung P, Bakalarski D, Yan K, Rehborg R, et al. Validity of respiratory scores in bronchiolitis. *Hosp Pediatr* 2012;2: 202-9.