

5-1-2010

## Cervical disc dimensions of the Thai population

U Wandee

P Itiravivong

T Tejapongvorachai

P. Tangpornprasert

Follow this and additional works at: <https://digital.car.chula.ac.th/clmjournal>



Part of the [Medicine and Health Sciences Commons](#)

---

### Recommended Citation

Wandee, U; Itiravivong, P; Tejapongvorachai, T; and Tangpornprasert, P. (2010) "Cervical disc dimensions of the Thai population," *Chulalongkorn Medical Journal*: Vol. 54: Iss. 3, Article 5.

DOI: <https://doi.org/10.58837/CHULA.CMJ.54.3.5>

Available at: <https://digital.car.chula.ac.th/clmjournal/vol54/iss3/5>

This Article is brought to you for free and open access by the Chulalongkorn Journal Online (CUJO) at Chula Digital Collections. It has been accepted for inclusion in Chulalongkorn Medical Journal by an authorized editor of Chula Digital Collections. For more information, please contact [ChulaDC@car.chula.ac.th](mailto:ChulaDC@car.chula.ac.th).

## Cervical disc dimensions of the Thai population

Udthapon Wandee\* Pibul Itiravivong \*

Tawechai Tejapongvorachai\* Pairat Tangpornprasert\*\*

**Wandee U, Itiravivong P, Tejapongvorachai T, Tangpornprasert P. Cervical disc dimensions of the Thai population. Chula Med J 2010 May – Jun; 54(3): 225 - 36**

- Problem** : *The incidence of complications in cervical disc arthroplasty was 6.2% per treat level. Perioperative kyphosis, heterotopic ossification and migration may occur be due to undersized prosthesis. Measurement of the dimensions of cervical discs of the Thai population was aimed to be a pilot study to know the estimation of the dimensions of cervical discs that could be referred to the size of disc prosthesis devices.*
- Objective** : *To provide data regarding the dimension of cervical discs dimensions that can be used in Thailand for preoperative planning and design of cervical disc prosthesis in Thailand.*
- Study design** : *Descriptive study.*
- Setting** : *Department of Orthopedics, King Chulalongkorn Memorial Hospital.*
- Method and Material** : *From January 2008 to December 2008, 60 consecutive C-spine magnetic resonance imaging studies were performed on Thai subjects. MRI was performed using a 1.5 Tesla whole body MR imaging system with an extremity coil. Pulse sequences were T2-weighted images. The direction of the axial slice imaging placed the slice perpendicular to the spinal mechanical axis in the coronal plane and perpendicular to the long axis of the spine in the sagittal plane. All 60 images were reconstructed at 3-mm intervals.*

\* Department of Orthopedics Surgery, Faculty of Medicine, Chulalongkorn University

\*\*Department of Engineer, Faculty of Engineer, Chulalongkorn University

- Result** : *Total mean maximum disc height in Thai population are as follows: C3-4 6.44 mm, C4 - 5 5.90 mm, C5 - 6 5.79 mm, C6 - 7 6.28 mm and C7 - T1 6.21 mm. The differences in the maximum disc heights between males and females were significant. The maximum and minimum distances from all samples are 6.63 mm and 5.47 mm, respectively. The total mean sagittal diameter in the Thai population are: C3i 14.77 mm, C4s 14.77 mm, C4i 15.23 mm, C5s 15.13 mm, C5i 15.74 mm, C6s 15.62 mm, C6i 15.72 mm, C7s 15.69 mm, C7i 15.79 mm and T1s 16.00 mm. The maximum and minimum distances from all samples are 16.47 mm and 13.33 mm respectively. The differences in maximum disc height between males and females were significant. The total mean transverse diameters in the Thai population are C3i 21.77 mm, C4s 22.15 mm, C4i 22.15 mm, C5s 22.95 mm, C5i 23.44 mm, C6s 24.23 mm, C6i 25.36 mm, C7s 26.28 mm, C7i 26.92 mm and T1s 26.95 mm. The maximum and minimum distances from all samples are 27.73 mm and 20.80 mm respectively. The differences in the maximum disc height between males and females were significant.*
- Conclusion** : *Measurement of disc dimensions can refer to the ranges of size to manufacturers for the devices to cover the use in Thai patients. The data fulfill the goal described for cervical disc dimension for the Thai population. The results of the project could provide design and data for the manufacturers and next step of experiment for cervical disc prosthesis suitable for the Thai population.*
- Keywords** : *Cervical disc dimensions, Cervical spondylosis, Cervical disc prosthesis.*

Reprint request: Wandee U. Department of Orthopedics Surgery, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. October 20, 2009.

อรรถพล วันดี, พิบูลย์ อธิธีระวิวงศ์, ทวีชัย เตชะพงศ์วรชัย, ไพรัช ตั้งพรประเสริฐ.  
มิติรูปร่างหมอนรองกระดูกของคนไทย. จุฬาลงกรณ์เวชสาร 2553 พ.ศ. - มิ.ย.; 54(3):  
225 - 36

- เหตุผลการทำวิจัย :** การใช้หมอนรองกระดูกเทียมจากต่างประเทศหากมีขนาดไม่เหมาะสมกับกายวิภาคของคนไทย อาจทำให้เกิดปัญหา subsidence และ migration ได้
- วัตถุประสงค์ :** เพื่อเป็นข้อมูลมิติรูปร่างหมอนรองกระดูกของคนไทย เพื่อการวางแผนก่อนการผ่าตัดและเป็นข้อมูลเบื้องต้นในการออกแบบหมอนรองกระดูกเทียมในคนไทยต่อไป
- รูปแบบการทำวิจัย :** การศึกษาเชิงพรรณนา
- สถานที่ทำการวิจัย :** ฝ่ายศัลยกรรมกระดูก โรงพยาบาลจุฬาลงกรณ์
- วิธีการศึกษา :** คนไทยที่มาทำการตรวจคลื่นแม่เหล็กไฟฟ้าที่ระดับคอในช่วงเดือน ม.ค. ถึงเดือน ธ.ค. พ.ศ. 2551 จากการถ่ายภาพด้วยคอมพิวเตอร์สแกนแม่เหล็กใช้ภาพ T2-weight images เป็นภาพตัดตั้งฉากกับแนวแกน mechanical axis ของกระดูกสันหลังในระนาบ coronal และตั้งฉากแนวตามยาวของกระดูกสันหลังในระนาบ sagittal ระยะห่างระหว่างภาพตัดห่างกัน 3 มม. ทั้งหมด 60 ภาพ
- ผลการศึกษา :** ผลของการวัดระยะคือค่าเฉลี่ยความสูงของหมอนรองกระดูกในคนไทย ค่าเฉลี่ยรวมทั้งหมดเป็นดังนี้ C3-4 6.44 mm, C4-5 5.90 mm, C5-6 5.79 mm, C6-7 6.28 mm และ C7-T1 6.21 mm. มีความแตกต่างระหว่างระยะในเพศชายและเพศหญิงอย่างมีนัยสำคัญ ค่าความสูงที่สุดและต่ำที่สุดจาก sample ทั้งหมดคือ 6.63 mm และ 5.47 mm ตามลำดับ ส่วนค่าเฉลี่ยความกว้างในแนวกลางของหมอนรองกระดูกในคนไทยค่าเฉลี่ยรวมทั้งหมดเป็นดังนี้ C3<sub>i</sub> 14.77 mm, C4<sub>i</sub> 14.77 mm, C4<sub>s</sub> 15.23 mm, C5<sub>i</sub> 15.13 mm, C5<sub>s</sub> 15.74 mm, C6<sub>i</sub> 15.62 mm, C6<sub>s</sub> 15.72 mm, C7<sub>i</sub> 15.69 mm, C7<sub>s</sub> 15.79 mm และ T1<sub>s</sub> 16.00 mm. มีความแตกต่างระหว่างระยะในเพศชายและเพศหญิงอย่างมีนัยสำคัญ ค่าความสูงที่สุดและต่ำที่สุดจาก sample ทั้งหมดคือ 16.47 mm และ 13.33 mm ตามลำดับ ส่วนค่าเฉลี่ยความกว้างในแนวขวางของหมอนรองกระดูกในคนไทยค่าเฉลี่ยรวมทั้งหมดเป็นดังนี้ C3<sub>i</sub> 21.77 mm, C4<sub>s</sub> 22.15 mm, C4<sub>i</sub> 22.15 mm, C5<sub>s</sub> 22.95 mm, C5<sub>i</sub> 23.44 mm, C6<sub>s</sub> 24.23 mm, C6<sub>i</sub> 25.36 mm, C7<sub>s</sub> 26.28 mm, C7<sub>i</sub> 26.92 mm และ T1<sub>s</sub> 26.95 mm. มีความแตกต่างระหว่างระยะในเพศชายและเพศหญิงอย่างมีนัยสำคัญ ค่าความสูงที่สุดและต่ำที่สุดจาก sample ทั้งหมดคือ 27.73 mm และ 20.80 mm ตามลำดับ

- สรุป** : มิติรูปร่างหมอนรองกระดูกสันหลังที่วัดได้ทำให้รู้กรอบกว้าง ๆ ของขนาดอุปกรณ์ในการวางแผนการรักษาและทำการผลิตในกรณีจะผลิตออกใช้จริง
- คำสำคัญ** : มิติรูปร่างหมอนรองกระดูก, กระดูกคอเสื่อม, การใส่ส่วยจะเทียม.

Symptoms suggestive of cervical and lumbar stenosis are relatively common among this cohort of older men, and generalized spinal stenosis may occur in as many as 4%.<sup>(1)</sup> The most common age group of the Thai population that needs the prosthesis is 30 - 60 years.<sup>(2)</sup>

Anterior cervical discectomy and fusion (ACDF) of one or two levels for spondylotic myelopathy or radiculopathy has been proved to be an extremely effective procedure in terms of clinical and radiographic outcomes.<sup>(3-5)</sup> The rate of adjacent segment degeneration has been reported to be as high as 3% to 11% per year for the first decade after fusion, with up to two-thirds of patients who required reoperation.<sup>(6,7)</sup> Hypermobility of segments adjacent to a fused segment is also often observed.<sup>(8)</sup> Advantages of cervical disc arthroplasty over anterior cervical discectomy and fusion.<sup>(9)</sup> Recently, cervical arthroplasty trends to be used more than ACDF. The incidences of complications in cervical disc arthroplasty were 6.2% per treat level.<sup>(10)</sup> The complications were, namely:

1. retropharyngeal hematoma;
2. neurological worsening;
3. intraoperative and delay migration;
4. postoperative segmental kyphosis;
5. heterotopic ossification and spontaneous fusion;
6. partial dislocation of the prosthesis in extension; and,
7. neck-and-shoulder pain

Some complications from arthroplasty such as subsidence may be occurred after prosthesis replacements.<sup>(11)</sup> There are 3 factors that were had been studied in cervical cage.<sup>(11)</sup>

1. Distance from anterior vertebral rim;
2. Spacer versus end-plate surface ratio;
3. Ratio of pre- and immediate postoperative height of the inter-vertebral space.

Perioperative kyphosis may occur after removing the anchoring pin and dual-track milling guide because of the mismatch of the size of the prosthesis.<sup>(10)</sup> Heterotopic ossification may occur because of limited motion of the prosthesis due to undersized prosthesis.<sup>(10)</sup> Delayed prosthesis migration was found in patients who had immediate kyphosis after operation.<sup>(10)</sup> In a 2 - year follow up period, a re-operative of 2.05% appears to be acceptable rate considering the investigative nature of this study.<sup>(12)</sup>

However, clinical study is still in need regarding the incomplete knowledge of the dimensions of cervical discs of the Thai population.<sup>(13)</sup> We, therefore, conducted a study of the dimensions cervical discs of the Thai population to prepare the data for preoperative planning and designs of cervical disc prostheses.

## Materials and Methods

### Patients Selection

From January 2008 to December 2008, 60 consecutive C-spine magnetic resonance imaging studies were performed in Thai subjects. The inclusion criteria of each subjects were: the subjects is between 22 to 76 years of age, having no congenital deformity, scoliosis, traumatic injury to the spine and/or having no previous spine surgery. From the criteria, 30 were males and 30 females were recruited. The demographic data are shown in Table 1.

**Table 1.** The basic information of the subjects\*

Sex	Total	Male	Female
Subjects (Number)	60	30	30
Age (year)	51.00 $\pm$ 12 (22 - 76)	52.47 $\pm$ 11.01 (30 - 72)	50.33 $\pm$ 13.13 (22 - 76)
Weight (Kg.)	60.75 $\pm$ 11.80 (40 - 100)	66.90 $\pm$ 10.00 (50 - 100)	54.60 $\pm$ 10.24 (40 - 88)

\*Represented as mean  $\pm$  standard deviation, median (range)

### MRI measurements

MRI was performed using a 1.5 Tesla whole body MR imaging system (Siemens 1.5 Tesla, Avanto, Germany) with an extremity coil. Pulse sequences were T2-weighted images.

The direction of the axial slice imaging placed the slice perpendicular to the spinal mechanical axis in the coronal plane and perpendicular to the long axis of the spine in the sagittal plane.

All 60 images were reconstructed at 3-mm intervals. These images were obtained from patients who attended clinics for neck pain. Patients below the age of 20, those with congenital cervical spine anomaly, having history of past surgery of the cervical spine, pregnancy, and/or any abnormal disc, which was outside the Woodend Classification I (Figure 1.), were all excluded. The Woodend Classification grades of the disc on a scale of 1- 4, where grade 1 is normal disc with a white nucleus, normal shape, and no annular tears.<sup>(14)</sup>

Measurement of the disc dimensions counted on 3-plane slice imaging at the most midline cut of MRI in T2 weighted image. Considering lordosis of the cervical spine, we measure the height of vertebral body at anteriorly, posteriorly, and maximum disc height. The data concerning the endplates were obtained from the caudal surfaces. On each surface

we measured three diameters; the median sagittal diameter (SD) which goes through the middle of the transverse diameter; the transverse diameter (TD) which is the maximum breadth across the vertebral body; the “diagonal diameter” (DD) which we considered the longest diameter running through the intersection of the two others and forming an angle between 30° and 60° with the sagittal plane. If we suspected a slight asymmetry, we would compare the diameter from the right anterior to the left posterior with the one from the left anterior in the right posterior. However, we never found any significant difference. Grossly asymmetrical cases would have been considered pathological and therefore excluded. The data were obtained by a computerized coordinate system from MRI images. All data were analyzed by Student's t test, ANOVA, and correlation between levels was analyzed. The validation of the data was done by intra-observer method and analyzed by Student pair t-test. A total of 300 measured cervical spine levels were done.

## Results and Discussion

### Disc dimensions

Measurement was done in 60 samples (30 males; 30 females) and the results are shown as table 3.









Grade	Disc Changes			Disc Height (H)
	Axial Sections	Sagittal view	Description	
Normal (DC-1)	White (Hypertensive) 		Flat or slightly convex, posterior annulus, nuclear cleft, on sagittal views. Annular margins well defined (bean shaped or rounded), no tears in axial views.	Normal (H1)
Mild (DC-2)	White or speckled 		Flat or bulge of posterior annulus on sagittal views. Distortion of the bean shape, or a rounded appearance of the annulus. Small radial tears not reaching the PLL on the axial views.	Reduced by 10% (H2)
Moderate (DC-3)	Speckled or dark 		Bulge or Prolapse of the posterior annulus on the sagittal views. Ill defined appearance of the annulus in axial views. Radial tears extending upto torn PLL on the sagittal/axial views, +/- prolapse or end plate changes.	Reduced by 10% - 50% (H3)
Severe (DC-4)	Dark (Hypointense) 		No difference between the appearance of annulus and nucleus, +/- complex tears, +/- Prolapse with or without end plate changes.	Reduced by 50% or more (H4)

Figure 1. Woodend Classification.

Table 2. Average anterior cervical disc height.

Subject	Subject number	Average anterior cervical disc height (mm)				
		C3-4	C4-5	C5-6	C6-7	C7-T1
Male	30	3.40	3.40	3.17	3.50	3.63
Female	30	3.03	2.77	2.47	3.03	2.93
Total	60	3.41	3.23	3.05	3.44	3.51

Table 3. Maximum cervical disc heights.

Subject	Subject number	Average maximum cervical disc height (mm)				
		C3-4	C4-5	C5-6	C6-7	C7-T1
Male	30	6.63	6.07	5.87	6.37	6.20
Female	30	5.83	5.53	5.47	5.87	6.10
Total	60	6.44	5.90	5.79	6.28	6.21



The maximum anterior cervical disc height in the male subjects was 3.63 mm at C3 - 4 level and minimum 2.47 mm at C5-6 level in the female subjects. There is statistical significantly difference between the male and the female subjects with was attested by ANOVA. (P-value <0.05). The mean difference is shown to be between male and female. Considering lordosis of the cervical spine, we measured the height of the vertebral body at anteriorly, posteriorly, and maximum disc height. The results are shown as table 3.

Maximum cervical disc height in the male subjects was 6.63 mm at C3-4 level and minimum 5.47 mm at C5-6 level in the female. There are statistical significantly differences between those of men and women attested by ANOVA (P-value <0.05) Regarding the lordosis of the cervical spine, we measured the heights of the vertebral body at

anteriorly, posteriorly, and the maximum disc height. The results are shown as table 4.

The maximum posterior cervical disc height in the male subjects was 3.30 mm at C3-4 level and the minimum of 2.37 mm at C5-6 level in the female. There are statistical significantly difference between men and women, attested by ANOVA (P-value <0.05). Regarding lordosis of the cervical spine, we measured the height of the vertebral body at anteriorly, posteriorly, and the maximum disc height. The results are shown as table 5.

The maximum sagittal diameter in the male subject was 16.47 mm at T1 level and the minimum of 13.33 mm at C3<sub>i</sub> level in the female. There are statistical significantly differences between men and women, attested by ANOVA. (P-value <0.05). The results are shown as table 6.

**Table 4.** Posterior cervical disc heights.

Subject	Subject number	Average posterior cervical disc height (mm)				
		C3-4	C4-5	C5-6	C6-7	C7-T1
Male	30	3.30	2.97	2.97	3.03	3.03
Female	30	2.80	2.63	2.37	2.53	2.67
Total	60	3.23	2.87	2.90	3.00	2.95

**Table 5.** Sagittal diameters.

Subject	Subject number	Average sagittal diameters (mm)									
		C3 <sub>i</sub>	C4 <sub>s</sub>	C4 <sub>i</sub>	C5 <sub>s</sub>	C5 <sub>i</sub>	C6 <sub>s</sub>	C6 <sub>i</sub>	C7 <sub>s</sub>	C7 <sub>i</sub>	T1 <sub>s</sub>
Male	30	15.17	15.1	15.67	15.63	16.17	15.9	16.2	16.07	16.33	16.47
Female	30	13.33	13.37	13.77	13.37	13.93	13.97	14.03	14.13	13.67	14.22
Total	60	14.77	14.77	15.23	15.13	15.74	15.62	15.72	15.69	15.79	16

**Table 6.** Transverse diameters.

Subject	Subject number	Average transverse diameters (mm)									
		C3 <sub>i</sub>	C4 <sub>s</sub>	C4 <sub>i</sub>	C5 <sub>s</sub>	C5 <sub>i</sub>	C6 <sub>s</sub>	C6 <sub>i</sub>	C7 <sub>s</sub>	C7 <sub>i</sub>	T1 <sub>s</sub>
Male	30	22.1	22.3	22.4	23.2	23.73	24.57	25.83	26.6	27.47	27.73
Female	30	20.8	21.5	21.7	22.2	22.83	23.4	24.13	25.43	25.57	24.8
Total	39	21.77	22.15	22.15	22.95	23.44	24.23	25.36	26.28	26.92	26.95

The maximum transverse diameter in the male subjects was 27.73 mm at T1 level and the minimum of 20.80 mm at C3<sub>i</sub> level in female. There are statistical significantly difference between men and women, attested by ANOVA. (P-value <0.05). The results are shown as table 7.

The maximum diagonal diameter in the male subjects was 25.57 mm at C7<sub>s</sub> level and the minimum of 21.37 mm at C3<sub>i</sub> level in female. There are statistical significantly difference between men and women with

test by ANOVA. (P-value <0.05)

The validation of data has been done by intraobserver method and analyzed by Student pair t-test. The results are shown as table 8.

From previous study of cervical disc dimension<sup>(15)</sup> (as table 2), we compared cervical discs dimensions data between sagittal diameter, trasnverse diameter, and diagonal diameter by using Student's t-test. The results are shown as table 9

**Table 7.** Diagonal diameters.

Subject	Subject number	Average diagonal diameters (mm)									
		C3 <sub>i</sub>	C4 <sub>s</sub>	C4 <sub>i</sub>	C5 <sub>s</sub>	C5 <sub>i</sub>	C6 <sub>s</sub>	C6 <sub>i</sub>	C7 <sub>s</sub>	C7 <sub>i</sub>	T1 <sub>s</sub>
Male	30	21.9	22.73	22	23.13	23.2	23.93	24.83	25.57	25.13	25.53
Female	30	21.37	22.4	21.57	22.47	23.3	23.17	23.83	24.5	24.43	23.8
Total	60	21.63	22.57	21.78	22.8	22.75	23.55	24.33	25.03	24.78	24.67

**Table 8.** Validation of intraobserver measurement.

Subject	Mean of 1 <sup>st</sup>	Mean 2 <sup>nd</sup>	P-value
	measurement (mm)	measurement (mm)	
Sagittal diameter	14.47	14.46	0.1
Transverse diameter	25.52	25.14	0.2
Diagonal diameter	22.49	21.73	0.00001
Disc height	5.79	5.95	0.08

**Table 9.** Compare previous study.

Subject	Previous study S. Aharinejad et al.	Cervical disc dimensions	P-value
Transverse diameter (mm)	21.2 $\pm$ 3.7	23.1 $\pm$ 3.6	<0.001
Diagonal diameter (mm)	19.6 $\pm$ 3.0	23.4 $\pm$ 2.3	<0.001
Sagittal diameter (mm)	16.2 $\pm$ 2.1	14.8 $\pm$ 1.9	<0.001

The cervical disc dimensions of Thai population is different from previous study that the sample data were collected from Austria (caucasian). This different dimensions of the cervical discs from caucasian may be inappropriate cervical discs prosthesis that be used in Thailand.

## Conclusion

This project aimed to know the cervical disc dimensions of Thai population and to compare with cervical disc prosthesis that be used in Thailand. As the cervical disc dimensions to use in cervical spine measurement of the cervical disc dimensions was done only in cervical segment. From the measurement result, it was recommended that the cervical disc prosthesis should cover 7 varies sizes (from 3 mm to 9 mm) for disc height, 6 sizes (from 10 mm to 20.50 mm) for sagittal diameter, and 8 sizes (from 17 mm to 31.20 mm) for transverse diameter.

The cervical disc dimensions in this research had been measured by MRI method that it could replace disc morphology in early degenerative motion segment and correct it to near normal. It allowed motion segment to move in flexion and extension.

There are different demographic data between male and female, may be limited to compare between male and female. More size disc may be

available in Thailand. To compare the cervical disc dimension with previous study was significantly different in sagittal diameter, transverse diameter, and diagonal diameter. The differences have been shown that cervical disc dimensions of Thai population seem to be smaller in disc height and transverse diameter than previous study.

Some complications from arthroplasty such as subsidence and migration may be occurred after prosthesis replacements.<sup>(11)</sup> Although, Failure of an intact cervical endplate occurs with an axial load of 634-745 N.<sup>(16,17)</sup> The cervical disc prosthesis subsidence likely stem from multiple causes including osteoporosis, aggressive endplate preparation, postoperative exogenous forces, and bracing.<sup>(4, 18)</sup> The improperly prosthesis disc height, the compressive force increase higher when increasing disc height.<sup>(19)</sup>

If we have a proper size of disc dimension, it may decrease incidence of subsidence or migration. The results of cervical disc dimensions may be useful for preoperative planning and data for cervical disc prosthesis design in the future.

## Acknowledgements

We thank Prachachuen MRI Center for permission to use MRI data. We also thank Mr. Suphap Ngoenthom and Mr. Sripet Chobaroon,

Managing Director of Prachachuen MRI Center and Radiographic of Technologist for their assistance in MRI measurements.

## References

1. Vogt MT, Cawthon PM, Kang JD, Donaldson WF, Cauley JA, Nevitt MC. Prevalence of symptoms of cervical and lumbar stenosis among participants in the Osteoporotic Fractures in Men Study. *Spine* 2006 Jun; 31(13): 1445-51
2. Bunnag S. Cervical spondylosis and the result of conservative treatment. *Bull Dept Med Serv* 1984 Oct; 9(10): 715-8
3. Emery SE, Bolesta MJ, Banks MA, Jones PK. Robinson anterior cervical fusion comparison of the standard and modified techniques. *Spine* 1994 Mar; 19(6): 660-3
4. Bohlman HH, Emery SE, Goodfellow DB, Jones PK. Robinson anterior cervical discectomy and arthrodesis for cervical radiculopathy. Long-term follow-up of one hundred and twenty-two patients. *J Bone Joint Surg Am* 1993 Sep; 75(9): 1298-307
5. Brodke DS, Zdeblick TA. Modified Smith-Robinson procedure for anterior cervical discectomy and fusion. *Spine* 1992 Oct; 17(10 Suppl): S427-30
6. Hilibrand AS, Carlson GD, Palumbo MA, Jones PK, Bohlman HH. Radiculopathy and myelopathy at segments adjacent to the site of a previous anterior cervical arthrodesis. *J Bone Joint Surg Am* 1999 Apr; 81(4): 519-28
7. Goffin J, Geusens E, Vantomme N, Quintens E, Waerzeggers Y, Depreitere B, Van Calenbergh F, van Loon J. Long-term follow-up after interbody fusion of the cervical spine. *J Spinal Disord Tech* 2004 Apr; 17(2): 79-85
8. Bryan VE Jr. Cervical motion segment replacement. *Eur Spine J* 2002 Oct; 11 Suppl 2: S92-7
9. Acosta FL Jr, Ames CP. Cervical disc arthroplasty: general introduction. *Neurosurg Clin N Am* 2005 Oct; 16(4): 603-7
10. Pickett GE, Sekhon LH, Sears WR, Duggal N. Complications with cervical arthroplasty. *J Neurosurg Spine* 2006 Feb; 4(2): 98-105
11. Barsa P, Suchomel P. Factors affecting sagittal malalignment due to cage subsidence in standalone cage assisted anterior cervical fusion. *Eur Spine J* 2007 Sep; 16(9): 1395-400
12. Pimenta L, Diaz R, McAfee CP, Cappuccino A, Cunningham B, Nicola H, Chamby JT, Guarzedin I. Cervical disc replacement revisions: Clinical and biomechanical considerations. In: Yue J, Bertagnoli R, McAfee P, An H, eds. *Motion Preservation Surgery of the Spine*. Philadelphia: Saunders Elsevier, 2008: 287-96
13. Pooni JS, Hukins DW, Harris PF, Hilton RC, Davies KE. Comparison of the structure of human intervertebral discs in the cervical, thoracic and lumbar regions of the spine. *Surg Radiol Anat* 1986; 8(3): 175-82
14. Askar Z, Wardlaw D, Muthukumar T, Smith F, Kader D, Gibson S. Correlation between inter-vertebral disc morphology and the results in patients undergoing Graf ligament stabilisation. *Eur Spine J* 2004 Dec; 13(8): 714-8

15. Aharinejad S, Bertagnoli R, Wicke K, Firbas W, Schneider B. Morphometric analysis of vertebrae and intervertebral discs as a basis of disc replacement. *Am J Anat* 1990 Sep; 189(1): 69-76
16. Truumees E, Demetropoulos CK, Yang KH, Herkowitz HN. Failure of human cervical endplates: a cadaveric experimental model. *Spine* 2003 Oct; 28(19): 2204-8
17. Lim TH, Kwon H, Jeon CH, Kim JG, Sokolowski M, Natarajan R, An HS, Andersson GB. Effect of endplate conditions and bone mineral density on the compressive strength of the graft-endplate interface in anterior cervical spine fusion. *Spine* 2001 Apr; 26(8): 951-6
18. Natarajan RN, Chen BH, An HS, Andersson GB. Anterior cervical fusion: a finite element model study on motion segment stability including the effect of osteoporosis. *Spine* 2000 Apr; 25(8): 955-61
19. Truumees E, Demetropoulos CK, Yang KH, Herkowitz HN. Effects of disc height and distractive forces on graft compression in an anterior cervical corpectomy model. *Spine* 2008 Jun; 33(13): 1438-41